

1-1-1984

# Changes in Spectral Properties of Detached Leaves

C. S. T. Daughtry

L. L. Biehl

Follow this and additional works at: <http://docs.lib.purdue.edu/larstech>

---

Daughtry, C. S. T. and Biehl, L. L., "Changes in Spectral Properties of Detached Leaves" (1984). *LARS Technical Reports*. Paper 64.  
<http://docs.lib.purdue.edu/larstech/64>


This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact [epubs@purdue.edu](mailto:epubs@purdue.edu) for additional information.

LARS Technical Report 0615 84  
NASA-Johnson Space Center  
Contract No. NAS9-16528

# Changes in Spectral Properties of Detached Leaves

by

C.S.T. Daughtry  
L.L. Biehl



Laboratory for Applications of Remote Sensing

West Lafayette, Indiana 47906  
USA



**PURDUE,**  
touching  
tomorrow  
today

### Star Information Form

|  |  |  |                                 |
|--|--|--|---------------------------------|
| 1. Report No.  | 2. Government Accession No                           | 3. Recipient's Catalog No                          |                                 |
| 4. Title and Subtitle<br><br>Changes in Spectral Properties of Detached Leaves   |  | 5. Report Date<br>6/15/84                          | 6. Performing Organization Code |
|  |  | 7. Author(s)<br>C.S.T. Daughtry and L.L. Biehl     |                                 |
| 9. Performing Organization Name and Address<br>Laboratory for Applications of Remote Sensing<br>Purdue University<br>1291 Cumberland Ave.<br>West Lafayette, IN 47906-1399   |  | 8. Performing Organization Report No<br>061584     |                                 |
|  |  | 10. Work Unit No.                                  |                                 |
|  |  | 11. Contract or Grant No.<br>NAS9-16528            |                                 |
| 12. Sponsoring Agency Name and Address<br>NASA Johnson Space Center<br>Houston, TX 77058   |  | 13. Type of Report and Period Covered<br>Technical |                                 |
|  |  | 14. Sponsoring Agency Code                         |                                 |
| 15. Supplementary Notes  |  |  |                                 |
| 16. Abstract<br><p style="margin: 0;">A leaf begins to senesce as soon as it is removed from the plant and changes in metabolic processes and spectral properties are inevitable. If senescence can be delayed for several days without significant changes in spectral properties, then samples of leaves at remote test sites could be prepared and shipped to laboratories which have the equipment to measure spectral properties. The objective of this study was to determine the changes in spectral properties of detached leaves. Leaves from red birch (<u>Betula nigra</u> L.) and red pine (<u>Pinus resinosa</u> Ait.) were immersed for 5 minutes in water or 0.001 M benzylaminopurine (BAP) and stored in plastic bags in the dark at either 5° or 25° C. Total directional-hemispherical reflectance and transmittance of the adaxial surface of birch leaves were measured over the 400 to 1100 nm wavelength region with a spectroradiometer and integrating sphere (LICOR LI-1800). Pine needles were taped together and reflectance of the mat of needles was measured. Spectral properties changed less than 5% of initial values during the first week when leaves were stored at 5° C. Storage at 25° C promoted rapid senescence and large changes in spectral properties. BAP delayed, but did not stop, senescence at 25° C. Low temperature was more effective than BAP in delaying senescence. It appears possible to store leaves at low temperatures in plastic bags in the dark for several days without significantly altering the spectral properties in the 400 to 1100 nm wavelength region.</p> |  |  |                                 |
| 17. Key Words (Suggested by Author(s))<br>Reflectance, Transmittance,<br>Red Birch, <u>Betula nigra</u> L.<br>Red Pine, <u>Pinus resinosa</u> Ait.   |  | 18. Distribution Statement                         |                                 |
| 19. Security Classif. (of this report)<br>Unclassified   | 20. Security Classif. (of this page)<br>Unclassified | 21. No. of Pages                                   | 22. Price                       |

NAS9-16528  
LARS 061584

Changes in Spectral Properties of Detached Leaves

C.S.T. Daughtry  
Research Agronomist, LARS

L.L. Biehl  
Research Engineer, LARS

Laboratory for Applications of Remote Sensing  
Purdue University  
West Lafayette, IN 47907-1399, U.S.A.

June 1984

## CHANGES IN SPECTRAL PROPERTIES OF DETACHED LEAVES

### ABSTRACT

A leaf begins to senesce as soon as it is removed from the plant and changes in metabolic processes and spectral properties are inevitable. If senescence can be delayed for several days without significant changes in spectral properties, then samples of leaves at remote test sites could be prepared and shipped to laboratories which have the equipment to measure spectral properties. The objective of this study was to determine the changes in spectral properties of detached leaves. Leaves from red birch (Betula nigra L.) and red pine (Pinus resinosa Ait.) were immersed for 5 minutes in water or 0.001 M benzylaminopurine (BAP) and stored in plastic bags in the dark at either 5° or 25° C. Total directional-hemispherical reflectance and transmittance of the adaxial surface of birch leaves were measured over the 400 to 1100 nm wavelength region with a spectroradiometer and integrating sphere (LICOR LI-1800). Pine needles were taped together and reflectance of the mat of needles was measured. Spectral properties changed less than 5% of initial values during the first week when leaves were stored at 5° C. Storage at 25° C promoted rapid senescence and large changes in spectral properties. BAP delayed, but did not stop, senescence at 25° C. Low temperature was more effective than BAP in delaying senescence. It appears possible to store leaves at low temperatures in plastic bags in the dark for several days without significantly altering the spectral properties in the 400 to 1100 nm wavelength region.

## INTRODUCTION

As models have been developed which describe the interactions of solar radiation with leaves, there has been increased need for accurate and reliable measurements of the spectral properties of leaves. Many factors affect the spectral properties of leaves, including chlorophyll content (Gausman et al., 1973; Hoffer and Johannsen, 1969; Knipling, 1970), water content (Sinclair et al., 1971; Thomas et al., 1971), leaf age (Gausman et al., 1971), and internal structure (Sinclair et al., 1971; Gausman et al., 1969). To describe variability of spectral properties that occur in nature, one must not only select representative samples but at the same time avoid introducing extraneous variation due to the measurement process.

When the spectroradiometer and its accessory equipment are located near the test site, spectral data of leaves can be acquired either in situ with leaves attached to the plant or immediately after removing the leaf from the plant. If precautions to minimize water loss are followed, then changes in spectral properties of the leaves probably will be small during the first several hours after excision (Sinclair et al., 1971; Gausman et al., 1969). However, when the spectroradiometer cannot be located near the test site, several days may elapse from the time leaves are removed from the plant until the spectral properties can be measured. During this time, significant metabolic, and possibly spectral, changes can occur. The objective of this study was to determine the changes in the spectral properties of detached leaves treated with a cytokinin and stored at different temperatures.

In oat leaves (Avena sativa L.), proteolysis may begin within 6 hours and chlorophyll degradation may be evident within 24 hours after excision (Tetley and Thimann, 1974; Martin and Thimann, 1972). In wheat leaves (Triticum aestivum L.) detached for 3 to 4 days, the cytoplasm, tonoplast, and cellular organelles were in various stages of degradation (Hurkman, 1979). Starch deposits in chloroplasts disappeared and the ultrastructure of chloroplasts, mitochondria, and other cellular organelles were swollen and distended, indicating loss of energy (Hurkman, 1979; Thimann, 1980).

Senescence is an active metabolic process and can be affected by temperature, light, plant hormones, and various other compounds (Thimann, 1980). As temperature of excised oat leaves kept in the dark for 3 days increased from 5° to 30° C, chlorophyll concentration decreased rapidly (Tetley and Thimann, 1974). Senescence proceeded at its maximum rate at 30° C. At higher temperatures, the rate of chlorophyll degradation decreased until temperatures exceeded 40° C and thermal effects began to cause secondary yellowing (Tetley and Thimann, 1974; Thimann, 1980).

Light delays both proteolysis and chlorophyll degradation, but the effect of light is readily reversible in the dark (Tetley and Thimann, 1974). Rapid senescence in dark may be due to deprivation of energy (Hurkman, 1979) or to accumulation of abscisic acid (Thimann, 1980). Photoperiod also influences the rate of senescence. For example, leaves of Hibiscus rosa-sinensis which senesce in 20 days in dark, survive to 35 days in 8-hour photoperiods and 45 days in 16-hour photoperiods (Misra and Biswal, 1973).

Cytokinins delay proteolysis and chlorophyll loss in leaves in the dark, perhaps by maintenance of a tight coupling between respiration and phosphorylation (Thimann, 1980). Although the relative activities of the cytokinins vary with plant species, benzylaminopurine (BAP) is generally several times more active than kinetin or zeatin (Tetley and Thimann, 1974). Treatment with BAP as preharvest spray or postharvest dip has been effective in prolonging storage life of some vegetables (Dedolph et al., 1962) and cut flowers (Heide and Oydvin, 1969).

When leaves are detached, they begin to senesce and changes in metabolic processes and physiological and spectral properties are inevitable. However, the rate of senescence can be altered. If senescence can be delayed for several days without significant changes in spectral properties, then samples of leaves from plants at remote test sites can be prepared and shipped to laboratories which have the equipment to measure spectral properties.

## MATERIALS AND METHODS

### Experimental Conditions

Two experiments were conducted during 1983 using leaves from red birch (Betula nigra L.) and red pine (Pinus resinosa Ait.). In the first experiment, 0.5 meter apical segments of branches with leaves fully exposed to the sun on the south side of the trees were cut within 3 hours of sunrise and the bases of the branches were immediately placed in water. Within an hour, four whole, fully-expanded leaves per branch were randomly selected, placed in resealable polyethylene bags, and stored in the dark at  $5^{\circ} \pm 2^{\circ}$  C. In the second experiment, leaves were collected and prepared as in the first experiment, except that the leaves were immersed for 5 minutes in either distilled water or 0.001 M BAP (Heide and Oydvin, 1969). Afterwards the leaves were blotted dry with paper towels, placed in resealable polyethylene bags and stored in the dark at either  $5^{\circ} \pm 2^{\circ}$  C or  $25^{\circ} \pm 3^{\circ}$  C. On selected days after excision, the leaves were removed from storage and their spectral properties were measured.

### Spectral Data

Total (diffuse and specular) directional-hemispherical reflectance and transmittance of the adaxial surface of the birch leaves were measured over the 400 to 1100 nm wavelength region with a spectroradiometer and integrating sphere (LICOR LI-1800). The incident angle of illumination was 5 from normal. An area between the major veins of each birch leaf was identified and measured each time to minimize within-leaf variation in spectral properties. Fascicles of red pine needles were laid next to each other and secured together with adhesive tape at both ends of the needles. An area in the middle of each set of needles was identified and only reflectance of the needles was measured. Spectral properties of the leaves were measured 0, 1, 3, 5, 8, 11, and 25 days after excision in the first experiment and 0, 3, 9, 17, and 23 days after excision in the second experiment. All data were corrected for reflectance of the BaSO<sub>4</sub> reference surface in the integrating sphere and expressed in directional-hemispherical reflectance and transmittance factor units. Absorptance was calculated as 1.0 minus the sum of reflectance and transmittance. Changes in spectral properties were calculated as differences and ratios and plotted as functions of wavelength and time after excision.



## RESULTS AND DISCUSSION

In the visible spectral region (400 to 700 nm), the low reflectance and transmittance and high absorptance (Figs. 1 and 2) of radiation by healthy, green leaves are due to pigments, primarily chlorophylls, carotenoids, xanthophylls, and anthocyanins (Hoffer and Johannsen, 1969; Knipling, 1970). Reflectance and transmittance of leaves in near infrared (700 to 1100 nm) are high and are related to internal cell structure of leaves. An important parameter in determining the level of near infrared reflectance is the number or total area of air-wall interfaces (i.e., discontinuities in refractive index) and not the volume of air space (Sinclair et al., 1971). Factors that change the number of interfaces also change the light-scattering properties of the leaves.

During initial stages of senescence, internal leaf volume may decrease but the number of interfaces may increase as cell walls dehydrate, split, and reorient themselves (Sinclair et al., 1971). These changes increase the radiation diffusing capacity and thus increase near infrared reflectance of leaves. In advanced stages of senescence, breakdown and deterioration of leaf tissue eventually decrease infrared reflectance (Colwell, 1956; Knipling, 1970).

In these birch leaves, no significant changes in reflectance, transmittance, or absorptance (Fig. 1) were evident within 3 days of excision, regardless of the treatment. Leaves stored longer than 3 days at 25° C showed marked changes in reflectance and transmittance, especially in the green (550 nm), red (670 nm), and near infrared (700 and 1100 nm) wavelengths. Reflectances of the pine needles changed only slightly during the 23 days (Fig. 2).

Changes in spectral properties of leaves are easily identified when the spectral property (reflectance, transmittance, or absorptance) on day  $n$  is divided by the spectral property on day 0. These ratios minimize differences in spectral properties among leaves and accentuate the changes in spectral properties within leaves. For birch leaves stored at 5° C, reflectance (Fig. 3A), transmittance (Fig. 4A), and absorptance (not shown) at all wavelengths did not change more than 10% of the value on day 0 until 9 days after the leaves were excised from the trees. After 17 days at 5° C, changes in visible reflectance (Fig. 3A) exceeded 20% of initial values.

For birch leaves stored at 25° C, changes in reflectance (Fig. 3B), and transmittance (Fig. 4B), exceeded 10% of initial values after 3 days. The greatest changes in reflectance (Fig. 3B) after 3 days occurred in green (550 nm), chlorophyll absorption (670 nm), and near infrared wavelengths. Leaves stored at 25° C for more than 9 days transmitted almost no visible (400 to 700 nm) radiation (Figs. 1 and 4B). Transmittance of near infrared (700 to 1100 nm) radiation was significantly reduced also (Fig. 4B) after 3 days at 25° C. Absorp-




Figure 1. Reflectance, transmittance, and absorptance of birch leaves, stored at 25° C, as a function of wavelength and time after excision. Data are means of three leaves. Absorptance may be calculated as 100 minus the sum of reflectance and transmittance.




Figure 2. Reflectance of pine needles, stored at 25° C, as a function of wavelength and time after excision. Data are means of three mats of needles.




Figure 3. Changes in reflectance of birch leaves as a function of wavelength and time after excision. Data are ratios of reflectance on day  $n$  divided by reflectance on day 0.




Figure 4. Changes in transmittance of birch leaves as a function of wavelength and time after excision. Data are ratios of reflectance on day  $n$  divided by reflectance on day 0.

tance of radiation by leaves stored at 25° C increased in the green and near infrared wavelengths (Fig. 1) as the leaves senesced and changed from a bright green to a dull olive color. Changes in optical properties in the visible wavelengths indicate significant chlorophyll degradation while the changes in near infrared (700 to 1100 nm) indicate alteration of cellular structure.

For the pine needles stored at 5° C, the changes in reflectance did not exceed 10% of value during the 23 days of the experiment (Fig. 5A). Changes were greater in the visible than in the near infrared wavelength (Figs. 5A and B). Even the changes in reflectance of the pine needles stored at 25° C (Fig. 5B) were less than the changes in the birch leaves stored at 5° C (Fig. 3A).

The effects of these treatments on the spectral properties were assessed by analysis of variance. Tables 1 and 2 illustrate the significance of F-ratio for main effects and interactions at selected wavelengths. Days after excision and storage temperatures were major factors affecting reflectance, transmittance, and absorptance of birch leaves (Table 1) and reflectance of pine needles (Table 2). The main effect of soaking solution (i.e., water or BAP) on reflectance, transmittance, and absorptance of birch leaves was significant only in the red or chlorophyll absorption wavelengths. For pine needles, the main effect of soaking solution was generally significant in both the visible and near infrared wavelengths. BAP tended to minimize changes in leaf spectral properties by delaying senescence, (Figs. 6 to 10) presumably by slowing chlorophyll loss (Tetley and Thimann, 1974).

When the changes in reflectance, transmittance, and absorptance of these birch leaves were examined at four representative wavelengths, the effects of temperature on rate of senescence are evident (Figs. 6 to 9). Changes in the spectral properties were significantly smaller for leaves stored at 5° C than at 25° C. Tetley and Thimann (1974) reported that senescence, as measured by chlorophyll degradation or  $\alpha$ -amino nitrogen accumulation, progressed very slowly at 5° C but proceeded at its maximum rate at 26 to 30° C. Thus, for leaves stored at 5° C, the increases in reflectance and transmittance and decreases in absorptance at 670 nm (Fig. 7) indicate a gradual loss of chlorophyll.

Leaves stored at 25° C senesced rapidly and the increases in reflectance and transmittance at 670 nm (Fig. 7) at 9 days were greater than those of leaves stored at 5° C for 23 days. However, after 9 days, transmittance of leaves at 25° C was nil (Fig. 1).

Greater changes in near infrared reflectance, transmittance, and absorptance occurred in birch leaves stored at 25° C than at 5° C (Figs. 8 and 9). Radiation scattering in the near infrared (700 to 1300 nm) wavelengths is strongly related to internal leaf structure, particularly the number of cell wall interfaces (Sinclair et al.,



Table 2. Significance of F-ratio in analysis of variance for days after excision (D), storage temperature (T), soaking solution (S), and their interaction. Dependent variables were changes in reflectance at selected wavelengths for pine needles. (n = 60)

| Wavelength<br>nm | Treatment |    |    |    |    | RMSE <sup>†</sup> | CV <sup>§</sup><br>% |
|------------------|-----------|----|----|----|----|-------------------|----------------------|
|                  | D         | T  | S  | DT | TS |                   |                      |
| 450              | *         | ** | ns | ns | *  | 0.043             | 4.3                  |
| 500              | **        | ** | ns | *  | *  | 0.042             | 4.1                  |
| 550              | **        | ** | ** | ns | ** | 0.031             | 3.0                  |
| 600              | **        | ** | *  | ns | ** | 0.036             | 3.6                  |
| 650              | **        | ** | *  | *  | ** | 0.041             | 4.1                  |
| 670              | **        | ** | ns | ns | ns | 0.046             | 4.5                  |
| 700              | **        | ** | ** | *  | *  | 0.032             | 3.1                  |
| 750              | **        | *  | ** | ns | ns | 0.027             | 2.7                  |
| 800              | *         | ns | *  | ns | ns | 0.028             | 2.7                  |
| 850              | *         | ns | *  | ns | ns | 0.028             | 2.8                  |
| 900              | *         | ns | *  | ns | ns | 0.028             | 2.8                  |
| 950              | *         | ns | *  | ns | ns | 0.028             | 2.7                  |
| 1000             | *         | *  | *  | ns | ns | 0.027             | 2.7                  |
| 1050             | *         | ns | *  | ns | ns | 0.028             | 2.8                  |
| 1100             | ns        | *  | *  | ns | ns | 0.031             | 3.0                  |

\*,\*\* = significant at  $\alpha = 0.05$  and  $0.01$ , respectively.

<sup>†</sup> Root mean square error includes DS and DTS interactions which were not significant at  $\alpha = 0.25$ .

<sup>§</sup> Coefficient of variation is RMSE divided by the mean.






Figure 5. Changes in reflectance of pine needles as a function of wavelength and time after excision. Data are ratios of reflectance on day  $n$  divided by reflectance on day 0.




Figure 6. Changes in reflectance (A), transmittance (B), and absorbance (C) of birch leaves at 550 nm as a function of time after excision. Solid lines represent leaves soaked in water before storage and dashed lines represent leaves soaked in BAP. Leaves were stored at either 5° C (squares) or 25° C (triangles).




Figure 7. Changes in reflectance (A), transmittance (B), and absorbance (C) of birch leaves at 670 nm as a function of time after excision. Solid lines represent leaves soaked in water before storage and dashed lines represent leaves soaked in BAP. Leaves were stored at either 5° C (squares) or 25° C (triangles).




Figure 8. Changes in reflectance (A), transmittance (B), and absorbance (C) of birch leaves at 750 nm as a function of time after excision. Solid lines represent leaves soaked in water before storage and dashed lines represent leaves soaked in BAP. Leaves were stored at either 5° C (squares) or 25° C (triangles).

1971, 1973; Gausman et al., 1969). This pattern of changes in spectral properties is consistent with other metabolic and structural changes that occur in senescing leaves. Among the first evidences of cellular degradation is the onset of chloroplast breakdown (Hurkman, 1979). Ultrastructural changes in chloroplast during senescence include increases in size of osmiophilic globuli, changes in organization of internal membranes, swelling of chloroplasts, and loss of stroma (Hurkman, 1979); Mittelheuser and Steveninck, 1971). After the tonoplast disintegrates, other cell constituents degenerate rapidly, a process attributed to release of lytic material from the vacuole (Matile and Winkenbach, 1971). The rapid changes in spectral properties of leaves stored at 25° C probably occurred during the terminal stages of senescence as cell membranes disintegrated.

The changes in reflectance at 670 nm were small for pine needles, averaging less than 5% for needles stored at 5° C and less than 10% for needles stored at 25° C (Fig. 10A). BAP tended to minimize the changes in reflectance at both temperatures. The arrangement of the needles in the mats shifted slightly with handling and probably contributed to measurement errors.

The changes in reflectance of the pine needles were smaller in the near infrared (Fig. 10B) than in the visible (Fig. 10A), indicating that cellular or structural changes were not occurring as rapidly as in the birch leaves (Fig. 8). Needles of pine trees typically live 3 to 5 years (Harlow and Harrar, 1969) and thus must survive much more severe environmental conditions than leaves of deciduous trees such as birch.

In summary, when leaves were stored at low temperatures (5° C) in the dark for one week, the changes in optical properties were less than 5% of the values on day 0. Storage at higher temperatures (i.e., 25° C.) promoted rapid senescence and large changes in optical properties. A prestorage treatment of the leaves with benzylaminopurine (BAP), a cytokinin, slowed but could not stop senescence at 25° C. Low temperature was more effective than BAP in delaying senescence. Thus it appears possible to store leaves at low temperatures in plastic bags for several days without significantly altering the optical properties in the 400 to 1100 nm region. Because of limitations of the spectroradiometer, we could not examine the 1300 to 2400 nm region where optical properties of leaves are dominated by water absorption. However, spectral properties of turgid leaves may not be significantly affected until water content decreases to less than 70% (Sinclair et al., 1971; Gausman et al., 1971; Knipling, 1970).




Figure 9. Changes in reflectance (A), transmittance (B), and absorbance (C) of birch leaves at 1050 nm as a function of time after excision. Solid lines represent leaves soaked in water before storage and dashed lines represent leaves soaked in BAP. Leaves were stored at either 5° C (squares) or 25° C (triangles).




Figure 10. Changes in reflectance of pine needles at 670 nm (A) and 750 nm (B) as a function of time after excision. Solid lines represent needles soaked in water prior to storage and dashed lines represent needles soaked in BAP. Needles were stored at either 5° C (squares) or 25° C (triangles).

## REFERENCES

- Colwell, R.N. (1956), Determining the prevalence of certain cereal crop diseases by means of aerial photography, *Hilgardia* 26:223-286.
- Dedolph, B.R., Wittwer, S.H., Tuli, V., and Gilbert, D. (1962), Effect of N<sup>6</sup>-benzylaminopurine on respiration and storage behavior of broccoli (*Brassica oleracea* var. *Italica*), *Plant Physiol.*, Lancaster 37:509-12.
- Gausman, H.W., Allen, W.A., and Cardenas, R. (1969), Reflectance of cotton leaves and their structure, *Remote Sens. Environ.* 1:19-22.
- Gausman, H.W., Allen, W.A., Cardenas, R., and Richardson, A.J. (1973), Reflectance discrimination of cotton and corn at four growth stages, *Agron. J.* 65:194-198.
- Gausman, H.W., Allen, W.A., Escobar, D.E., Rodriguez, Z.Z., and Cardenas, R. (1971), Age effects of cotton leaves on light reflectance, transmittance, and absorptance and on water content and thickness, *Agron. J.* 63:465-469.
- Harlow, W.M., and Harrar, E.S. (1969), Textbook of Dendrology, McGraw-Hill, New York, NY
- Heide, D.M., and Oydvin, J. (1969), Effects of 6-benzylaminopurine on the keeping quality and respiration of glasshouse carnations, *Hort. Res.* 9:26-36.
- Hoffer, R.M., and Johannsen, C.J. (1969), Ecological potentials in spectral signature analysis, pp. 1-16. In Johnson, Philip (ed.), Remote Sensing in Ecology (Philip Johnson, ed.), University of Georgia Press, Athens, GA.
- Hurkman, W.J. (1979), Ultrastructural changes of chloroplasts in attached and detached, aging primary wheat leaves, *Amer. J. Botany* 66:64-70.
- Knipling, E.B. (1970), Physical and physiological basis for the reflectance of visible and near-infrared radiation from vegetation, *Remote Sens. Environ.* 1:155-159.
- Martin, C., and Thimann, K.V. (1972), The role of protein synthesis in the senescence of leaves. I. The formation of protease, *Plant Physiol.* 49:64-71.
- Misra, G., and Biswal, U.C. (1972), Factors concerned in leaf senescence. I. Effects of age, chemicals, petiole, and photoperiod on



- senescence in detached leaves of Hibiscus rosa-sinesis L., Bot. Gaz. (Chicago) 138:5-11.
- Sinclair, T.R., Hoffer, R.M., and Schreiber, M.M. (1971), Reflectance and internal structure of leaves from several crops during a growing season, Agron. J. 63:864-868.
- Sinclair, T.R., Schreiber, M.M., and Hoffer, R.M. (1973), Diffuse reflectance hypothesis for the pathway of solar radiation through leaves, Agron. J. 65:276-283.
- Tetley, R.M., and Thimann, K.V. (1974), The metabolism of oat leaves during senescence. I. Respiration, carbohydrate metabolism, and the action of cytokinins, Plant Physiol. 54:294-303.
- Thimann, K.V. (1980), The senescence of leaves. In Senescence in Plants (K.V. Thimann, Ed.), CRC Press, Boca Raton, FL. 33432, pp. 86-115.
- Thomas, J.R., Namken, L.N., Oerther, G.F., and Brown, R.G. (1971), Estimating leaf water content by reflectance measurements, Agron. J. 63:845-847.