

1-1-1975

Delineation of Udolls from Udalfs Using Multi-Spectral Scanner Data

J. B. Peterson

F. E. Goodrick

W. N. Melhorn

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Peterson, J. B.; Goodrick, F. E.; and Melhorn, W. N., "Delineation of Udolls from Udalfs Using Multi-Spectral Scanner Data" (1975).
LARS Symposia. Paper 79.
http://docs.lib.purdue.edu/lars_symp/79

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

Symposium on

Machine Processing of

Remotely Sensed Data

June 3 - 5, 1975

The Laboratory for Applications of
Remote Sensing

Purdue University
West Lafayette
Indiana

IEEE Catalog No.
75CH1009-0 -C

Copyright © 1975 IEEE
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

THE USE OF THE ANALYSIS OF COVARIANCE
MODEL IN SPATIAL POINT ESTIMATION

Randall W. Thomas

Remote Sensing Research Program
University of California, Berkeley

DELINEATION OF UDOLLS FROM UDALFS
USING MULTI-SPECTRAL SCANNER DATA

J. B. Peterson, F. E. Goodrick
W. N. Melhorn

LARS
1220 Potter Drive
West Lafayette, IN

In many resource management oriented disciplines, a spatial array of point estimates for ecosystem parameters and productive capacities is required over a given political/administrative unit. Statistical statements of confidence about these point estimates must be made and the relative importance of factors involved in their prediction should be testable. Moreover, the point estimates must be generated in such a way that differences between values over space can be assessed for statistical significance.

Current remote sensing data automatic classification algorithms employ weighted probability density comparisons of clustered/unclustered hypervolume look up functions for class selection. As such, these discriminant models do not produce true point estimates or estimates that may be readily subjected to the statistical analysis necessary in some resource management disciplines.

The use of another discriminant model documented in the statistical literature and giving point estimate arrays with the aforementioned desirable properties is presented. This function is commonly referred to as the analysis of covariance model.

The model may either be developed from raw, spectral/resource parameter channel data obtained from "training" fields or from previously classified data. It may then be used to generate point estimates of such quantities as forage/timber yield and snow water content.

REFERENCE

- Scheffe, H. 1959. The Analysis of Variance. John Wiley & Sons, Inc., New York. 477 pp.