

1-1-1976

A Remote Sensing-Aided System for Evaporation and Watershed-Wide Evapotranspiration Estimation

Siamak Khorram

Randall W. Thomas

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Khorram, Siamak and Thomas, Randall W., "A Remote Sensing-Aided System for Evaporation and Watershed-Wide Evapotranspiration Estimation" (1976). *LARS Symposia*. Paper 112.
http://docs.lib.purdue.edu/lars_symp/112

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

Symposium on

Machine Processing of

Remotely Sensed Data

June 29 - July 1, 1976

The Laboratory for Applications of
Remote Sensing

Purdue University
West Lafayette
Indiana

IEEE Catalog No.
76CH1103-1 MPRSD

Copyright © 1976 IEEE
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

A REMOTE SENSING-AIDED SYSTEM FOR
EVAPORATION AND WATERSHED-WIDE
EVAPOTRANSPIRATION ESTIMATION

Siamak Khorram and Randall W. Thomas
Space Sciences Laboratory
Berkeley, California

ABSTRACT

Timely, spatial, and relatively inexpensive remotely sensed data when combined with other conventional ground meteorological data can potentially provide accurate, location specific estimates of evaporation and evapotranspiration. Reliable evaporation data are required for planning, designing and operating reservoirs, ponds, shipping canals, and irrigation and drainage systems. Data on evapotranspiration are useful for estimating irrigation requirements, rainfall disposition, yield of ground water basins, water yields from mountain watershed, streamflow depletion in river basins, and in crop production management.

The approach uses a multistage, multi-phase sampling technique employing three increasingly involved levels of information. The input variables for level I models are surface temperature, radiation components, relative humidity, and cloud cover; the data came from Landsat, meteorological satellites, meteorological stations, and topographic data. The data for level II models are temperature, humidity, and wind profiles in the soil-vegetation zone along with wind friction velocity, radiation, rainfall, and ground cover. At level III, evapotranspiration models employing a combination of empirical, energy balance, and aerodynamic methods will be applied.