

1-1-1976

Digital Image Processing of LANDSAT I MSS Data Specifically Designed for Linear Enhancement in Southwestern Jordan I MSS Data Specifically Designed for Linear Enhancement in Southwestern Jordan

Pat S. Chavez

Lennis G. Berlin

Alex V. Acosta

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Chavez, Pat S.; Berlin, Lennis G.; and Acosta, Alex V., "Digital Image Processing of LANDSAT I MSS Data Specifically Designed for Linear Enhancement in Southwestern Jordan I MSS Data Specifically Designed for Linear Enhancement in Southwestern Jordan" (1976). *LARS Symposia*. Paper 128.
http://docs.lib.purdue.edu/lars_symp/128

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

Symposium on

Machine Processing of

Remotely Sensed Data

June 29 - July 1, 1976

The Laboratory for Applications of
Remote Sensing

Purdue University
West Lafayette
Indiana

IEEE Catalog No.
76CH1103-1 MPRSD

Copyright © 1976 IEEE
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

DIGITAL IMAGE PROCESSING OF LANDSAT I MSS
DATA SPECIFICALLY DESIGNED FOR LINEAR
ENHANCEMENT IN SOUTHWESTERN JORDAN

Pat S. Chavez, Jr.
G. Lennis Berlin
Alex V. Acosta
U. S. Geological Survey
Flagstaff, Arizona

LINEAR ATMOSPHERIC TRANSFORM
ON LANDSAT MEASUREMENTS

Richard K. Kiang
GTE Information Systems
Goddard Institute for Space Studies
New York, N.Y. 10025

William E. Collins
Goddard Institute for Space Studies
New York, N.Y. 10025

ABSTRACT

This study was designed to take LANDSAT-1 multispectral scanner digital data and use digital image processing techniques to highlight linears. Two computer processing techniques were used on the data: (1) two dimensional high-pass filtering of MSS bands 5, 6, and 7 and (2) the horizontal derivative of MSS band 6.

The primary objective of this research was to evaluate structural reconnaissance of an arid area in southwestern Jordan. Interpretive analyses indicate that the structural pattern appears to be much more detailed and complex than indicated by presently available maps. The types of lines that were identifiable as lineaments include faults, joints, fractures, topographic crests, lithologic contacts, bedding traces and flexures. A noticeable difference between published maps and maps from the high-pass filtered images is the number of long lineaments that have never been mapped as faults. High-pass filtered images displayed the largest linears in various lithologies. The horizontal derivative image proved to be an excellent data source for obtaining a perspective for the structural fabric.

ABSTRACT

A problem exists when applying a set of ground training signatures determined under one atmospheric condition to the same area under another atmospheric condition. Sometimes the training signatures need to be adapted to the atmospheric condition of the subject area before classification. Other times the radiance measurements of an area with variable atmospheric conditions need to be adjusted to a common atmospheric condition before classification.

In order to describe the effects of the atmosphere on the ground-reflected radiance measured by the LANDSAT-1 satellite, a radiative transfer model was developed. The model is a combination of a doubling model, which describes the effects of molecular and aerosol scattering in the atmosphere, and a scaling-approximation absorption model which takes into account the gaseous absorption due to oxygen, ozone and water vapor in the LANDSAT-1 spectral region.

The model indicates that atmospheric effects can be approximated by linear transforms. The transform shows that the atmosphere always degrades the albedo difference but not necessarily the color difference. Signature transformations of Lambertian surfaces due to changes in the atmospheric condition can be determined.