

1-1-1976

Improved Signature Definition through Boundary-Edited Clustering

Daniel P. Rice

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Rice, Daniel P., "Improved Signature Definition through Boundary-Edited Clustering" (1976). *LARS Symposia*. Paper 138.
http://docs.lib.purdue.edu/lars_symp/138

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

Symposium on

Machine Processing of

Remotely Sensed Data

June 29 - July 1, 1976

The Laboratory for Applications of
Remote Sensing

Purdue University
West Lafayette
Indiana

IEEE Catalog No.
76CH1103-1 MPRSD

Copyright © 1976 IEEE
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

IMPROVED SIGNATURE DEFINITION THROUGH
BOUNDARY-EDITED CLUSTERING*

Daniel P. Rice
Environmental Research Institute
of Michigan
Ann Arbor, Michigan

ABSTRACT

It is desirable to enhance the performance of clustering techniques, so that cluster statistics more usefully represent the true spectral classes in the scene. If this is done, then improvements in classification can be realized, leading for example to more accurate large area crop surveys.

To improve the statistics available from a clustering algorithm, an edge-detection algorithm is used. The edge detector provides the cluster routine with inputs to filter out pixels which are likely to represent spectral response to a mixture of more than one ground cover type. The result is that the proportion of pixels representing pure classes in the scene is enriched.

While nearly all boundary detection algorithms are candidates for such use, this paper deals primarily with one which uses a gradient technique. Briefly, the gradient technique measures the spatial rate of change of signal value, averaged over all spectral bands. This is done by performing subtractions between neighboring pixels, and combining the along-scan and across-scan differences. Once the "gradient" at each point is calculated, a threshold is established so as to reject a specified percentage of the pixels to be used by the cluster routine. The paper will discuss the choice of the threshold, so as to filter most "mixture" pixels and yet avoid discarding too much, and perhaps losing high-texture ground cover types.

The effect of using the gradient filter on clustering seems to lie primarily in causing a greater proportion of pixels to fall in clusters which are near modes representing pure cover types, but also in bunching of the cluster centers somewhat closer to the modes. This aspect is discussed in conjunction with how the gradient filter affects the overall distribution function. In the quest toward improving techniques of clustering, a useful goal should be to improve the positioning of cluster centers relative to meaningful modes in the distribution. The enhancement of modes by the gradient filtering is an important step.

Two significant advantages are believed to result from the technique. First, it provides a suitable means of data reduction when clustering large scenes, so as to save processing costs. And second, the clusters which result more accurately represent the true ground classes, leading to better classification performance and more effective use of cluster-based signature extension techniques.

* This work is being performed under Contract NAS9-14123 for the Earth Observations Division of the NASA Johnson Space Center, Houston, Texas.