

1-1-1976

Statistical Analysis of Texture in LANDSAT Images of the United Kingdom

R. M. Lee

P. Gray

M. E. Barnett

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Lee, R. M.; Gray, P.; and Barnett, M. E., "Statistical Analysis of Texture in LANDSAT Images of the United Kingdom" (1976). *LARS Symposia*. Paper 145.

http://docs.lib.purdue.edu/lars_symp/145

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

Symposium on

Machine Processing of

Remotely Sensed Data

June 29 - July 1, 1976

The Laboratory for Applications of
Remote Sensing

Purdue University
West Lafayette
Indiana

IEEE Catalog No.
76CH1103-1 MPRSD

Copyright © 1976 IEEE
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

STATISTICAL ANALYSIS OF TEXTURE IN
LANDSAT IMAGES OF THE UNITED KINGDOM

R. M. Lee, P. Gray and M. E. Barnett
Blackett Laboratory
London SW7 2BZ

ABSTRACT

The agricultural regions of England appear in LANDSAT IMAGES as extended mosaic patterns of small fields. Individual fields, especially in the hillier north and west of the country may be as small as a few pixels and thus it is the field patterns themselves which make up the texture of the image. We have been concerned with the problem of mapping the average orientation of fields (a secondary indicator for geomorphology) and the distribution of field size.

Two approaches are being used:

- (i) Holographic derivation of local autocorrelation information.
- (ii) Digital extraction of statistical parameters.

For our test area in East Cornwall we have compared optically and digitally derived autocorrelation functions. We have also studied the local stationarity of the first and second order statistical measures which is crucial when evaluating statistical classification strategies.

A REAL TIME FREEZE PREDICTION MODEL BASED
UPON REMOTELY SENSED SURFACE TEMPERATURES

R. A. Sutherland¹ and J. F. Bartholic²

ABSTRACT

A mathematical model based on a numerical solution to the earth-air interface energy flux density equation is described. The model uses synoptic information, real time satellite data and standard meteorological measurements at key locations to produce short-range air and surface temperature predictions for the Florida peninsula during critical freeze nights. A statistical approach is then used to produce a detailed lattice of temperatures with the values from the key locations. The lattice of values is used to produce a thermal map of the Florida peninsula that is displayed on a television screen in a format identical to that used presently to display the satellite thermal data. The work stems from a cooperative project with NOAA, NASA and the University of Florida, to improve freeze forecasts.

¹Assistant Research Scientist, Fruit Crops Department, IFAS, University of Florida.

²Associate Professor, Fruit Crops Department, IFAS, University of Florida.