### Purdue University Purdue e-Pubs

LARS Symposia

Laboratory for Applications of Remote Sensing

1-1-1976

# Computer Analysis of Remotely Sensed Wheat Canopy Data

M. S. Sohel

Follow this and additional works at: http://docs.lib.purdue.edu/lars\_symp

Sohel, M. S., "Computer Analysis of Remotely Sensed Wheat Canopy Data" (1976). *LARS Symposia*. Paper 160. http://docs.lib.purdue.edu/lars\_symp/160

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

# Symposium on

# **Machine Processing of**

# **Remotely Sensed Data**

### June 29 - July 1, 1976

The Laboratory for Applications of Remote Sensing

Purdue University West Lafayette Indiana

IEEE Catalog No. 76CH1103-1 MPRSD

Copyright © 1976 IEEE The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

#### COMPUTER ANALYSIS OF REMOTELY SENSED

#### WHEAT CANOPY DATA

M. S. Sohel Prairie View A & M University Prairie View, Texas

#### ABSTRACT

The existing mathematical models of wheat growth or yield are based on air temperatures two meters above the ground from standard nationwide or worldwide meteorological networks, whereas temperatures obtained by thermal infrared scanners on satellites, such as the current NOAA series are surface radiative temperatures. These temperatures are correlated, but are not necessarily the same. In order to use satellite-derived surface radiative temperatures in wheat growth or yield models developed using standard air temperatures, the empirical relationship between these two temperatures needs to be established for the range of anticipated conditions.

Experiments were performed for bare soil and wheat canopies under irrigated and non-irrigated conditions. Measurements were made between 0900 and 1500 hours of radiative temperatures, air and soil temperatures. Precipitation, soil moisture, relative humidity, sunlight and windspeed were also recorded.

Analysis of the data shows that a relationship of the form Y = AX + B exists between radiative temperature and air temperature of wheat canopies. Additional effects of diurnal and seasonal variations have been found.