

1-1-1979

# Estimation of Soil Moisture Status and Actual Evapotranspiration Using Remotely-Sensed Data

R. J. Gurney

Follow this and additional works at: [http://docs.lib.purdue.edu/lars\\_symp](http://docs.lib.purdue.edu/lars_symp)

---

Gurney, R. J., "Estimation of Soil Moisture Status and Actual Evapotranspiration Using Remotely-Sensed Data" (1979). *LARS Symposia*. Paper 265.  
[http://docs.lib.purdue.edu/lars\\_symp/265](http://docs.lib.purdue.edu/lars_symp/265)

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact [epubs@purdue.edu](mailto:epubs@purdue.edu) for additional information.

Reprinted from

**Symposium on  
Machine Processing of  
Remotely Sensed Data**

**June 27 - 29, 1979**

The Laboratory for Applications of  
Remote Sensing

Purdue University  
West Lafayette  
Indiana 47907 USA

IEEE Catalog No.  
79CH1430-8 MPRSD

Copyright © 1979 IEEE  
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to [pubs-permissions@ieee.org](mailto:pubs-permissions@ieee.org).

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

# ESTIMATION OF SOIL MOISTURE STATUS AND ACTUAL EVAPOTRANSPIRATION USING REMOTELY-SENSED DATA

R. J. GURNEY  
Institute of Hydrology

Two digital models are described which relate soil surface temperatures estimated from aerial radiometric measurements to average daily soil moisture status and actual evapotranspiration. Both models estimate times series of surface temperatures from meteorological and soil measurements. The surface temperatures estimated using the models and the radiometer are compared, and the parameters of the models optimized until the differences are small. The parameters are carefully chosen to have unique relationships with soil moisture status and actual evapotranspiration, which may then be found.

The numerical models are expensive to implement, as are the models of atmospheric effects which must be used when estimating the surface temperatures from radiometric data, and so look-up versions of the models are used. The models are implemented for those input data which occur most frequently, these inputs being found by estimating joint frequency distributions from the inputs, and results for other inputs are found by interpolation within the look-up tables. In order to do this, the input data must be accurately registered geometrically and procedures for doing this, for estimating the joint frequency distributions efficiently, for constructing the look-up tables and then making maps of actual evapotranspiration and soil moisture status are described. The procedures are illustrated using data from a field experiment in Buckinghamshire, England in September 1977.