

9-2008

# Global Transcriptional Response of the Alkalitolerant Cyanobacterium *Synechocystis* sp. Strain PCC 6803 to pH 10.

Tina C. Summerfield

Louis A. Sherman  
*Purdue University*, lsherman@purdue.edu

Follow this and additional works at: <http://docs.lib.purdue.edu/bioscipubs>

---

## Recommended Citation

Summerfield, Tina C. and Sherman, Louis A., "Global Transcriptional Response of the Alkalitolerant Cyanobacterium *Synechocystis* sp. Strain PCC 6803 to pH 10." (2008). *Department of Biological Sciences Faculty Publications*. Paper 30.  
<http://dx.doi.org/10.1128/AEM.00883-08>

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact [epubs@purdue.edu](mailto:epubs@purdue.edu) for additional information.

## Global Transcriptional Response of the Alkali-Tolerant Cyanobacterium *Synechocystis* sp. Strain PCC 6803 to a pH 10 Environment

Tina C. Summerfield and Louis A. Sherman  
*Appl. Environ. Microbiol.* 2008, 74(17):5276. DOI:  
10.1128/AEM.00883-08.  
Published Ahead of Print 7 July 2008.

---

Updated information and services can be found at:  
<http://aem.asm.org/content/74/17/5276>

---

**SUPPLEMENTAL MATERIAL**

*These include:*

[Supplemental material](#)

**REFERENCES**

This article cites 62 articles, 26 of which can be accessed free  
at: <http://aem.asm.org/content/74/17/5276#ref-list-1>

**CONTENT ALERTS**

Receive: RSS Feeds, eTOCs, free email alerts (when new  
articles cite this article), [more»](#)

---

---

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>  
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

---

# Global Transcriptional Response of the Alkali-Tolerant Cyanobacterium *Synechocystis* sp. Strain PCC 6803 to a pH 10 Environment<sup>∇†</sup>

Tina C. Summerfield and Louis A. Sherman\*

Department of Biological Sciences, Purdue University, 201 S. University St., Hansen Hall, West Lafayette, Indiana 47907

Received 17 April 2008/Accepted 26 June 2008

Many cyanobacterial strains are able to grow at a pH range from neutral to pH 10 or 11. Such alkaline conditions favor cyanobacterial growth (e.g., bloom formation), and cyanobacteria must have developed strategies to adjust to changes in CO<sub>2</sub> concentration and ion availability. *Synechocystis* sp. strain PCC 6803 exhibits similar photoautotrophic growth characteristics at pH 10 and pH 7.5, and we examined global gene expression following transfer from pH 7.5 to pH 10 to determine cellular adaptations at an elevated pH. The strategies used to develop homeostasis at alkaline pH had elements similar to those of many bacteria, as well as components unique to phototrophic microbes. Some of the response mechanisms previously identified in other bacteria included upregulation of Na<sup>+</sup>/H<sup>+</sup> antiporters, deaminases, and ATP synthase. In addition, upregulated genes encoded transporters with the potential to contribute to osmotic, pH, and ion homeostasis (e.g., a water channel protein, a large-conductance mechanosensitive channel, a putative anion efflux transporter, a hexose/proton symporter, and ABC transporters of unidentified substrates). Transcriptional changes specific to photosynthetic microbes involved NADH dehydrogenases and CO<sub>2</sub> fixation. The pH transition altered the CO<sub>2</sub>/HCO<sub>3</sub><sup>-</sup> ratio within the cell, and the upregulation of three inducible bicarbonate transporters (BCT1, SbtA, and NDH-1S) likely reflected a response to this perturbed ratio. Consistent with this was increased transcript abundance of genes encoding carboxysome structural proteins and carbonic anhydrase. Interestingly, the transition to pH 10 resulted in increased abundance of transcripts of photosystem II genes encoding extrinsic and low-molecular-weight polypeptides, although there was little change in photosystem I gene transcripts.

Cyanobacteria are among the most alkaliphilic microbes, and they frequently dominate alkaline environments, such as soda lakes and microbial mats (30, 36). In addition to pH, key parameters that include nutrient availability and temperature influence the population composition of phytoplankton communities (18). However, cyanobacterial bloom formation is usually accompanied by an elevated pH that results from increased photosynthesis that depletes CO<sub>2</sub>. Many cyanobacterial strains are alkali tolerant and grow at pHs ranging from neutral to 10 to 11, so that cyanobacteria both generate and thrive in alkaline conditions. There have been numerous reports of habitats where photosynthetic rates are high (such as shallow lakes), pH values exceed pH 10, and cyanobacteria become the major phytoplankton species (7, 27). Such populations of cyanobacteria are frequently associated with the production of a range of secondary metabolites, including nuisance and toxic compounds (5). This has led to experiments aimed at reducing cyanobacterial populations; e.g., it has been established for a long time that adding carbon dioxide or acid to lower the pH of lake samples can increase the abundance of green algae relative to that of cyanobacteria (42).

One reason that cyanobacteria have an advantage over other phytoplankton species at high pH is that the carbon-concen-

trating mechanism of cyanobacteria is better able to utilize bicarbonate than the mechanism in green algae (21). It is anticipated that cyanobacteria must employ additional mechanisms to maintain pH homeostasis in order to flourish at high pH. Many nonphotosynthetic bacteria are able to survive or grow at alkaline pH, and they respond to increased pH using a variety of mechanisms to maintain homeostasis within the cell; the best characterized of these strategies is the increased expression and activity of monovalent cation/proton antiporters (33). These transmembrane proteins maintain the intracellular pH through the uptake of protons, utilizing outward monovalent cation gradients. Multiple cation/proton antiporters have been identified in cyanobacteria, and their involvement in pH homeostasis has been suggested by gene knockout studies with different strains exhibiting altered pH- and NaCl-sensitive phenotypes (3, 14, 57). Additional bacterial responses aimed at regulating intracellular pH include elevated metabolic acid production (via amino acid deaminases and sugar fermentation), increased ATP synthase activity (H<sup>+</sup> entry coupled to ATP generation), and altered cell surface properties (33, 52). However, the extent to which these strategies are employed and their impact in different bacterial strains remain to be determined.

Compared to other alkaliphilic bacteria, cyanobacteria have two additional complexities, photosynthetic (thylakoid) membranes and the presence of ATP synthase in both thylakoid and plasma membranes (43). Compartments within the cell are maintained at different pHs, and the thylakoid lumen has a pH that is ~2 units lower than the pH of the cytosol (2). Changes in the external pH have been

\* Corresponding author. Mailing address: Department of Biological Sciences, Purdue University, 201 S. University St., Hansen Hall, West Lafayette, IN 47907. Phone: (765) 494-8106. Fax: (765) 496-1496. E-mail: lsherman@purdue.edu.

† Supplemental material for this article may be found at <http://aem.asm.org/>.

∇ Published ahead of print on 7 July 2008.

shown to alter both the cytoplasmic and thylakoid lumen pHs, with an increase in the external pH of 2 pH units resulting in an increase of  $\sim 0.2$  pH unit (2, 40). Therefore, growth of cyanobacteria in alkaline environments requires maintenance of pH gradients across multiple membrane systems, regulation of inorganic carbon uptake, and adjustment to changes in the abundance of different ions.

We investigated the impact of a pH transition from pH 7.5 to pH 10 in *Synechocystis* sp. strain PCC 6803. This freshwater cyanobacterium is a halo- and alkali-tolerant strain which exhibits similar growth at pH 7.5 and pH 10 (8). Six genes have been annotated as genes that encode sodium/proton antiporters in *Synechocystis* sp. strain PCC 6803 (19). It is likely that these proteins have overlapping functions, and this has made it difficult to define their role in pH homeostasis (59). In addition, *Synechocystis* sp. strain PCC 6803 accumulates acetolactate under alkaline conditions, and it has been suggested that this is a mechanism for pH homeostasis (28).

The importance of external pH has been demonstrated in *Synechocystis* sp. strain PCC 6803 by identification of a number of pH-sensitive photosystem II (PSII) mutants that are able to grow photoautotrophically at pH 10 but not at pH 7.5 (8, 54). Each of these pH-sensitive strains contains two mutations in PSII, including the absence of either the PsbO or PsbV luminal protein. The cellular adaptations that occur during changes in the external pH that make the differential growth possible have not been identified. We examined global gene expression in *Synechocystis* sp. strain PCC 6803 following a transition to high pH by establishing a time course to identify genes that showed pH-dependent expression at 1 h ( $t_1$ ), 2 h ( $t_2$ ), and 6 h ( $t_6$ ) following transfer from pH 7.5 to pH 10. This study revealed that the response of *Synechocystis* sp. strain PCC 6803 to alkaline conditions was cell-wide and included strategies typical of many bacteria, as well as strategies specific to phototrophic microbes. The levels of transcripts of a number of the components involved in acclimation to alkaline pH in other bacteria, such as monovalent cation/proton antiporters and ATP synthase, were elevated at pH 10. We observed increased abundance of transcripts of additional transporters with the potential to contribute to osmotic, pH, and ion homeostasis. Changes specific to photosynthesis included the upregulation of genes encoding three bicarbonate transport systems, probably in response to a perturbed  $\text{CO}_2/\text{HCO}_3^-$  ratio within the cell. Consistent with this was increased abundance of transcripts of genes encoding carboxysome structural proteins and carbonic anhydrase. The transcripts of a number of genes encoding transcriptional regulators were differentially regulated at pH 10. Furthermore, we observed that at an elevated pH, the levels of transcripts of genes encoding PSII extrinsic and low-molecular-weight polypeptides were increased.

#### MATERIALS AND METHODS

**Growth conditions.** The glucose-tolerant organism *Synechocystis* sp. strain PCC 6803 (60) was grown at  $30 \pm 2^\circ\text{C}$  using cool white fluorescent light at an intensity of  $\sim 30$  microeinsteins  $\text{m}^{-2} \text{s}^{-1}$  with shaking at 125 rpm in BG-11 medium (4). The pH of the BG-11 medium was maintained by addition of either 25 mM HEPES (pH 7.5) or 25 mM 3-(cyclohexylamino)-1-propanesulfonic acid (CAPS) (pH 10) (8). The cell

densities of the cultures were determined by measuring the optical density at 750 nm ( $\text{OD}_{750}$ ) as previously described (6, 29).

**RNA isolation.** Total RNA was extracted and purified using phenol-chloroform extraction and CsCl gradient purification as previously described (39, 48).

**Microarray design and analysis.** The microarray platform used and construction of this platform were described previously by Postier et al. (38), and the cDNA labeling, prehybridization, and hybridization protocols were described in detail by Singh et al. (47). The microarray experiment involved a loop design that allowed comparison of *Synechocystis* sp. strain PCC 6803 at different time points following a transition from pH 7.5 to pH 10 by using an analysis of variance model (26, 47). Cells were grown in BG-11 medium at pH 7.5 until the  $\text{OD}_{750}$  was  $\sim 0.2$  (approximately  $8 \times 10^7$  cells/ml) before they were harvested by centrifugation ( $5,000 \times g$  for 5 min) and transferred to pH 10 at an  $\text{OD}_{750}$  of  $\sim 0.2$ . Samples were collected immediately after transfer to pH 10 ( $t_0$ ) and at  $t_1$ ,  $t_2$ , and  $t_6$  after transfer.

Data acquisition and analysis were performed as described by Singh et al. (47); this included an analysis of variance model approach to test the null hypothesis that a particular gene's expression level was not different over time, and a  $P$  value was calculated. We used a false discovery rate (FDR) of 5% to control the proportion of significant results that were type I errors (false rejection of the null hypothesis) as described previously (55). Genes that had an FDR of 0.05 (corresponding to 5% expected false positives) and that exhibited a change of at least 1.5-fold were considered interesting and retained for further analysis. The  $P$  values for these genes ranged from  $6.0 \times 10^{-3}$  to  $1.4 \times 10^{-13}$ .

## RESULTS AND DISCUSSION

**Global transcriptional response to the transition from pH 7.5 to pH 10.** Approximately 7, 12, and 10% of the chromosomal genes were differentially regulated at  $t_1$ ,  $t_2$ , and  $t_6$  after transfer from pH 7.5 to pH 10, respectively. Genes were divided into functional categories according to Cyanobase (<http://bacteria.kazusa.or.jp/cyanobase>), and the number of differentially expressed genes in each category is shown in Table 1. Excluding hypothetical and unknown genes, photosynthesis and respiration was the category with the largest number of differentially regulated genes following transfer to pH 10, and these genes were almost all upregulated. Other categories with elevated levels of transcripts after transfer to pH 10 included proteins with regulatory functions and transport and binding proteins (Table 1).

**pH-independent gene expression.** Genes that were upregulated both after transfer from pH 7.5 to pH 10 and after transfer from pH 10 to pH 7.5 were designated pH independent and are listed separately from the pH 10-responsive data set in Table 1. These genes were identified by combining data from the microarray experiment examining transfer from pH 7.5 to pH 10 with data from a similar experiment examining the transition from pH 10 to pH 7.5. There were 198 genes whose transcription was found to change independent of the direction of the pH transition, and  $>75\%$  of these genes were upregulated after transfer. Upregulated genes encoding ribosomal proteins accounted for approximately one-quarter of the differentially expressed genes. Other categories with differentially expressed genes included photosynthesis and respiration, energy metabolism, and biosynthesis of cofactors, prosthetic groups, and carriers (Table 2). Increased abundance of the transcripts of a number of these genes, such as those encoding ribosomal proteins and ATP synthase, has been associated with favorable growth conditions, including light-versus-dark transition and log-phase growth versus stationary-phase growth (10, 53). However, the doubling times following the transition from pH 7.5 to pH 10 and the transition from pH 10 to pH 7.5 remained  $\sim 12$  h (data not shown). Additional pH-

TABLE 1. Functional categories of genes differentially regulated at pH 10 compared to pH 7.5 in *Synechocystis* sp. strain PCC 6803<sup>a</sup>

General pathway	No. of genes <sup>b</sup>	No. of differentially regulated genes (no. of genes upregulated) <sup>c</sup>			
		pH independent ( $t_2/t_0$ ) <sup>d</sup>	pH 10 compared to pH 7.5		
			$t_1/t_0$	$t_2/t_0$	$t_6/t_0$
Amino acid biosynthesis	97	6 (5)	6 (6)	10 (9)	9 (8)
Biosynthesis of cofactors, prosthetic groups, and carriers	124	11 (11)	10 (6)	14 (8)	11 (8)
Cell envelope	67	2 (1)	2 (1)	4 (1)	3 (2)
Cellular processes	76	5 (4)	7 (6)	8 (8)	4 (4)
Central intermediary metabolism	31	4 (3)	0 (0)	1 (1)	1 (1)
DNA replication, restriction, recombination, and repair	60	0 (0)	4 (4)	11 (4)	9 (2)
Energy metabolism	132	11 (10)	14 (10)	13 (7)	15 (12)
Hypothetical	1,076	40 (22)	73 (37)	130 (71)	96 (62)
Other categories	175 <sup>e</sup>	10 (8)	15 (7)	28 (8)	27 (11)
Photosynthesis and respiration	141	23 (23)	23 (22)	30 (28)	34 (31)
Purines, pyrimidines, nucleosides, and nucleotides	41	3 (3)	2 (1)	4 (2)	5 (2)
Regulatory functions	146	4 (2)	14 (13)	18 (13)	19 (15)
Transcription	30	3 (3)	2 (2)	2 (1)	3 (2)
Translation	168	51 (51)	12 (11)	15 (12)	15 (13)
Transport and binding proteins	196	3 (3)	13 (10)	19 (10)	20 (15)
Unknown	474	22 (9)	19 (4)	63 (15)	45 (17)
Total	3,165	198 (158)	216 (140)	370 (198)	316 (205)

<sup>a</sup> Genes were considered differentially regulated when the FDR was 0.05 (change, >1.5-fold).

<sup>b</sup> Total number of genes based on Kazusa annotation prior to May 2002.

<sup>c</sup> The upregulated genes were the genes upregulated at pH 10 compared to pH 7.5.

<sup>d</sup> Differentially regulated genes after transfer from pH 7.5 to pH 10 and after transfer from pH 10 to pH 7.5.

<sup>e</sup> The number does not include genes encoding transposases.

independent transcriptional changes likely include mechanisms to maintain cellular homeostasis and are discussed in more detail below.

**Photosynthesis and respiration.** Cyanobacterial NADH dehydrogenase complexes (NDH-1) are composed of a multiprotein core (NDH-1M) and NdhD and NdhF subunits. Different NdhD and NdhF subunits determine whether an NDH-1 complex functions in cyclic electron transfer around PSI and respiratory electron transfer or CO<sub>2</sub> uptake (1). Genes encoding 7 of the 13 NDH-1M core proteins, including *ndhD2*, were upregulated at pH 10 (Table 3). The protein encoded by *ndhD2* is hypothesized to associate with the NDH-1M core along with NdhF1 to form the NDH-1L' complex, and gene knockout studies indicated that this subunit is involved in PSI cyclic electron transfer (32). The NdhD3 and NdhF3 subunits, together with the proteins designated CupA and CupS, associate with NDH-1M to form a low-CO<sub>2</sub>-inducible transporter (Sll1732 to Sll1735) (62). The operon consisting of *sll1732* to *sll1735* was upregulated both after transfer to pH 10 and after transfer to pH 7.5 (Table 2).

The transcript encoding a  $\beta$ -type carbonic anhydrase (*slr1347*) was upregulated after transfer to pH 10 (Table 3). This protein is located in the carboxysome, where it catalyzes the conversion of HCO<sub>3</sub><sup>-</sup> to CO<sub>2</sub> and both activates ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) and compensates for the low affinity of Rubisco for CO<sub>2</sub> (21, 51). The assembly and structural protein components of the carboxysome are encoded by the carbon-concentrating mechanism (*ccm*) genes; the abundance of the transcripts was increased after transfer to pH 10, and the abundance of many of them was increased after transfer to pH 7.5 (Table 2).

The genes encoding the PSII extrinsic proteins PsbO, PsbP, PsbQ, and PsbU were upregulated at pH 10. In addition, the genes encoding seven low-molecular-weight proteins implicated in assembly or stability of PSII centers were upregulated at pH 10 (44). These genes comprised the *psbEFLJ* operon, whose genes encode the cytochrome *b*<sub>559</sub>  $\alpha$  and  $\beta$  subunits, PsbL, and PsbJ, respectively, and the *psbK*, *psbM*, and *psbX* genes. Increased abundance was observed for the *sll1414* transcript encoding the PSII-associated protein Psb29, which is thought to be involved in PSII biogenesis (22). There was no change in the abundance of transcripts for the core proteins D1, D2, CP47, and CP43, and the genes encoding these proteins were highly expressed at both pH 7.5 and pH 10. This is consistent with the similar oxygen evolution rates and PSII abundance observed at pH 10 and pH 7.5 (54).

ATP synthase genes were upregulated after transfer to both pH 7.5 and pH 10 (Table 2). As ATP synthase is associated with both the thylakoid and plasma membranes, the pH regulation of this complex may be different from that observed for nonphotosynthetic bacteria.

**Regulatory genes.** Genes encoding 21 regulatory components were upregulated after transfer to pH 10, including genes that had different transcriptional kinetics (Table 3; see Table S1 in the supplemental material). Following transfer to pH 10, increased abundance was observed for the transcripts of two-component system histidine kinases (*hik9* and *hik12*) and response regulators (*rre7* and *rre5*), a serine/threonine kinase, the global nitrogen regulator, *ntcA*, and several putative transcription factors (Table 3). The response regulator *rre5* gene was upregulated under inorganic carbon (C<sub>i</sub>) limitation conditions

TABLE 2. Selected differentially regulated pH-independent genes in *Synechocystis* sp. strain PCC 6803 after transfer from pH 7.5 to pH 10 and after transfer from pH 10 to pH 7.5<sup>a</sup>

Gene	Designation or function	Change (fold) after transfer to:			
		pH 10			pH 7.5 (t <sub>2</sub> )
		t <sub>1</sub>	t <sub>2</sub>	t <sub>6</sub>	
<b>Photosynthesis and respiration</b>					
<b>ATP synthase</b>					
sll1321	Hypothetical	1.7	1.4	1.4	3.0
sll1322	<i>atpI</i>	1.3	1.2	1.2	2.9
ssl2615	<i>atpH</i>	2.1	1.8	1.8	3.7
sll1323	<i>atpG</i>	2.1	1.7	1.7	4.7
sll1324	<i>atpF</i>	2.5	2.1	2.0	5.5
sll1325	<i>atpD</i>	3.1	2.8	2.6	4.5
sll1326	<i>atpA</i>	2.0	1.8	1.7	3.1
sll1327	<i>atpC</i>	1.9	1.6	1.6	3.6
slr1329	<i>atpB</i>	2.1	1.7	1.8	2.6
slr1330	<i>atpE</i>	3.0	2.5	2.4	3.1
<b>CO<sub>2</sub> fixation: carboxysome</b>					
sll1028	<i>ccmK2</i>	2.0	2.0	2.1	2.4
sll1029	<i>ccmK1</i>	2.4	2.4	2.6	2.2
sll1030	<i>ccmL</i>	1.8	1.8	2.3	2.4
sll1031	<i>ccmM</i>	2.3	1.9	2.2	2.1
sll1032	<i>ccmN</i>	1.7	1.4	1.9	1.9
slr1838	<i>ccmK3</i>	1.3	1.8	1.8	1.3
slr1839	<i>ccmK4</i>	1.3	1.7	1.4	1.6
<b>NADH dehydrogenase: bicarbonate transport</b>					
sll1732	<i>ndhF3</i>	2.1	1.4	3.7	1.3
sll1733	<i>ndhD3</i>	2.8	2.1	4.6	1.8
sll1734	<i>cupA</i>	3.3	2.7	5.3	2.9
sll1735	Hypothetical	2.0	1.6	3.8	3.2
<b>Regulatory functions</b>					
slr0473	<i>cph1</i>	-2.4	-1.9	-1.9	-1.8
slr0474	<i>rcp1</i>	-4.6	-3.6	-3.4	-2.3
<b>Transport proteins: bicarbonate</b>					
slr0040	<i>cmpA</i>	1.4	1.1	5.3	2.9
slr0041	<i>cmpB</i>	1.4	1.2	9.4	2.8
slr0043	<i>cmpC</i>	1.5	1.4	2.4	1.6
slr0044	<i>cmpD</i>	1.3	1.5	3.3	2.0
slr1512	<i>sbtA</i>	2.9	2.1	5.5	3.7
slr1513	<i>sbtB</i>	2.5	2.1	6.5	2.9
<b>Gene clusters</b>					
slr1667	Hypothetical	-1.8	-2.5	-1.7	-5.8
slr1668	Hypothetical	-1.5	-1.7	-1.3	-1.5
sll1077	<i>speB2</i>	1.7	2.2	-1.1	2.2
sll1078	<i>hypA2</i>	2.2	3.5	-1.1	1.6
sll1079	<i>hypB2</i>	2.5	3.9	-1.4	1.6
sll1080	Transport	2.3	2.7	-1.1	-1.4
sll1081	Transport	1.8	2.4	-1.3	ND <sup>b</sup>
sll1082	Transport	1.4	1.9	-1.2	-1.1

<sup>a</sup> Genes were considered differentially regulated when the FDR was 0.05 (change, >1.5 fold) (indicated by bold type).

<sup>b</sup> ND, not determined.

and was suggested previously to be involved in CO<sub>2</sub> uptake and pH homeostasis (58). In addition, sll1937 and sll0567, encoding putative ferric uptake regulation (Fur) proteins, were upregulated at pH 10 from t<sub>1</sub> to t<sub>6</sub> (1.7- to 2.9-fold). Fur proteins are metal ion uptake regulators, and the sll0567 product is essential for growth under normal culture conditions and is part of the iron-responsive regulation mechanism (25). The role of

sll1937 is not clear, as deletion of this gene did not alter iron-stress-induced gene expression (25). Transcripts of a histidine kinase and response regulator (*cph1/rcp1*) exhibited strong pH-independent downregulation (Table 2). These transcripts are upregulated in the dark and are thought to be involved in regulation at light-dark transitions (49).

**Stress response.** The slr1516 gene, encoding superoxide dismutase (*sodB*), was upregulated at pH 10 (1.5-fold). Expression of this gene, which encodes an antioxidant, is induced by various stress conditions, including temperature (high or low), salt, hydrogen peroxide, and light (57). Superoxide dismutase converts reactive oxygen species to hydrogen peroxide, which then is scavenged by catalases or peroxidases or both. The slr1992 gene, encoding glutathione peroxidase, was upregulated at t<sub>1</sub> through t<sub>6</sub> (1.9- to 1.8-fold). Two genes encoding thioredoxin (slr0623 and slr1139) were upregulated at t<sub>2</sub> and t<sub>6</sub> (1.5- to 1.7-fold). The levels of the slr0623 transcript were high, consistent with the hypothesis that it encodes the most abundant of the four *Synechocystis* sp. strain PCC 6803 thioredoxins. Furthermore, Slr0623 has been suggested to have a major role in supplying reducing equivalents to the antioxidant systems (13). In addition, the genes encoding the chaperones HspA, DnaK1, and HtpG were upregulated at pH 10 (Table 3), and previous reports indicated that these genes are upregulated under various stress conditions, including oxidative stress (20, 26, 34, 45, 50). Two of these genes, *hspA* and *hspG*, were upregulated in a pseudorevertant of a PSII mutant, ΔPsbO:ΔPsbU. The original ΔPsbO:ΔPsbU strain is able to grow at pH 10, although it does not grow at pH 7.5, but the pseudorevertant was able to grow at both pH 10 and pH 7.5 (53).

**Monovalent cation/proton antiporters.** Six genes have been annotated as genes encoding Na<sup>+</sup>/H<sup>+</sup> antiporters in *Synechocystis* sp. strain PCC 6803, including sll0689 (*nhaS3*), whose transcript level was increased twofold at pH 10 (Table 3). Unlike four of the Na<sup>+</sup>/H<sup>+</sup> antiporters, sll0689 appeared to be essential for cell viability, as mutants lacking this gene could not be fully segregated (16, 59). Moreover, the partially segregated ΔSll0689 strain was sensitive to high-salt conditions at pH 9, and Sll0689 has a high affinity for both Na<sup>+</sup> and Li<sup>+</sup> ions (16, 59).

The eight-gene cluster containing slr2006 to slr2012 (including *ssr3410*) was upregulated 1.7- to 2.3-fold within 1 h at pH 10, and the transcript level remained elevated at 6 h (Table 3). Two of these genes were annotated as genes encoding NDH subunits; however, this cluster has similarity to genes encoding a putative multiprotein cation/H<sup>+</sup> antiporter in *Anabaena* sp. strain PCC 7120 (3). Interruption of one of these genes in *Anabaena* sp. strain PCC 7120 resulted in a strain that exhibited retarded growth at elevated pH and enhanced salt sensitivity at pH 10.5 (3). Blanco-Rivero et al. (3) designated this cluster *mup* (multiple resistance and pH adaptation) due to similarity to a *Bacillus subtilis* *mup* operon involved in Na<sup>+</sup> resistance, particularly at high pH (17). The similarity between subunits of the Mrp and NDH complexes has been described previously and may reflect a common origin and similar functions of the two complexes (12).

**Other transporters.** The transfer to pH 10 resulted in differential regulation of a number of transporters that may contribute to osmotic, pH, and ion homeostasis (Table 3). This

TABLE 3. Selected differentially regulated genes in *Synechocystis* sp. strain PCC 6803 after transition from pH 7.5 to pH 10<sup>a</sup>

Gene	Designation or function	Change (fold) at:			Gene	Designation or function	Change (fold) at:		
		t <sub>1</sub>	t <sub>2</sub>	t <sub>6</sub>			t <sub>1</sub>	t <sub>2</sub>	t <sub>6</sub>
Cellular processes: chaperones					slr1937	<i>fur</i>	<b>2.9</b>	<b>2.7</b>	<b>2.8</b>
slr1514	<i>hspA</i>	1.4	<b>1.9</b>	1.4	slr1423	<i>nicA</i>	1.2	<b>1.5</b>	<b>1.8</b>
slr0058	<i>dnaK1</i>	<b>1.7</b>	<b>1.7</b>	<b>1.7</b>	slr0594	Treg <sup>c</sup>	<b>1.5</b>	<b>1.7</b>	<b>1.6</b>
slr0430	<i>htpG</i>	<b>1.9</b>	<b>1.5</b>	1.3	slr0782	Treg	<b>1.6</b>	<b>1.9</b>	<b>1.6</b>
Photosynthesis and respiration					slr0599	<i>spkC</i>	1.3	<b>1.6</b>	<b>1.6</b>
PSII					slr0210	<i>hik9</i>	1.4	<b>1.9</b>	<b>1.7</b>
Oxygen-evolving complex					slr1672	<i>hik12</i>	<b>1.5</b>	1.2	<b>1.9</b>
slr0427	<i>psbO</i>	<b>1.5</b>	<b>1.9</b>	<b>1.7</b>	slr1042	<i>rre7</i>	<b>1.7</b>	<b>1.5</b>	<b>2.7</b>
slr1194	<i>psbU</i>	<b>1.5</b>	<b>1.6</b>	<b>1.7</b>	slr1594	<i>rre5</i>	<b>1.8</b>	1.2	-1.2
slr1418	<i>psbP</i>	1.4	<b>1.5</b>	<b>1.6</b>	Transport and binding proteins				
slr1638	<i>psbQ</i>	<b>1.6</b>	<b>2.1</b>	<b>1.7</b>	slr0689	<i>nhaS3</i>	<b>2.1</b>	1.4	<b>2.0</b>
Low-molecular-mass polypeptides					slr0771	<i>glcP</i>	<b>1.7</b>	<b>1.5</b>	<b>1.5</b>
slr0002	<i>psbX</i>	<b>2.4</b>	<b>2.2</b>	<b>2.1</b>	slr0753	Transport	1.3	1.3	<b>2.2</b>
slr0003	<i>psbM</i>	1.2	<b>1.5</b>	<b>1.5</b>	slr0875	<i>mscL</i>	1.4	<b>1.5</b>	<b>1.7</b>
slr0005	<i>psbK</i>	1.3	<b>1.5</b>	<b>1.7</b>	slr2057	<i>apqZ</i>	<b>1.9</b>	<b>2.2</b>	<b>2.9</b>
slr3451	<i>psbE</i>	1.4	<b>1.8</b>	<b>1.5</b>	Gene cluster				
slr0006	<i>psbF</i>	1.4	<b>1.9</b>	<b>1.6</b>	Putative cation/H <sup>+</sup> antiporter				
slr0007	<i>psbL</i>	1.2	<b>1.7</b>	<b>1.6</b>	slr2006	<i>mrpC</i>	<b>1.8</b>	1.2	<b>1.9</b>
slr0008	<i>psbJ</i>	1.3	<b>2.2</b>	<b>1.9</b>	slr2007	<i>mrpD</i>	<b>2.0</b>	<b>1.7</b>	<b>2.3</b>
Putative assembly protein					slr2008	<i>mrpC</i>	<b>1.6</b>	<b>1.6</b>	<b>1.8</b>
slr1414	<i>psb29</i>	1.4	<b>1.8</b>	<b>1.8</b>	slr2009	<i>mrpD</i>	<b>1.7</b>	<b>1.6</b>	<b>1.7</b>
NADH dehydrogenase: core subunits					slr2010	<i>mrpE</i>	<b>2.3</b>	<b>1.8</b>	<b>1.8</b>
slr0223	<i>ndhB</i>	<b>1.6</b>	1.4	<b>1.7</b>	slr3409	<i>mrpF</i>	1.4	1.4	<b>1.5</b>
slr0520	<i>ndhI</i>	<b>2.0</b>	<b>1.8</b>	<b>2.3</b>	slr3410	<i>mrpG</i>	<b>1.8</b>	<b>1.7</b>	<b>1.7</b>
slr0521	<i>ndhG</i>	<b>1.8</b>	<b>2.0</b>	<b>2.3</b>	slr2011	<i>mrpA</i>	<b>2.1</b>	<b>1.7</b>	<b>1.9</b>
slr0522	<i>ndhE</i>	<b>1.6</b>	<b>1.9</b>	<b>1.9</b>	slr2012	<i>mrpB</i>	<b>1.7</b>	1.4	<b>1.5</b>
slr1279	<i>ndhC</i>	<b>1.6</b>	<b>1.5</b>	<b>1.7</b>	slr1501	Other	<b>3.9</b>	<b>3.9</b>	<b>6.4</b>
slr1280	<i>ndhK</i>	<b>2.0</b>	<b>1.8</b>	<b>2.1</b>	slr1113	Transport	1.4	1.4	<b>2.0</b>
slr1281	<i>ndhJ</i>	<b>1.6</b>	1.4	<b>2.0</b>	slr1114	Hypothetical	1.3	1.3	<b>1.6</b>
slr1291	<i>ndhD2</i>	<b>2.5</b>	<b>1.6</b>	<b>1.8</b>	slr1392	Treg	1.3	<b>1.6</b>	<b>1.8</b>
CO <sub>2</sub> fixation					slr0408	Unknown	-1.4	-2.1	-1.9
slr1347	<i>cab</i>	<b>1.6</b>	<b>1.6</b>	<b>1.5</b>	slr0142	Hypothetical	ND <sup>b</sup>	ND	ND
slr0436	<i>ccmO</i>	<b>1.5</b>	<b>1.6</b>	<b>1.8</b>	slr0143	<i>hat</i>	ND	ND	ND
slr0934	<i>ccmA</i>	1.2	<b>1.5</b>	-1.1	slr0144	4VR <sup>d</sup>	ND	ND	ND
Regulatory functions					slr0145	Unknown	-1.5	1.4	1.4
slr0567	<i>fur</i>	<b>1.6</b>	<b>1.7</b>	<b>1.6</b>	slr0146	Hypothetical	-2.4	-1.2	-1.1
					slr0147	4VR	-1.9	-1.0	-1.1
					slr0148	Hypothetical	-2.6	-2.0	-2.0
					slr0149	Hypothetical	-2.7	-2.2	-2.1
					slr0150	<i>petF</i>	-1.8	-1.6	-1.6
					slr0151	Unknown	-2.1	-3.9	-3.5

<sup>a</sup> Genes were considered differentially regulated when the FDR was 0.05 (change, >1.5-fold) (indicated by bold type).

<sup>b</sup> ND, not determined.

<sup>c</sup> Treg, transcriptional regulator.

<sup>d</sup> 4VR, 4-vinyl reductase.

differential regulation included upregulation (1.9- to 2.9-fold) of the slr2057 transcript encoding a water channel protein (ApqZ). Water channels allow bidirectional movement of water and often highly specific movement of other compounds, although they are usually impermeable to ions (41). In cyanobacteria these channels have been postulated to have a role in CO<sub>2</sub> uptake (56). A ΔSlr2057 strain exhibited altered cell shrinkage and altered gene expression under hyperosmotic stress conditions (41). Another channel protein that is respon-

sive to osmotic shock had elevated transcript levels at pH 10. The gene (slr0875) encoded a high-conductance mechanosensitive channel; such channels are present in bacterial membranes and open in response to stretch forces in the lipid bilayer, preventing cell lysis (23). In addition, activation of Slr0875 was shown to result in Ca<sup>2+</sup> release (31). Increased abundance of the transcript for this channel may be part of a response directed at maintaining ion homeostasis at high pH. The same may be true of the slr0753 gene, encoding a putative




FIG. 1. Coordinately regulated gene clusters of *Synechocystis* sp. strain PCC 6803. (a) Gene cluster upregulated at pH 10 compared to pH 7.5. (b) Gene cluster showing pH-independent regulation. (c) Gene cluster containing many genes that exhibit upregulation after transfer to pH 10.

anion efflux transporter, which was upregulated >2-fold following 6 h at pH 10. Mutations in this gene removed the chloride requirement of a PSII strain lacking PsbV, suggesting that the gene encodes a chloride extrusion protein (24). Finally, the gene encoding a glucose transporter (*glcP*) was upregulated at pH 10. This transporter has been suggested to be a hexose/proton symporter and therefore may be involved in pH homeostasis (9).

In addition to the NDH-1-associated bicarbonate transport, two low- $C_i$ -inducible bicarbonate uptake systems were upregulated after transfer to both pH 10 and pH 7.5. These systems are BCT1, encoded by the *cmp* operon, and SbtA, plus the neighboring gene *sbtB*. Changes in the external pH have been shown to alter the cytosolic and thylakoid lumen pH (2); thus, the transfer of cells to a different pH alters the  $CO_2/HCO_3^-$  ratio within the cell. The upregulation of three inducible bicarbonate transporters (BCT1, SbtA, and NDH-1S) after transfer to pH 7.5 and pH 10 may reflect a response to this perturbed ratio.

**Metabolic acid production.** Deaminases have been suggested to play a role in acclimation to alkaline pH in some bacteria, and this appears to be true in *Synechocystis* sp. strain PCC 6803 as well. Upon transfer to pH 10, the level of the transcript encoding L-threonine deaminase (slr2072) was increased. In addition, a number of genes involved in the biosynthesis of valine, leucine, and isoleucine were upregulated (see Table S1 in the supplemental material). These genes included the gene encoding acetolactate synthase, *ilvB* (sll1981), which may play a role in pH homeostasis (26).

**Gene clusters.** The slr1501 gene was upregulated after transfer to pH 10 and was rapidly downregulated after transfer to pH 7.5. The adjacent gene sll1392 was similarly regulated, suggesting that there may be a divergent promoter. In addition, two genes downstream of slr1501 (slr1113 and slr1114) were upregulated at pH 10, and slr1113 was also downregulated after transfer to pH 7.5 (Fig. 1a and Table 3). The designations of these genes in Cyanobase are as follows: probable acetyltransferase gene, slr1501; transcriptional regulator gene, sll1392; ABC transporter ATP-binding protein gene, slr1113; and permease gene, slr1114. The accumulation of these transcripts at pH 10 and their striking downregulation at pH 7.5 suggest that this gene cluster has a role in growth at high pH.

The cluster containing genes slr0145 to slr0151 was downregulated at pH 10 (Fig. 1c and Table 3). Both slr0148 and slr0150 encode ferredoxinlike proteins, and slr0150 has been shown to be downregulated following high-light treatment (37). Two genes in this cluster (slr0144 and slr0147) contain 4-vinyl reductase motifs predicted to be involved in small-molecule binding (no data were available for slr0144 from this experiment due to a high *P* value). Another two genes (slr0146 and slr0149) encode proteins containing bilin-binding domains. This cluster was previously shown to be downregulated in iron-deficient media and in the presence of hydrogen peroxide, and it was suggested that the proteins may be involved in PSI function and assembly (46). This function would be consistent with the downregulation of this cluster and the upregulation of a number of PSII genes at pH 10. The slr0408 gene is upstream of this cluster and was upregulated at pH 10 (Fig. 1c and Table 3). This large gene encodes an unknown protein with a putative  $Ca^{2+}$  expulsion domain.

The slr1667 and slr1668 genes encoding a hypothetical protein and an unknown protein, respectively, were downregulated after transfer to pH 10 and pH 7.5 (Table 2). These genes may be regulated through 3',5'-cyclic AMP (cAMP) signaling as the levels of both transcripts were decreased in a strain lacking a cAMP receptor protein encoded by *syrcp1* (sll1371) (61). In addition, the adenyl cyclase Cya1 (Slr1991), which synthesizes cAMP, is activated by  $CO_2$  (11). This raises the possibility that Cya1 may sense the altered  $HCO_3^-/CO_2$  ratio, resulting in the downregulation of slr1667 and slr1668.

Genes in the cluster consisting of sll1077 to sll1082 were upregulated for the first 2 h of the transition to pH 10, and sll1077, sll1078, and sll1079 were upregulated after transfer to pH 7.5 (Fig. 1b and Table 2). The sll1077 gene product is annotated as an agmatinase (EC 3.5.3.11), an enzyme that catalyzes the conversion of agmatine to putrescine and urea. The polyamine putrescine forms a necessary component of the outer membrane of some gram-negative bacteria during biofilm formation (35). The sll1078 and sll1079 genes were annotated as genes encoding hydrogenase formation proteins. However, deletion of these proteins did not alter hydrogenase activity, leading to the suggestion that they are metallochaperones of the protein encoded by sll1077 and not hydrogenases (14). The sll1080, sll1081, and sll1082 genes are predicted to encode the substrate-binding, permease, and ATP-binding subunits of an ABC transporter, respectively. The coordinated regulation of genes sll1077 to sll1082 at pH 10 suggests that this transporter may be involved in putrescine transport to the outer membrane.

**Validation of microarray results.** The microarray data were validated by semiquantitative reverse transcription-PCR, and good correspondence was observed for all genes examined (see Fig. S1 in the supplemental material).

**Summary.** Figure 2 illustrates the upregulation of transcripts encoding structural proteins that may be involved in the maintenance of cellular homeostasis following a pH transition. The response of *Synechocystis* sp. strain PCC 6803 to high pH had similarities to the response reported for other bacteria. This included the pH 10 upregulation of genes encoding two cation/ $H^+$  antiporters (NhaS3 and Mrp), ATP synthase (also upregulated after transfer to pH 7.5), and at least one amino acid




FIG. 2. Model of a *Synechocystis* sp. strain PCC 6803 cell showing the transcriptional response to pH change. Genes upregulated after transfer to pH 10 are indicated by dark green, and genes upregulated after transfer to both pH 10 and pH 7.5 are indicated by light green. Genes that are not differentially regulated are indicated by open boxes (e.g., Rubisco). The upregulated genes include the genes that encode two cation/proton antiporters (NhaS3 and Mrp), a putative chloride extrusion protein (Slr0753), a mechanosensitive channel that may also act as a calcium channel (MscL), a hexose/proton symporter (GlcP), and a water channel (ApqZ). In addition, genes encoding subunits of NADH dehydrogenase (NDH) and PSII, including the oxygen evolving center (OEC), are upregulated. PGA, phosphoglyceric acid; Cyt  $b_6/f$ , cytochrome  $b_6/f$  complex.

deaminase. One important difference between the response of *Synechocystis* sp. strain PCC 6803 and the response of other bacteria may be related to the lack of transcript level changes for genes involved in the cell surface; however, the cluster containing the agmatinase gene was upregulated, indicating that the cell may produce polyamines destined for the outer membrane. Furthermore, the cell envelope of cyanobacteria has characteristics associated with both gram-negative and gram-positive bacteria, as well as cyanobacterium-specific characteristics, which result in distinct cell wall properties (15). In addition to the general bacterial response, we identified other transporters that may be involved in maintaining pH and ion homeostasis following a transition to pH 10. These include a putative chloride extrusion protein (Slr0753), a mechanosensitive channel that may also act as a calcium channel (MscL), a hexose/proton symporter (GlcP) (Fig. 2), and several ABC transporter subunits for unidentified substrates that are not shown in Fig. 2. A major result of these changes is that they permit the cell to more readily dissipate a buildup of protons in the cytoplasm.

The most important cyanobacterium-specific findings were related to transcriptional changes involved in the maintenance of photosynthetic capability. This study focused on NADH dehydrogenases and the carbon-concentrating mechanism, including the genes encoding carboxysome components and carbonic anhydrase (Fig. 2). In cyanobacteria, changes in the external pH alter the intracellular pH, and an increase in the external pH of 2 pH units results in an increase of  $\sim 0.2$  pH unit in both the cytosol and the thylakoid lumen (2). Such changes

alter the  $\text{CO}_2/\text{HCO}_3^-$  ratio within the cell, and regulation of  $\text{CO}_2/\text{HCO}_3^-$  concentration is essential for maintaining the carboxylase activity of Rubisco. The upregulation of three inducible bicarbonate transporters (BCT1, SbtA, and NDH-1S) and many of the transcripts encoding the structural components of the carboxysome after transfer to pH 7.5 and pH 10 may reflect a response to this perturbed ratio. The pH 10-specific regulation may be a response to increased external pH that decreases  $\text{CO}_2$  levels in the cell (e.g., upregulation of the transcripts encoding the carboxysome  $\beta$ -type carbonic anhydrase and the water channel protein [ApqZ] that has been implicated in  $\text{CO}_2$  import).

The transcriptional response to the transition from pH 7.5 to pH 10 was not the same as the response to  $\text{C}_i$  limitation. The levels of a number of genes that were upregulated under  $\text{C}_i$  limitation conditions were unchanged by the pH transition, and a number of genes that were downregulated by  $\text{C}_i$  were upregulated by the pH transition (e.g., many of the genes encoding low-molecular-weight PSII polypeptides [Table 3]) (58). The ATP synthase,  $\beta$ -type carbonic anhydrase, and ribosomal genes were downregulated at low  $\text{C}_i$  levels but upregulated at pH 10. In contrast, the flavoprotein-encoding transcripts slr0217 and slr0219 were upregulated at low  $\text{C}_i$  levels but were downregulated at high pH. These differences may reflect the decreased growth that was observed after transition to a low  $\text{C}_i$  level but that was not observed at high pH. The pH change had another impact specific to photosynthesis, namely, the pH 10 upregulation of genes encoding the extrinsic and low-molecular-weight

intrinsic proteins of PSII, including the *psbO* and *psbU* transcripts. This was surprising as photoautotrophic growth of a  $\Delta$ PsbO: $\Delta$ PsbU strain at pH 10 but not at pH 7.5 had been interpreted as indicating that PsbO and PsbU may be more important at lower pH. Furthermore, this upregulation was specific to the PSII genes, and very little change in the abundance of the PSI gene transcripts was observed. The mutant lacking luminal proteins may require quantitatively greater enhancement of some of the components highlighted in Fig. 2, and comparisons to examine this possibility will be the objective of future experiments.

#### ACKNOWLEDGMENT

This research was funded by grant DE-FG02-99ER20342 from the Department of Energy.

#### REFERENCES

- Battchikova, N., and E. M. Aro. 2007. Cyanobacterial NDH-1 complexes: multiplicity in function and subunit composition. *Physiol. Plant.* **131**:22–32.
- Belkin, S., and L. Packer. 1988. Determination of pH gradients in intact cyanobacteria by electron spin resonance spectroscopy. *Methods Enzymol.* **167**:677–685.
- Blanco-Rivero, A., F. Leganes, E. Fernandez-Valiente, P. Calle, and F. Fernandez-Pinas. 2005. *mnpA*, a gene with roles in resistance to Na<sup>+</sup> and adaptation to alkaline pH in the cyanobacterium *Anabaena* sp. PCC7120. *Microbiology* **151**:1671–1682.
- Castenholz, R. W. 1988. Culturing methods for cyanobacteria. *Methods Enzymol.* **167**:68–93.
- Codd, G. A., S. G. Bell, K. Kaya, C. J. Ward, K. A. Beattie, and J. S. Metcalf. 1999. Cyanobacterial toxins, exposure routes and human health. *Eur. J. Phycol.* **34**:405–415.
- Colon-Lopez, M. S., D. M. Sherman, and L. A. Sherman. 1997. Transcriptional and translational regulation of nitrogenase in light-dark- and continuous-light-grown cultures of the unicellular cyanobacterium *Cyanothece* sp. strain ATCC 51142. *J. Bacteriol.* **179**:4319–4327.
- de Figueiredo, D. R., M. J. Pereira, A. Moura, L. Silva, S. Barrios, F. Fonseca, I. Henriques, and A. Correia. 2007. Bacterial community composition over a dry winter in meso- and eutrophic Portuguese water bodies. *FEMS Microbiol. Ecol.* **59**:638–650.
- Eaton-Rye, J. J., J. A. Shand, and W. S. Nicoll. 2003. pH-dependent photoautotrophic growth of specific photosystem II mutants lacking luminal extrinsic polypeptides in *Synechocystis* PCC 6803. *FEBS Lett.* **543**:148–153.
- Flores, E., and G. Schmetterer. 1986. Interaction of fructose with the glucose permease of the cyanobacterium *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **166**:693–696.
- Foster, J. S., A. K. Singh, L. J. Rothschild, and L. A. Sherman. 2007. Growth-phase dependent differential gene expression in *Synechocystis* sp. strain PCC 6803 and regulation by a group 2 sigma factor. *Arch. Microbiol.* **187**:265–279.
- Hammer, A., D. R. W. Hodgson, and M. J. Cann. 2006. Regulation of prokaryotic adenyllyl cyclases by CO<sub>2</sub>. *Biochem. J.* **396**:215–218.
- Hiramatsu, T., K. Kodama, T. Kuroda, T. Mizushima, and T. Tsuchiya. 1998. A putative multisubunit Na<sup>+</sup>/H<sup>+</sup> antiporter from *Staphylococcus aureus*. *J. Bacteriol.* **180**:6642–6648.
- Hishiyama, S., W. Hatakeyama, Y. Mizota, N. Hosoya-Matsuda, K. Motohashi, M. Ikeuchi, and T. Hisabori. 2008. Binary reducing equivalent pathways using NADPH-thioredoxin reductase and ferredoxin-thioredoxin reductase in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Plant Cell Physiol.* **49**:11–18.
- Hoffmann, D., K. Gutekunst, M. Klissenbauer, R. Schulz-Friedrich, and J. Appel. 2006. Mutagenesis of hydrogenase accessory genes of *Synechocystis* sp. PCC 6803. Additional homologues of hypA and hypB are not active in hydrogenase maturation. *FEBS J.* **273**:4516–4527.
- Hoiczky, E., and A. Hansel. 2000. Cyanobacterial cell walls: news from an unusual prokaryotic envelope. *J. Bacteriol.* **182**:1191–1199.
- Inaba, M., A. Sakamoto, and N. Murata. 2001. Functional expression in *Escherichia coli* of low-affinity and high-affinity Na<sup>+</sup>(Li<sup>+</sup>)/H<sup>+</sup> antiporters of *Synechocystis*. *J. Bacteriol.* **183**:1376–1384.
- Ito, M., A. A. Guffanti, B. Oudega, and T. A. Krulwich. 1999. *mnp*, a multi-gene, multifunctional locus in *Bacillus subtilis* with roles in resistance to cholate and to Na<sup>+</sup> and in pH homeostasis. *J. Bacteriol.* **181**:2394–2402.
- Jacoby, J. M., D. C. Collier, E. B. Welch, F. J. Hardy, and M. Crayton. 2000. Environmental factors associated with a toxic bloom of *Microcystis aeruginosa*. *Can. J. Fish. Aquat. Sci.* **57**:231–240.
- Kaneko, T., S. Sato, H. Kotani, A. Tanaka, E. Asamizu, Y. Nakamura, N. Miyajima, M. Hirose, M. Sugiura, S. Sasamoto, T. Kimura, T. Hosouchi, A. Matsuno, A. Muraki, N. Nakazaki, K. Naruo, S. Okumura, S. Shimpo, C. Takeuchi, T. Wada, A. Watanabe, M. Yamada, M. Yasuda, and S. Tabata. 1996. Sequence analysis of the genome of the unicellular cyanobacterium *Synechocystis* sp. strain PCC6803. II. Sequence determination of the entire genome and assignment of potential protein-coding regions. *DNA Res.* **3**:109–136.
- Kanesaki, Y., H. Yamamoto, K. Paithoonrangsarit, M. Shoumskaya, I. Suzuki, H. Hayashi, and N. Murata. 2007. Histidine kinases play important roles in the perception and signal transduction of hydrogen peroxide in the cyanobacterium, *Synechocystis* sp. PCC 6803. *Plant J.* **49**:313–324.
- Kaplan, A., and L. Reinhold. 1999. CO<sub>2</sub> concentrating mechanisms in photosynthetic microorganisms. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **50**:539–570.
- Keren, N., H. Ohkawa, E. A. Welsh, M. Liberton, and H. B. Pakrasi. 2005. Psb29, a conserved 22-kD protein, functions in the biogenesis of photosystem II complexes in *Synechocystis* and *Arabidopsis*. *Plant Cell* **17**:2768–2781.
- Kloda, A., E. Petrov, G. R. Meyer, T. Nguyen, A. C. Hurst, L. Hool, and B. Martinac. 2008. Mechanosensitive channel of large conductance. *Int. J. Biochem. Cell Biol.* **40**:164–169.
- Kobayashi, M., H. Katoh, and M. Ikeuchi. 2006. Mutations in a putative chloride efflux transporter gene suppress the chloride requirement of photosystem II in the cytochrome c550-deficient mutant. *Plant Cell Physiol.* **47**:799–804.
- Kunert, A., J. Vinnemeier, N. Erdmann, and M. Hagemann. 2003. Repression by Fur is not the main mechanism controlling the iron-inducible *istAB* operon in the cyanobacterium *Synechocystis* sp. PCC 6803. *FEMS Microbiol. Lett.* **227**:255–262.
- Li, H., A. K. Singh, L. M. McIntyre, and L. A. Sherman. 2004. Differential gene expression in response to hydrogen peroxide and the putative PerR regulon of *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **186**:3331–3345.
- Lopez-Archilla, A. I., D. Moreira, P. Lopez-Garcia, and C. Guerrero. 2004. Phytoplankton diversity and cyanobacterial dominance in a hypereutrophic shallow lake with biologically produced alkaline pH. *Extremophiles* **8**:109–115.
- Maestri, O., and F. Joset. 2000. Regulation by external pH and stationary growth phase of the acetolactate synthase from *Synechocystis* PCC6803. *Mol. Microbiol.* **37**:828–838.
- Meunier, P. C., M. S. Colon-Lopez, and L. A. Sherman. 1997. Temporal changes in state transitions and photosystem organization in the unicellular, diazotrophic cyanobacterium *Cyanothece* sp. ATCC 51142. *Plant Physiol.* **115**:911–1000.
- Mogelhoj, M. K., P. J. Hansen, P. Henriksen, and N. Lundholm. 2006. High pH and not allelopathy may be responsible for negative effects of *Nodularia spumigena* on other algae. *Aquat. Microb. Ecol.* **43**:43–54.
- Nazarenko, L. V., I. M. Andreev, A. A. Lyukevich, T. V. Pisareva, and D. A. Los. 2003. Calcium release from *Synechocystis* cells induced by depolarization of the plasma membrane: MscL as an outward Ca<sup>2+</sup> channel. *Microbiology* **149**:1147–1153.
- Ohkawa, H., H. B. Pakrasi, and T. Ogawa. 2000. Two types of functionally distinct NAD(P)H dehydrogenases in *Synechocystis* sp. strain PCC6803. *J. Biol. Chem.* **275**:31630–31634.
- Padan, E., E. Bibi, M. Ito, and T. A. Krulwich. 2005. Alkaline pH homeostasis in bacteria: new insights. *Biochim. Biophys. Acta* **1717**:67–88.
- Paithoonrangsarit, K., M. A. Shoumskaya, Y. Kanesaki, S. Satoh, S. Tabata, D. A. Los, V. V. Zinchenko, H. Hayashi, M. Tanticharoen, I. Suzuki, and N. Murata. 2004. Five histidine kinases perceive osmotic stress and regulate distinct sets of genes in *Synechocystis*. *J. Biol. Chem.* **279**:53078–53086.
- Patel, C. N., B. W. Wortham, J. L. Lines, J. D. Fetherston, R. D. Perry, and M. A. Oliveira. 2006. Polyamines are essential for the formation of plague biofilm. *J. Bacteriol.* **188**:2355–2363.
- Pikuta, E. V., R. B. Hoover, and J. Tang. 2007. Microbial extremophiles at the limits of life. *Crit. Rev. Microbiol.* **33**:183–209.
- Poncelet, M., C. Cassier-Chauvat, X. Leschelle, H. Bottin, and F. Chauvat. 1998. Targeted deletion and mutational analysis of the essential (2Fe-2S) plant-like ferredoxin in *Synechocystis* PCC6803 by plasmid shuffling. *Mol. Microbiol.* **28**:813–821.
- Postier, B. L., H. L. Wang, A. Singh, L. Impson, H. L. Andrews, J. Klahn, H. Li, G. Risinger, D. Pesta, M. Deyholos, D. W. Galbraith, L. A. Sherman, and R. L. Burnap. 2003. The construction and use of bacterial DNA microarrays based on an optimized two-stage PCR strategy. *BMC Genomics* **4**:23.
- Reddy, K. J., R. Webb, and L. A. Sherman. 1990. Bacterial RNA isolation with one hour centrifugation in a table-top ultracentrifuge. *BioTechniques* **8**:250–251.
- Ritchie, R. J. 1991. Membrane-potential and pH control in the cyanobacterium *Synechococcus* R-2 (*Anacystis nidulans*) PCC 7942. *J. Plant Physiol.* **137**:409–418.
- Shapiguzov, A., A. A. Lyukevich, S. I. Allakhverdiev, T. V. Sergeev, I. Suzuki, N. Murata, and D. A. Los. 2005. Osmotic shrinkage of cells of *Synechocystis* sp. PCC 6803 by water efflux via aquaporins regulates osmotic-stress-inducible gene expression. *Microbiology* **151**:447–455.
- Shapiro, J. 1973. Blue-green algae: why they become dominant. *Science* **179**:382–384.

43. Sherman, D. M., T. A. Troyan, and L. A. Sherman. 1994. Localization of membrane proteins in the cyanobacterium *Synechococcus* sp. PCC7942 (radial asymmetry in the photosynthetic complexes). *Plant Physiol.* **106**:251–262.
44. Shi, L. X., and W. P. Schroder. 2004. The low molecular mass subunits of the photosynthetic supracomplex, photosystem II. *Biochim. Biophys. Acta* **1608**:75–96.
45. Shoumskaya, M. A., K. Paithoonrangsarid, Y. Kanesaki, D. A. Los, V. V. Zinchenko, M. Tanticharoen, I. Suzuki, and N. Murata. 2005. Identical Hik-Rre systems are involved in perception and transduction of salt signals and hyperosmotic signals but regulate the expression of individual genes to different extents in *Synechocystis*. *J. Biol. Chem.* **280**:21531–21538.
46. Singh, A. K., H. Li, and L. A. Sherman. 2004. Microarray analysis and redox control of gene expression in the cyanobacterium *Synechocystis* sp. PCC 6803. *Physiol. Plant.* **120**:27–35.
47. Singh, A. K., L. M. McIntyre, and L. A. Sherman. 2003. Microarray analysis of the genome-wide response to iron deficiency and iron reconstitution in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Physiol.* **132**:1825–1839.
48. Singh, A. K., and L. A. Sherman. 2002. Characterization of a stress-responsive operon in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Gene* **297**:11–19.
49. Singh, A. K., and L. A. Sherman. 2005. Pleiotropic effect of a histidine kinase on carbohydrate metabolism in *Synechocystis* sp. strain PCC 6803 and its requirement for heterotrophic growth. *J. Bacteriol.* **187**:2368–2376.
50. Singh, A. K., T. C. Summerfield, H. Li, and L. A. Sherman. 2006. The heat shock response in the cyanobacterium *Synechocystis* sp. strain PCC 6803 and regulation of gene expression by HrcA and SigB. *Arch. Microbiol.* **186**:273–286.
51. So, A. K., and G. S. Espie. 1998. Cloning, characterization and expression of carbonic anhydrase from the cyanobacterium *Synechocystis* PCC6803. *Plant Mol. Biol.* **37**:205–215.
52. Stancik, L. M., D. M. Stancik, B. Schmidt, D. M. Barnhart, Y. N. Yoncheva, and J. L. Slonczewski. 2002. pH-dependent expression of periplasmic proteins and amino acid catabolism in *Escherichia coli*. *J. Bacteriol.* **184**:4246–4258.
53. Summerfield, T. C., J. Eaton-Rye, and L. A. Sherman. 2007. Global gene expression of a  $\Delta$ PsbO: $\Delta$ PsbU mutant and a spontaneous revertant in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Photosynth. Res.* **94**:265–274.
54. Summerfield, T. C., J. A. Shand, F. K. Bentley, and J. J. Eaton-Rye. 2005. PsbQ (Sll1638) in *Synechocystis* sp. PCC 6803 is required for photosystem II activity in specific mutants and in nutrient-limiting conditions. *Biochemistry* **44**:805–815.
55. Summerfield, T. C., and L. A. Sherman. 2007. Role of sigma factors in controlling global gene expression in light/dark transitions in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **189**:7829–7840.
56. Tchernov, D., Y. Helman, N. Keren, B. Luz, I. Ohad, L. Reinhold, T. Ogawa, and A. Kaplan. 2001. Passive entry of CO<sub>2</sub> and its energy-dependent intracellular conversion to HCO<sub>3</sub><sup>-</sup> in cyanobacteria are driven by a photosystem I-generated DmH<sup>+</sup>. *J. Biol. Chem.* **276**:23450–23455.
57. Ushimaru, T., Y. Nishiyama, H. Hayashi, and N. Murata. 2002. No coordinated transcriptional regulation of the *sod-kat* antioxidative system in *Synechocystis* sp. PCC 6803. *J. Plant Physiol.* **159**:805–807.
58. Wang, H. L., B. L. Postier, and R. L. Burnap. 2004. Alterations in global patterns of gene expression in *Synechocystis* sp. PCC 6803 in response to inorganic carbon limitation and the inactivation of *ndhR*, a LysR family regulator. *J. Biol. Chem.* **279**:5739–5751.
59. Wang, H. L., B. L. Postier, and R. L. Burnap. 2002. Polymerase chain reaction-based mutageneses identify key transporters belonging to multigene families involved in Na<sup>+</sup> and pH homeostasis of *Synechocystis* sp. PCC 6803. *Mol. Microbiol.* **44**:1493–1506.
60. Williams, J. G. K. 1988. Construction of specific mutations in photosystem II photosynthetic reaction center by genetic engineering methods in *Synechocystis* 6803. *Methods Enzymol.* **167**:766–778.
61. Yoshimura, H., S. Yanagisawa, M. Kanehisa, and M. Ohmori. 2002. Screening for the target gene of cyanobacterial cAMP receptor protein SYCRP1. *Mol. Microbiol.* **43**:843–853.
62. Zhang, P., N. Battchikova, T. Jansen, J. Appel, T. Ogawa, and E. M. Aro. 2004. Expression and functional roles of the two distinct NDH-1 complexes and the carbon acquisition complex NdhD3/NdhF3/CupA/Sll1735 in *Synechocystis* sp. PCC 6803. *Plant Cell* **16**:3326–3340.