

6-26-2013

Ultradian Metabolic Rhythm in the Diazotrophic Cyanobacterium *Cyanothece* sp. ATCC 51142.

Jan Cervený

Maria A. Sinetova

Luis Valledor

Louis A. Sherman

Purdue University, lsherman@purdue.edu

Ladislav Nebal

Follow this and additional works at: <http://docs.lib.purdue.edu/bioscipubs>

Recommended Citation

Cervený, Jan; Sinetova, Maria A.; Valledor, Luis; Sherman, Louis A.; and Nebal, Ladislav, "Ultradian Metabolic Rhythm in the Diazotrophic Cyanobacterium *Cyanothece* sp. ATCC 51142." (2013). *Department of Biological Sciences Faculty Publications*. Paper 49. <http://dx.doi.org/10.1073/pnas.1301171110>

1 *Classification:*

2 *BIOLOGICAL SCIENCES: Cell Biology*

3 **Title: Ultradian metabolic rhythm in the diazotrophic cyanobacterium *Cyanothece* sp.**

4 **ATCC 51142**

5 *Short title:*

6 Ultradian metabolic rhythm in *Cyanothece*

7 *Authors:*

8 Jan Červený^{a,2}, Maria A. Sinetova^{a,b,2}, Luis Valledor^{a,c}, Louis A. Sherman^d, and Ladislav

9 Nedbal^{a,e,1}

10 *Author affiliation:*

11 ^a Global Change Research Centre - CzechGlobe, Academy of Sciences of the Czech Republic,

12 Drásov 470, 664 24 Drásov, Czech Republic

13 ^b Laboratory of Intracellular Regulation, Institute of Plant Physiology, Russian Academy of

14 Sciences, Botanicheskaya street 35, 127276, Moscow, Russian Federation

15 ^c Molecular Systems Biology MOSYS, University of Vienna, Vienna, Austria

16 ^d Department of Biological Sciences, Purdue University, West Lafayette, IN 47907, USA

17 ^e Systems Biology Laboratory, Faculty of Informatics, Masaryk University, Brno, Czech

18 Republic

19 ¹ To whom correspondence should be addressed: nedbal.lad@gmail.com

20 ² JC and MS contributed equally to this work

21 *Corresponding author:*

22 Ladislav Nedbal, Global Change Research Centre - CzechGlobe, Academy of Sciences of the

23 Czech Republic, Drásov 470, 664 24 Drásov, Czech Republic

24 Ph. +420 606 690028, e-mail: nedbal.lad@gmail.com

- 1 *Keywords:* cyanobacteria, nitrogen fixation, oscillation, photosynthesis, respiration, ultradian
- 2 rhythm
- 3

1 **Abstract**

2 The unicellular cyanobacterium *Cyanothece* sp. ATCC 51142 is capable of performing oxygenic
3 photosynthesis during the day and microoxic nitrogen fixation at night. These mutually exclusive
4 processes are possible only by temporal separation by circadian clock or another cellular
5 program. We report an identification of a temperature-dependent ultradian metabolic rhythm that
6 controls the alternating oxygenic and microoxic processes of *Cyanothece* sp. ATCC 51142 under
7 continuous high irradiance and in high CO₂ concentration. During the oxygenic photosynthesis
8 phase, nitrate deficiency limited protein synthesis and CO₂ assimilation was directed towards
9 glycogen synthesis. The carbohydrate accumulation reduced over-excitation of the
10 photosynthetic reactions until a respiration burst initiated a transition to microoxic N₂-fixation. In
11 contrast to the circadian clock, this ultradian period is strongly temperature-dependent: 17h at
12 27°C, which continuously decreased to 10h at 39°C. The cycle was expressed by an oscillatory
13 modulation of net O₂ evolution, CO₂ uptake, pH, fluorescence emission, glycogen content, cell
14 division, and culture optical density. The corresponding ultradian modulation was also observed
15 in the transcription of nitrogenase-related *nifB* and *nifH* genes and in nitrogenase activities. We
16 propose that the control by the newly identified metabolic cycle adds another rhythmic
17 component to the circadian clock that reflects the true metabolic state depending on the actual
18 temperature, irradiance, and CO₂ availability.

19

1 \body

2 **Introduction**

3 Cyanobacteria helped to form the Earth's biosphere since they first evolved some 3 billion years
4 ago. These microorganisms made our atmosphere oxygenic and their fixation of CO₂ and N₂ has
5 made important contributions to the elemental cycles. In the past decades, cyanobacteria of the
6 genus *Cyanothece* have been attracting a strong attention by their significant contributions to the
7 nitrogen cycle (1), as unique models to study relationship between N₂ fixation, photosynthesis
8 and respiration (2), as well as promising candidates for bioenergy production (3-5). Distinct
9 potential has been found particularly in the strain *Cyanothece* ATCC 51142 (2, 4). *Cyanothece*
10 sp. ATCC 51142 is a unicellular cyanobacterium where spatial compartmentalization of the
11 mutually exclusive oxygenic photosynthesis and microoxic nitrogen fixation is impossible (6).
12 The strategy used by this organism is to temporally separate the molecular oxygen released by
13 photosynthesis from the nitrogenase that would otherwise be irreversibly O₂-inactivated. The
14 capacity to separate the antagonistic metabolic processes in time is usually attributed to circadian
15 control. The circadian clock in cyanobacteria relies on cyclic (de-) phosphorylation involving
16 complexes of the KaiA, KaiB, KaiC proteins (7-9). The clock mechanism has been studied to a
17 great detail in *Synechococcus elongatus* PCC 7942 and also *in vitro* (10). The clock period in this
18 organism has been shown to be temperature compensated – a feature essential for controlling the
19 daily rhythm particularly in organisms that do not sustain a stable temperature for their
20 metabolism (11).

21 In *Cyanothece* sp. ATCC 51142, the *kai* genes exist in multiple copies (12) although the Kai
22 proteins have not been studied in the detail achieved for *S. elongatus*. The daily modulation of
23 the metabolic activity and gene transcription, namely alternation of photosynthetic and N₂-

1 fixation phases, has been attributed to the control by the circadian clock largely because the
2 observed period was approximately 24 h, even in continuous light following a 12 h light / 12 h
3 dark entrainment (13). Certain genes, especially those associated with nitrogenase assembly and
4 function, demonstrated a circadian-type of regulation when the light-dark pattern was modified
5 for growth under continuous light or short day-night periods (14).

6 However, there are no published data on *Cyanothece* sp. ATCC 51142 to show that the period
7 of the oscillatory modulation in continuous light is temperature compensated as expected for
8 circadian control. One can expect an involvement of metabolic processes (see also (2)) that may
9 result in distinct ultradian rhythms such as known in yeast, which occur with temperature-
10 dependent periods that are significantly shorter than 24 h. The hypothesis suggesting an
11 involvement of an ultradian metabolic cycle in *Cyanothece* sp. ATCC 51142 is supported by
12 oscillations with ca. 12 h periods that occur in continuous light following an initial 12 h light/ 12
13 h dark entrainment (15) as well as in continuous light in a batch culture (16). Components of
14 about 12 h period have also been detected by a Fourier transform analysis in transcript data in
15 continuous light following a 12 h light / 12 h dark entrainment (17).

16 In a search for a potential ultradian metabolic rhythm in *Cyanothece* sp. ATCC 51142, we
17 have performed experiments similar to those in yeast in which ultradian oscillations were
18 induced by a starvation period (18). *Cyanothece* sp. ATCC 51142 was grown to late exponential
19 or linear phase in regular medium containing nitrate and supplied with saturating CO₂ and light.
20 Strong ultradian oscillations occurred after the cells were moved to a minus-nitrate medium. The
21 newly found ultradian metabolic rhythm is strongly temperature-dependent. We also show that
22 the circadian cycle is well temperature-compensated. The contrasting temperature dependence
23 documents that the ultradian and circadian cycles are independent. The ultradian rhythm

1 dominates in saturating CO₂ and light while the circadian rhythm prevails when irradiance and/or
2 CO₂ concentration are lowered.

3 **Results**

4 *Ultradian and circadian rhythms in a diurnally entrained culture.* *Cyanothece* sp. ATCC 51142
5 (hereafter *Cyanothece*) was grown in flat-panel photobioreactors with highly time-resolved,
6 automated sampling to follow cyclic processes over days and weeks (19). Before the experiment
7 shown in Fig.1, the culture was entrained in 12 h light / 12 h dark cycles in a turbidostat mode in
8 which the culture optical density was kept in a narrow range by a photobioreactor-controlled
9 feedback dilution. The experiment started after the last period of the diurnal entrainment (interval
10 0-24h in Fig.1) by switching off the dilution allowing the culture batch growth and by keeping
11 the culture in continuous light for the subsequent 10 days (interval 24 - 264h in Fig.1). In
12 response to the pre-treatment, the culture exhibited a complex oscillatory pattern in which both,
13 the shorter ultradian and the longer circadian rhythms combined. The periods in which the
14 ultradian rhythm dominated (see the 72-96 h interval in the bottom left panel) were followed by
15 periods dominated by the circadian rhythm (see the 216-264 h interval in the bottom right panel).

16 The concentration of dissolved oxygen oscillated with a minimum in the first half of
17 subjective night in the circadian phase (216-264 h, see open circles in the lower right panel in
18 Fig. 1). The minimum in the concentration of the dissolved oxygen during the ultradian phase
19 (approx. 48–144 h, see the open circles in the lower left panel in Fig. 1) was not firmly anchored
20 to the subjective day/night cycle, because its period was neither equal to 12 h nor to 24 h. The
21 minima of the dissolved oxygen preceded by 1 to 2 hours the maxima of the dissolved CO₂
22 concentration (thin full line in Fig. 1). The oscillations of the medium pH (short-dashed line in
23 the lower panels of Fig. 1) were nearly anti-parallel to the dissolved CO₂ concentration, as

1 anticipated from the water chemistry of inorganic carbon forms. During the rapid growth (lower
2 left panel in Fig. 1) the optical density at 680 nm of the suspension (OD_{680}) was weakly
3 modulated with a period of the ultradian rhythm so that the maxima of the dissolved oxygen
4 concentration matched the steepest increase of the optical density. This suggests that high
5 photosynthetic activity indicated by the high concentration of dissolved O_2 correlated with an
6 increase in chlorophyll concentration represented by the surrogate OD_{680} . The maxima of the
7 stationary pigment fluorescence emission (the long-dashed line in the lower panels in Fig. 1)
8 approximately coincided with the O_2 minima and with the CO_2 maxima, a pattern that was not
9 always reproduced in parallel experiments. This dynamic pattern can be tentatively interpreted as
10 high fluorescence due to a reduction of the plastoquinone pool by high rates of respiration and
11 low rates of photosynthesis and *vice versa*.

12

1 ***Metabolic rhythms in a diurnally non-entrained culture.*** A complex oscillatory pattern similar
2 to the diurnally entrained culture (Fig.1) was also observed in continuous light, i.e. without the
3 diurnal entrainment, when cyanobacteria rapidly growing in a nitrate-rich medium were
4 transferred to a medium without nitrate (time 0 in Fig. 2). The dynamic pattern during the first
5 3 days following the media replacement exhibited a dominant ultradian character (period
6 ~10.5 h). As the culture grew ($OD_{680} \sim 2.2$), the oscillations became dominated by the circadian
7 rhythm (lower right panel in Fig. 2). Note that the residual involvement of the ultradian
8 dynamics remained also in the dense culture, as demonstrated by the two high amplitudes at
9 160.5th and 234th hours of the experiment. The time period separating the two high amplitudes
10 corresponded to 7 ultradian periods ($7 \times 10.5 \text{ h} = 73.5 \text{ h}$) and to, roughly, 3 circadian periods
11 ($3 \times 24 \text{ h} = 72 \text{ h}$). Thus, at every 7th ultradian and every 3rd circadian period, the respective
12 amplitudes combined rather than cancelled each other.

13 We argue that the transition from the dominant ultradian to circadian rhythm in Fig. 2 was
14 induced by decreasing the mean effective irradiance in the dense culture. This hypothesis is
15 supported by the results in Fig. 3 in which the culture density was stabilized by the turbidostat
16 that was diluting the suspension to the same optical density and, thus, the mean irradiance was
17 fully controlled by the photobioreactor to high irradiance in the initial days and to low irradiance
18 towards the end of the experiment. The data demonstrate that the ultradian rhythm can be
19 converted to circadian cycle by lowering of the incident irradiance in the photobioreactor. This
20 result implies a question what is the decisive factor controlling the transition of the ultradian
21 phenomena to the circadian rhythm: Is it light as the result in Fig. 3 suggests, CO₂ availability, or
22 is it fast growth that requires both, high light and high CO₂? In Supplementary information (*SI*
23 *Appendix*, Fig. S1 and Fig. S2), we suggest that high light and high CO₂, rather than fast growth,
24 lead to the dominating ultradian rhythm. Another condition for the ultradian rhythm to occur is

1 lack of nitrate in the medium. An addition of nitrate quickly damped the ultradian oscillations to
2 zero (*SI Appendix*, Fig. S3). Also, the ultradian oscillations in Fig. 2 were induced by transferring
3 the cells to a nitrate-free medium.

4 Temperature is another important factor affecting the ultradian metabolic rhythm. The
5 oscillations shown in Figs. 1-3 occurred at 36 °C with an ultradian period of 10.55 ± 0.05 hour
6 (mean and standard deviation of 5 biological replicates). We found that the period of ultradian
7 oscillations was strongly temperature dependent. It increased to 17 hours at 27 °C and decreased
8 to 10 hours at 39 °C (Fig. 4). The temperature coefficient Q_{10} was 1.55, documenting the
9 ultradian temperature dependence. We also measured the periods of the 24h rhythms in different
10 temperatures and found Q_{10} close to 1, typical for the temperature-compensated circadian
11 rhythm. The different response to temperature change shows that the ultradian rhythm is truly
12 independent phenomenon that is not derived from the circadian clock.

13 Data in Fig. 5A demonstrate the ultradian oscillations of dissolved O₂ concentration, pH and
14 partial synchronization of cell division (bars). The increasing dissolved oxygen and rising pH
15 (i.e., decreasing dissolved CO₂) reflected active photosynthesis leading to an accumulation of
16 biomass carbon between ca. 7th and 13th hour (closed squares in Fig. 5B). A large part of the
17 organic carbon was stored in form of semi-amylopectin granules (20), frequently referred to as
18 glycogen (open squares in the Fig. 5B). The proportion of glycogen carbon to total biomass
19 carbon increased during the photosynthesis phase, likely due to the lack of nitrogen that limited
20 protein synthesis. The imbalance between the surplus of fixed carbon and the lack of nitrate in
21 the medium grew to a point where photosynthesis was down-regulated, the dissolved oxygen
22 and, ca. 2 h later, also the pH started decreasing and the accumulation of organic carbon was
23 suspended. The glycogen content started decreasing and oxygen dropped to a level allowing
24 nitrogen fixation. Nitrogen fixation dominated between ca. 3rd and 7th hour and between 13.5th

1 and 18th hour of the experiment – periods indicated by the dashed line rectangle surrounding the
2 respective segments in Fig. 5. The rising nitrogen content of the biomass is indicated by the
3 dashed line and full circles in Fig. 5C. The genes *nifH* and *nifB* necessary for nitrogen fixation
4 are expressed to higher mRNA levels in the early phase of the nitrogen fixation periods.

5

6 **Discussion**

7 *Linking the ultradian metabolic rhythm to other cyclic processes.* This paper has
8 demonstrated that *Cyanothece* sp. ATCC 51142 displays ultradian rhythms of alternating
9 photosynthesis and N₂-fixation when grown without nitrate in continuous culture conditions.
10 Whether the cultures were entrained (Fig. 1) or not (Fig. 2), the exponential growth phase
11 showed ultradian rhythms that eventually degenerated into more circadian-like rhythms. This
12 finding ties together many disparate results for *Cyanothece* sp. ATCC 51142 over the years and
13 may provide a basis for the interrelationship of the circadian and ultradian types of temporal
14 regulation.

15 The earlier experiments typically used low light / low CO₂ and showed circadian patterns
16 without an obvious ultradian modulation (3, 21). Here, we demonstrated that the ultradian
17 rhythm occurs when photosynthesis is saturated by high light and high CO₂ concentration. In
18 respect to high carbon availability, it is important to note that an ultradian component was also
19 observed over a circadian pattern when *Cyanothece* was grown heterotrophically with glycerol
20 (22).

21 Both, the ultradian and circadian cycles exhibit many similar features. During the
22 photosynthesis phase, cells accumulate large amounts of glycogen. The transition from
23 photosynthetic to nitrogen fixation phase correlate with degradation of glycogen and a

1 respiration burst. When enough ATP and NADH/NADPH are accumulated and intracellular
2 oxygen is lowered by respiration, the cells start fixing nitrogen.

3 Earlier studies of *Cyanothece* sp. ATCC 51142 have been performed mostly on 12 h light/
4 12 h dark grown cells and also on cells grown under continuous light and short light/dark periods
5 (21). Invariably, the nitrogenase genes in the 35-gene cluster demonstrated the most consistent
6 circadian behavior (13, 14, 21). Interestingly, the nitrogenase genes were among the large
7 majority of genes displaying a circadian rhythm even in experiments in which some other genes
8 oscillated with a 12h period. The observation of gene oscillation with a 12h period led to an
9 earlier proposal of an ultradian rhythm (17). Here, we probed expression of the *nifB* and *nifH*
10 genes and both were clearly modulated by the ultradian rhythm, a feature consistent also with the
11 ultradian modulation of the nitrogenase activity (Fig. 5). The ultradian modulation of the
12 nitrogenase genes supports the notion that the ultradian rhythm is, under favorable conditions,
13 dominant in all aspects attributed earlier exclusively to circadian control. In this sense, it is
14 highly relevant to ask if the ultradian and circadian processes are not reflecting a single control
15 mechanism (23).

16 To clarify the interrelationship between the ultradian and circadian rhythms, we measured
17 temperature dependence of the respective periods (Fig. 4). The circadian period was shown to be
18 well temperature-compensated ($Q_{10} \approx 1$) while the ultradian period was found to be strongly
19 temperature-dependent ($Q_{10} = 1.55$). We conclude that the ultradian metabolic oscillation
20 represents a process that is truly independent from the circadian cycle.

21 The ultradian rhythms were replaced or dominated by circadian rhythms when the cell
22 irradiance was lowed either by self-shading in high cell density (Figs. 1,2) or by dimming the
23 irradiance incident at the photobioreactor cuvette (Fig. 3). In Supplementary information, we

1 show an experiment which supports the conclusion that the ultradian rhythm is largely
2 suppressed when photosynthesis and growth are not saturated by light (*SI Appendix, Fig. S1A*).

3 The ultradian rhythm may also depend on fast growth which requires both, high light and
4 high CO₂. However, we show in (*SI Appendix, Fig. S1B*) that the growth rate is maximal at ca.
5 33°C and declines towards 36°C and 39°C. This is a qualitatively contrasting temperature
6 dependence compared to that of the ultradian period that declines monotonously between 27°C
7 and 39°C. We conclude that the growth rate is not a driver of the ultradian phenomena. This
8 conclusion is further supported by showing (*SI Appendix, Fig. S2*) that the change of the specific
9 growth rate by increasing the sparging gas CO₂ concentration in the range from 700 to 1000 ppm
10 is small compared to the much steeper decrease of the ultradian period.

11 It appears that two independent mechanisms of temporal separation of photosynthesis and
12 nitrogen fixation are possible in *Cyanothece*: the ultradian rhythm that prevails when
13 photosynthesis is saturated by light and CO₂ and the circadian rhythm that dominates when the
14 rate of photosynthesis is limited.

15 Interestingly, the experiments reported here also shed light at an earlier observation that the
16 circadian patterns of photosynthesis and nitrogen fixation often appeared even during the first
17 entrainment period, a feature that was hard to understand without context shown in Fig. 1. The
18 experimental cultures used to be inoculated with batch cultures that were 10-14 days old and,
19 thus, closely resembled ca. >200 h cells in Fig. 1 that were probably already poised for the
20 circadian regime prior to inoculation. It was the new photobioreactor that permitted us to work
21 under a variety of environmental conditions and we can now assimilate the information from
22 these different experiments.

23

1 *On the homology between the metabolic rhythms in Cyanotheca and in Saccharomyces*
2 *cerevisiae.*

3 Aerobically-growing, dense continuous cultures of *Saccharomyces cerevisiae* exhibit
4 autonomous energy-metabolite ultradian oscillations of ca. 2-5 hours period that are observed
5 under low nitrogen and low carbon conditions (18, 24-26). Though yeast are eukaryotic
6 organisms that separate incompatible processes in different compartments, they also use
7 temporal separation to carry out incompatible metabolic reaction and further alleviate futile
8 cycles (27). A difference between the ultradian oscillations in yeast and in *Cyanotheca* relates to
9 the culture density. In contrast to yeast, the *Cyanotheca* ultradian oscillations do not occur in
10 high cell density that leads to self-shading of the penetrating light and to a reduction of the
11 ultradian oscillatory amplitudes. Thus, this difference is only due to the role of light penetration
12 through the culture that is critical for cyanobacteria and irrelevant to yeast.

13 In other aspects, several principal homologous features of metabolic oscillations of
14 *Cyanotheca* sp. ATCC 51142 and *Saccharomyces cerevisiae* can be found in spite of their largely
15 different prokaryotic and eukaryotic character. The yeast oscillations, similarly to the oscillations
16 reported here for *Cyanotheca*, consisted of two distinct phases: the anabolic, carbohydrate-
17 storing phase (reductive in the yeast and photosynthetic in the cyanobacterium) and catabolic
18 phase (oxidative in the yeast and N₂-fixing in the cyanobacterium). The ultradian oscillations of
19 dissolved O₂ and pH (CO₂) were observed in both organisms, accompanied by accumulation of
20 carbohydrates in the anabolic/photosynthetic phase that was concluded by a burst of respiration,
21 see (24) and (18) for yeast. Also, cell division occurred only in the anabolic/photosynthetic
22 phase, see (18) for yeast. In yeast, many genes participating in the regulation of metabolic
23 processes were oscillating with the same ultradian period as the energetic-metabolic oscillations.
24 In *Cyanotheca*, the nitrogenase-related genes *nifB* and *nifH* were also clearly modulated by the

1 ultradian rhythm, a feature consistent with the ultradian modulation of the nitrogenase activity.
2 The homology between the metabolic ultradian oscillations in yeast and in cyanobacteria
3 suggests that both organisms may have the same strategy of benefiting from the ultradian cycling
4 in separating the mutually exclusive metabolic processes, anabolic or photosynthetic and
5 catabolic or nitrogen-fixing reactions, and avoiding futile cycles (18, 27).

6 The finding of the ultradian metabolic cycle *Cyanothece* sp. ATCC 51142 also resonates with
7 the recent report (2) about significant differences between six *Cyanothece* strains subjected to
8 identical external cues. The differences were interpreted as indicating importance of metabolic
9 signals that are specific for each strain rather than pure regulation by the circadian clock. An
10 important question to ask in future research is on the role of such metabolic signals and of the
11 ultradian cycle in natural conditions. *Cyanothece* sp. ATCC 51142 is a benthic organism that
12 lives in mats or clusters on rocks in intertidal areas (28). Ultradian synchronization of alternating
13 oxygenic and microoxic conditions in the *Cyanothece*-containing biofilms can be considered as
14 an potentially important supplement to the circadian control in adjusting to actual temperature,
15 light and, carbon availability in the environment.

16

17 **Materials and Methods**

18 *Growth conditions, real time measurements and sampling.* The *Cyanothece* sp. strain
19 ATCC 51142 (28) was obtained from Jana Stöckel, Washington University, USA and maintained
20 in the complete ASP2 medium (29, 30) buffered with TAPS and TAPSO as described earlier
21 (16).

22 For growing and real time monitoring of cultures, we used the flat panel photobioreactors
23 (FMT-150, Photon Systems Instruments, Brno, Czech Republic), which were described in detail

1 in (19). Unless specified otherwise, the culture temperature was stabilized at 36.0 ± 0.3 °C,
2 irradiance was $130\text{-}160$ $\mu\text{mol}(\text{photons})\text{ m}^{-2}\text{ s}^{-1}$ of red light (≈ 627 nm) supplemented with
3 25 $\mu\text{mol}(\text{photons})\text{ m}^{-2}\text{ s}^{-1}$ of cool white or blue light (≈ 455 nm). High concentration of dissolved
4 CO_2 was generated by sparging with 0.5% CO_2 .

5 For diurnal entrainment, the culture grown previously on a shaker in an Erlenmeyer flask was
6 inoculated in a nitrate-free ASP2 medium in the photobioreactor with alternating 12 h light and
7 12 h dark periods. After the entrainment lasting at least 5 days, the lights were switched on
8 continuously.

9 For ultradian metabolic cycles induced by removal of nitrate from the medium, the culture
10 was first grown in photobioreactors in ASP2 medium with nitrate until late exponential or early
11 linear phase. The cells were harvested by centrifugation and re-suspended in nitrate-free ASP2
12 medium. Cells continued to grow in continuous light without nitrate in batch or in turbidostat
13 mode ($\text{OD}_{680} = 0.7$). Aliquots for cell counting, for quantification of chlorophyll, carbohydrate
14 storage, total carbon and total nitrogen content, and for the measurement of nitrogenase activity
15 were collected every 2-2.5 hours during 24-54 hours. Concentrations of dissolved oxygen and
16 carbon dioxide were measured by the photobioreactor *in situ* every minute as described in (31).

17

18 **Analytical measurements.** Estimation of chlorophyll, carbohydrate, carbon and nitrogen content
19 and cell counts were done as described previously (16).

20 The nitrogenase activity was determined using the standard acetylene reduction method:
21 Culture aliquots of 15 ml were incubated in gas-tight glass vials of total 40 ml volume with 8 ml
22 of acetylene for 2 hours in light and temperature that were used for growing. Then, 4 ml of

1 headspace gas were assayed by gas-chromatography. Potential ethylene contamination of
2 acetylene was checked in 4 ml of gas phase taken immediately after the addition of the acetylene.

3 Quantitative Real Time PCRs were performed in a Mini-Opticon System (Bio-Rad).
4 Normalized Relative Quantities (NRQ) and Standard errors of RQ were determined according to
5 (32). Expression levels of *16S* and *rpnA* genes were used as endogenous controls for normalizing
6 the abundance of *nifB* and *nifH* transcripts. Detailed information about RNA extraction, RT, and
7 PCR conditions is available in *SI Appendix*.

8

9 **Acknowledgements**

10 This publication is an output of the CzechGlobe Centre that is being developed within the OP
11 RDI and co-financed from EU funds and the State Budget of the Czech Republic (Project:
12 CzechGlobe – Centre for Global Climate Change Impacts Studies), Reg. No.
13 CZ.1.05/1.1.00/02.0073 and an output of the EC OP project financed from EU with the support
14 of MEYS CR: Local Team and International Consortium for Computational Modelling of a
15 Cyanobacterial Cell, Reg. No. CZ.1.07/2.3.00/20.0256 (to JC, MS, LV and LN). This work was
16 also supported by research project RFBR 12-04-32148 to MS, by a grant GAČR 206/09/1284 to
17 LN, and by a grant from the US DOE Genomics: GTL program to LS.

18 We thank Kristina Felcmanová for contributions during nitrogenase activity samples analysis.

19

20 **References**

- 21 1. Zehr JP (2011) Nitrogen fixation by marine cyanobacteria. *Trends Microbiol* 19(4):162-
22 173.

- 1 2. Bandyopadhyay A, Elvitigala T, Liberton M, Pakrasi HB (2013) Variations in the rhythms
2 of respiration and nitrogen fixation in members of the unicellular diazotrophic
3 cyanobacterial genus *Cyanothece*. *Plant Physiol* 161(3):1334-1346.
- 4 3. Sherman LA, Min H, Toepel J, Pakrasi HB (2010) in *Recent Advances in Phototrophic*
5 *Prokaryotes*, ed Hallenbeck PC (Springer, New York, NY), pp 275-290.
- 6 4. Bandyopadhyay A, Stöckel J, Min H, Sherman LA, Pakrasi HB (2010) High rates of
7 photobiological H₂ production by a cyanobacterium under aerobic conditions. *Nat*
8 *Commun* 1(9):139.
- 9 5. Aryal UK, et al. (2013) Proteome analyses of strains ATCC 51142 and PCC 7822 of the
10 diazotrophic cyanobacterium *Cyanothece* sp. under culture conditions resulting in
11 enhanced H₂ production. *Appl Environ Microbiol* 79(4):1070-1077.
- 12 6. Schneegurt MA, Sherman DM, Nayar S, Sherman LA (1994) Oscillating behavior of
13 carbohydrate granule formation and dinitrogen fixation in the cyanobacterium
14 *Cyanothece* sp. strain ATCC 51142. *J Bacteriol* 176(6):1586-1597.
- 15 7. Dong G, Golden SS (2008) How a cyanobacterium tells time. *Curr Opin Microbiol*
16 11(6):541-546.
- 17 8. Iwasaki H, Kondo T (2004) Circadian timing mechanism in the prokaryotic clock system
18 of cyanobacteria. *J Biol Rhythms* 19(5):436-444.
- 19 9. Johnson CH, Mori T, Xu Y (2008) A cyanobacterial circadian clockwork. *Curr Biol*
20 18(17):R816-R825.
- 21 10. Nakajima M, et al. (2005) Reconstitution of circadian oscillation of cyanobacterial KaiC
22 phosphorylation *in vitro*. *Science* 308(5720):414-415.

- 1 11. Johnson CH, Elliott J, Foster R, Honma K-I, Kronauer R (2004) in *Chronobiology:*
2 *Biological Timekeeping*, eds Dunlap JC, Loros JJ, DeCoursey PJ (Sinauer Associates,
3 Sunderland), pp 66-105.
- 4 12. Welsh EA, et al. (2008) The genome of *Cyanothece* 51142, a unicellular diazotrophic
5 cyanobacterium important in the marine nitrogen cycle. *Proc Natl Acad Sci*
6 105(39):15094-15099.
- 7 13. Toepel J, Welsh E, Summerfield TC, Pakrasi HB, Sherman LA (2008) Differential
8 transcriptional analysis of the cyanobacterium *Cyanothece* sp. strain ATCC 51142 during
9 light-dark and continuous-light growth. *J Bacteriol* 190(11):3904-3913.
- 10 14. Toepel J, McDermott JE, Summerfield TC, Sherman LA (2009) Transcriptional analysis
11 of the unicellular, diazotrophic cyanobacterium *Cyanothece* sp. ATCC 51142 grown
12 under short day/night cycles. *J Phycol* 45(3):610-620.
- 13 15. Červený J, Nedbal L (2009) Metabolic rhythms of the cyanobacterium *Cyanothece* sp.
14 ATCC 51142 correlate with modeled dynamics of circadian clock. *J Biol Rhythms*
15 24(4):295-303.
- 16 16. Sinetova MA, Červený J, Zavřel T, Nedbal L (2012) On the dynamics and constraints of
17 batch culture growth of the cyanobacterium *Cyanothece* sp. ATCC 51142. *J Biotechnol*
18 162(1):148-155.
- 19 17. Elvitigala T, Stockel J, Ghosh B, Pakrasi H (2009) Effect of continuous light on diurnal
20 rhythms in *Cyanothece* sp. ATCC 51142. *BMC Genomics* 10(1):226.
- 21 18. Tu BP, Kudlicki A, Rowicka M, McKnight SL (2005) Logic of the yeast metabolic cycle:
22 temporal compartmentalization of cellular processes. *Science* 310(5751):1152-1158.

- 1 19. Nedbal L, Trtílek M, Červený J, Komárek O, Pakrasi HB (2008) A photobioreactor
2 system for precision cultivation of photoautotrophic microorganisms and for high-content
3 analysis of suspension dynamics. *Biotechnol Bioeng* 100(5):902-910.
- 4 20. Suzuki E, et al. (2013) Physicochemical variation of cyanobacterial starch, the insoluble
5 α -Glucans in cyanobacteria. *Plant Cell Physiol* 54(4):465-473.
- 6 21. McDermott JE, et al. (2011) A model of cyclic transcriptomic behavior in the
7 cyanobacterium *Cyanothece* sp. ATCC 51142. *Mol BioSyst* 7(8):2407-2418.
- 8 22. Schneegurt MA, Tucker DL, Ondr JK, Sherman DM, Sherman LA (2000) Metabolic
9 rhythms of a diazotrophic cyanobacterium, *Cyanothece* sp. strain atcc 51142,
10 heterotrophically grown in continuous dark. *J Phycol* 36(1):107-117.
- 11 23. Elvitigala T, Pakrasi HB, Ghosh BK (2009) in *Emergent Problems in Nonlinear Systems*
12 *and Control*, eds Ghosh B, Martin CF, Yishao Z (Springer, Heildeberg), pp 21-41.
- 13 24. Chen C-I, McDonald KA (1990) Oscillatory behavior of *Saccharomyces cerevisiae* in
14 continuous culture: II. Analysis of cell synchronization and metabolism. *Biotechnol*
15 *Bioeng* 36(1):28-38.
- 16 25. Müller D, Exler S, Aguilera-Vázquez L, Guerrero-Martín E, Reuss M (2003) Cyclic AMP
17 mediates the cell cycle dynamics of energy metabolism in *Saccharomyces cerevisiae*.
18 *Yeast* 20(4):351-367.
- 19 26. Saito T, Mitsui K, Hamada Y, Tsurugi K (2002) Regulation of the Gts1p level by the
20 ubiquitination system to maintain metabolic oscillations in the continuous culture of
21 yeast. *J Biol Chem* 277(37):33624-33631.
- 22 27. Tu BP, McKnight SL (2006) Metabolic cycles as an underlying basis of biological
23 oscillations. *Nat Rev Mol Cell Biol* 7(9):696-701.

- 1 28. Reddy KJ, Haskell JB, Sherman DM, Sherman LA (1993) Unicellular, aerobic nitrogen-
2 fixing cyanobacteria of the genus *Cyanothece*. *J Bacteriol* 175(5):1284-1292.
- 3 29. Provasoli L, McLaughlin JJA, Droop MR (1957) The development of artificial media for
4 marine algae. *Arch Mikrobiol* 25(4):392-428.
- 5 30. van Baalen C (1962) Studies on marine blue-green algae. *Bot Mar* 4:129-139.
- 6 31. Červený J, Šetlík I, Trtílek M, Nedbal L (2009) Photobioreactor for cultivation and real-
7 time, *in-situ* measurement of O₂ and CO₂ exchange rates, growth dynamics, and of
8 chlorophyll fluorescence emission of photoautotrophic microorganisms. *Eng Life Sci*
9 9(3):247-253.
- 10 32. Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J (2007) qBase relative
11 quantification framework and software for management and automated analysis of real-
12 time quantitative PCR data. *Genome Biol* 8(2):R19.

13

1 **Figure legends**

2 **Fig. 1.** Transition from ultradian to circadian rhythms in diurnally entrained batch culture.
3 *Cyanothece* culture was diurnally entrained in a turbidostat during 28 days with 12 h light /12 h
4 dark, 36 °C, in the ASP2 medium without nitrate. The dark/light entrainment was followed by
5 constant light $136 \mu\text{mol}(\text{photons})\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ of 627 nm and $10 \mu\text{mol}(\text{photons})\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ of cool white
6 light with turbidostat switched off. The upper panel shows the experiment period starting with
7 the last 24 h entrainment period and ending after 10 days of continuous light. The black rectangle
8 between 12 and 24 h shows the last night dark period and the subsequent striped rectangles show
9 subjective (anticipated) night periods. The lower two panels show typical ultradian (72-96 h) and
10 circadian (216-264 h) rhythms. The solid line shows the culture growth dynamics as measured
11 by optical density at 680 nm, OD_{680} . The thick line with open circles shows relative changes in
12 the dissolved O_2 . The other measurements are only shown in the lower two panels: The thin full
13 line represents relative change in the dissolved CO_2 , the dashed line represents relative change of
14 pH and the long-dashed line shows relative change of the steady state pigment fluorescence
15 emission.

16

17 **Fig. 2.** Batch culture transition from ultradian to circadian rhythms after an exchange to nitrate-
18 less medium. The culture growing rapidly in nitrate supplemented ASP2 medium was harvested
19 and re-suspended in nitrate-less medium (time 0). The transition introduced oscillations in
20 dissolved O_2 , in OD_{680} optical density, steady-state fluorescence emission F_t , and in pH. The
21 dissolved CO_2 is represented here by the anti-parallel pH dynamics. The symbols are the same as
22 in Fig. 1.

23

1 **Fig. 3.** Transition from ultradian to circadian rhythms after decrease of light in a turbidostat. The
2 oscillations of dissolved O₂, CO₂, stationary fluorescence Ft, and pH were dominated by the
3 ultradian rhythm during the first 64 hours when exposed to the high irradiance
4 (160 μmol(photons).m⁻².s⁻¹). Switching to low irradiance (50 μmol(photons).m⁻².s⁻¹) lead to an
5 decrease in the ultradian rhythm amplitude and to prevalence of the circadian oscillation.
6 Noteworthy, the saw-like OD₆₈₀ curve shows that the culture was automatically diluted whenever
7 its density grew over 0.7. The dilutions were less frequent around minima in the concentration of
8 dissolved O₂ indicating slow growth when photosynthesis is suppressed.

9

10 **Fig. 4.** Temperature dependence of the ultradian (open circles) and circadian (closed circles)
11 cycle periods. The data were obtained as averages of at least two independent experiments. In 33
12 and 36°C, more independent experiments were performed to determine the standard deviation of
13 ±0.05 h. The temperature coefficients (Q₁₀) were calculated to quantify the temperature
14 dependence of a rate proxy 1/period.

15

16 **Fig. 5.** The processes occurring during the ultradian oscillations relative to the photosynthesis
17 and nitrogen fixation phases. (A) Oscillation of the dissolved O₂ (open circles), pH (closed
18 circles) and percentage of dividing cells as counted in a microscope (bars). (B) Total carbon in
19 cells (closed rectangles) increased during the photosynthesis phase whereas it remained stagnant
20 during the nitrogen fixation phase. Glycogen content (open rectangles) decreased during the
21 nitrogen fixing phase. (C) Dynamics of the total nitrogen content of photobioreactor biomass
22 together with expression levels of nitrogenase genes. The dashed line shows the increase of N
23 content during the nitrogen fixation phase and stagnant level during the oxygenic photosynthesis

- 1 phase. The solid lines show expression levels (NRQ) of the *nifB* (x marks) and *nifH* (+ marks)
- 2 genes together with measured nitrogenase activity (open triangles).