

2003

Interposed Abdominal Compression CPR: A Comprehensive Evidence Based Review

Charles F. Babbs

Purdue University, babbs@purdue.edu

Follow this and additional works at: <http://docs.lib.purdue.edu/bmepubs>



Part of the [Biomedical Engineering and Bioengineering Commons](#)

Recommended Citation

Babbs, Charles F, "Interposed Abdominal Compression CPR: A Comprehensive Evidence Based Review" (2003). *Weldon School of Biomedical Engineering Faculty Publications*. Paper 63.
<http://docs.lib.purdue.edu/bmepubs/63>

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Interposed Abdominal Compression CPR: A Comprehensive Evidence Based Review

Charles F. Babbs, MD, PhD

Weldon School of Biomedical Engineering, Purdue University,
and Indiana University School of Medicine

(Resuscitation, 59, 71-82, 2003)

Abstract

Interposed abdominal compression (IAC)-CPR includes all steps of standard external CPR with the addition of manual mid-abdominal compressions in counterpoint to the rhythm of chest compressions. IAC-CPR can increase blood flow during CPR about two fold compared to standard CPR without IAC, as shown by 6 of 6 studies in computer models and 19 of 20 studies in various animal models. The addition of IAC has clinical benefit in humans, as indicated in 10 of 12 small to medium sized clinical studies. The technique increases the frequency of immediate return of spontaneous circulation for in-hospital resuscitations from roughly 25% to 50%. Improved survival to discharge is also likely on the basis of two small in-hospital trials. Possible harm from abdominal compression is minimal on the basis of 426 humans, 151, dogs and 14 pigs that received IAC in published reports. The complexity of performing IAC is similar to that of opening the airway and is less than that of other basic life support maneuvers. The aggregate evidence suggests that IAC-CPR is a safe and effective means to increase organ perfusion and survival, when performed by professionally trained responders in a hospital and when initiated early in the resuscitation protocol. Cost and logistical considerations discourage use of IAC-CPR outside of hospitals.

Keywords: Abdomen, Clinical trials, Guidelines, Interposed abdominal compression-CPR, IAC-CPR, Review, Statistical analysis

1. Introduction

Interposed abdominal compression (IAC) – CPR includes all the steps of ordinary CPR with the addition of external mid-abdominal compressions by a second or third rescuer, timed between chest compressions. The abdominal compressions are delivered in counterpoint to the rhythm of chest compressions, so that abdominal pressure is maintained whenever the chest is not being compressed. Pulses of central abdominal pressure are applied with overlapping hands just headward of the umbilicus. Hand position, depth, rhythm, and rate of abdominal compression are similar to those for chest compression.

The force of abdominal compression is similar to that needed to palpate the abdominal aortic pulse. The technique of IAC-CPR achieved some notoriety in the 1990's after the clinical studies of Sack and coworkers, involving several hundred patients, suggested a rough doubling of immediate resuscitation success and neurologically intact, long term survival¹.

The present paper is an extensive evidence based review. Relevant full length, peer reviewed publications were identified using evidence evaluation methods described previously^{2,3} and developed specifically for topics in emergency cardiovascular care. Full-length, peer reviewed publications were obtained from MEDLINE searches, the author's files, and reference lists of review articles on newer techniques in resuscitation as referenced in "Guidelines 2000 for cardiopulmonary resuscitation and emergency cardiovascular care: international consensus on science"⁴.

The results of the search turned up a rather remarkable accumulation of positive evidence published since 1980, documenting the hemodynamic and clinical benefits of IAC-CPR, when compared to conventional standard CPR in a variety of pre-clinical and clinical models (Table 1). Despite the relatively small numbers of patients reported in human clinical trials, the overall tone of this review is therefore positive and rather different from that of the usual conservative review of controversial topics in science: "John found this, but Mary found the opposite. More research is needed." Of the 38 full-length, peer reviewed studies (Table 2) relevant to IAC-CPR, 34 obtained results indicating benefit of IAC. Two found no difference between IAC and standard CPR, and two found results worse for IAC than for standard CPR. Only a single case report describes an adverse complication attributable to external abdominal compression—possible traumatic pancreatitis in a child. Even the authors of this report stated that their finding was "not meant to condemn a technique which may ultimately prove superior to conventional CPR"⁵.

Table 1. Broad summary of research on IAC-CPR in 3 classes of experimental models.

Model Class	Subtypes	Results
Mathematical, computer, and mechanical models	Paper and pencil ⁶	6 of 6 studies positive (100 percent)
	Analog ^{7, 8}	
	Digital ^{9, 10}	
	Digital spreadsheet ¹¹	
Animal models (dogs and pigs)	Blood flow ¹²⁻²⁰	19 of 20 studies positive (95 percent)
	Blood pressure ²¹⁻²⁴	
	Oxygen uptake ²⁵	
	CO ₂ excretion/blood gasses ²⁶	
	Survival and complications ²⁷⁻³¹	
Clinical models	Pediatric anesthesia overdose ³²	10 of 12 studies positive (83 percent)
	Acute hemodynamics (BP, ETCO ₂ ± crossover) ³³⁻³⁹	
	All arrests (mostly VF) ^{1, 40}	
	Difficult arrests (EMD, asystole, prolonged) ^{41, 42}	

For most end points results for IAC-CPR were between 1.5 and 2.0 times those for comparable standard CPR without abdominal compression.

EMD=electromechanical dissociation, ETCO₂=end tidal carbon dioxide concentration, SPICE=simulation program for integrated circuit evaluation, VF=ventricular fibrillation.

Table 2. Data base composition by level of evidence *

LEVEL	CAPSULE DESCRIPTION OF LEVEL	# OF PAPERS
1	Larger randomized clinical trials ^{1, 40, 41}	3
2	Smaller randomized clinical trials ³⁴	1
3	Prospective, non-randomized cohort studies ^{36, 43}	2
4	Historic, non-randomized cohort studies ^{32, 37, 44}	3
5	Human case series ^{38, 39, 45}	3
6	Animal or mechanical model studies ⁶⁻³¹	25
7	Rational extrapolations or quasi-experimental designs ⁴⁶	1
8	Common sense, etc.	0

* Performance sites for reviewed studies included USA 28 studies, UK 2, Israel 2, Canada 2, Japan 1, Germany 1, Netherlands 1, Spain 1, and Italy 1. (Some studies had multiple sites.)

Table 3: Departmental affiliations of investigators publishing on IAC-CPR

Anesthesiology	27
Biomedical Engineering	10
Cardiology	3
Critical Care	2
Emergency Medicine	8
Industry	1
Internal Medicine	7
Neurobiology	1
Physiology	6
Radiology	1
Surgery	6

Why has IAC-CPR not been enthusiastically adopted already for widespread clinical use? Why is the method only recommended in current guidelines as an optional alternative technique rather than standard of care? Probably in part because nearly all of the 38 studies were small, and inexpensive affairs, often conducted by researchers fundamentally interested in other research topics or who came upon the phenomenon of IAC by accident. Moreover, as is typical in the field of cardiopulmonary resuscitation, there is a paucity of human data. Large multi-center clinical trials were never conducted. Moreover, in the United States changed federal regulations in the 1990s, making clinical studies of resuscitation exceedingly difficult with respect to informed consent, were promulgated just as clinical evidence favoring IAC started to accumulate^{47, 48}.

In this sense the technique of IAC-CPR is analogous to an orphan drug. No device or product is needed to perform IAC, and so there is no multi-national corporation to fund or support relevant research and development. Appreciation of the science supporting abdominal compression as an adjunct to chest compression requires the aggregation and synthesis of scattered studies published in diverse journals by investigators from 9 countries in 11 different sub-specialties of science and medicine (Tables 2 and 3). Such a comprehensive review has not been published heretofore.

2. Early work on abdominal counterpulsation

Three years before external CPR was described by Kouwenhoven, Jude, and Knickerbocker in 1960⁴⁹, Rainer and Bullough in the United Kingdom developed a jackknife maneuver for resuscitation of children from anesthetic overdose, which involved rhythmically compressing first the abdomen and then the chest³². The technique required the surgeon to fold the knees of a child as far as possible toward the chin, forcing the thighs against the abdomen. Then "when full flexion is reached, the surgeon compresses the patient's chest with the bent knees, using some of his own weight judiciously to add to the pressure".³² This method can be regarded as a version of phased abdominal and chest compression and was successful in 8 of 8 reported cases.

In 1972 Molokhia, Norman and coworkers at Boston City Hospital²¹ described a similar means of generating an artificial circulation during open chest surgery by alternately compressing the aortic arch and the left ventricle. The addition of interposed aortic compressions augmented coronary perfusion by about 50 percent. In 1976 Ohomoto²² at Tokyo Women's medical college resuscitated anesthetized dogs with two motor driven pistons, one compressing the abdomen, the other the chest. The addition of phased abdominal compression for 80 percent of cycle time to chest compression for 25 percent of cycle time increased carotid artery blood pressure and improved short-term survival in anesthetized dogs with ventricular fibrillation.

Subsequently Rosborough and coworkers in Houston Texas⁵⁰ while attempting to develop an animal model of cough CPR⁵¹ combined simultaneous high-pressure lung inflation with abdominal compression. They found to their surprise that abdominal compression and

ventilation alone could maintain carotid flow and aortic pressure during ventricular fibrillation in dogs. They suggested the technique as a new CPR modality. Contemporaneously, Coletti, Bregman, and coworkers in New York were studying a canine model of cardiogenic shock in the laboratory one day, when the mechanical intra-aortic balloon pump failed to work properly¹³. As a substitute they tried manual external abdominal compressions interposed between heartbeats, as judged from the electrocardiogram. Both cerebral and coronary perfusion, measured with electromagnetic flowmeters, increased.

Ralston et al.¹⁴ described the modern idea of alternating compression of the abdomen and the chest for the purpose of external cardiopulmonary resuscitation in the year 1982. The first graphic record demonstrating the phenomenon is reproduced in reference²⁰. Subsequently a burst of preclinical studies, followed by clinical studies, appeared in the 1980's and 1990's. These included a wide variety of model systems and experimental end points (Table 1).

The following summary of evidence is organized according to the "Levels of Evidence" paradigm for topics in resuscitation research, which is described in detail in reference³. Research reports are assigned to a particular level according to the study design and methodology. Level 1 studies are considered the most relevant to human medicine, providing the strongest evidence favoring a novel intervention. Level 2 studies are the next most relevant, etc., continuing to Level 8 reports, which are speculative opinion pieces without actual data. Table 2 gives an overview of reviewed studies in terms of their levels of evidence, together with brief definitions of the levels.

3. Mechanical, electrical, and computer models (Level 6)

A number of investigators with a biomedical engineering bent have investigated IAC-CPR by building mechanical, electrical, or computer models⁶⁻¹¹. A good example of this genre is provided by the electrical model of Ralston and Babbs⁸, who appreciated the analogy between the flow of current around an electrical circuit and the circulation of blood through systemic and pulmonary vessels. In their simulations electrical resistance mimicked physiological resistance of vascular beds. Resistive-capacitive networks modeled the heart and blood vessels. Voltages represented intravascular pressures; electric current represented blood flow, inductance modeled blood inertia, and diodes served as the cardiac and venous valves. The application of half-sinusoidal voltage pulses to the model represented the effects of compression of the chest and abdomen during CPR. In this model the addition of IAC to standard CPR produced flow augmentation according to the expression

$$\text{flow} = \alpha P_{\text{th}} + \beta P_{\text{abd}},$$

where P_{th} is peak intrathoracic pressure, P_{abd} is peak intraabdominal pressure, and α and β are constants ($\alpha > \beta$). For typical adult CPR intra-abdominal pressure can be made

somewhat greater than intrathoracic pressure ($P_{abd} > P_{th}$), so that chest and abdominal flow components become roughly equal. Thus the addition of IAC doubled flow in this electrical model of the adult human.

Such theoretical models also provided a detailed glimpse into hemodynamic mechanisms of abdominal counterpulsation. Figure 1 shows simulated pressure waveforms in a more recent model of a typical adult human circulation¹¹ during standard and IAC-CPR. Two mechanisms are evident upon close inspection. The first is the obvious hump in aortic blood pressure (squares) during the abdominal compression phase, which is greater than the corresponding rise in central venous pressure (circles). The second, and subtler, mechanism is the augmentation of the filling phase of the chest pump (i.e. pulmonary vasculature), represented by the positive difference between right heart and chest pump pressure ($P_{rh} - P_{pump}$) during the last one fifth of the compression cycle. Thus external abdominal compression acts like an intra-aortic balloon pump, to squeeze more blood from the aorta into systemic resistance vessels. It also acts to squeeze more venous blood into the chest pump, priming it before the next chest compression. In this way phased abdominal pressure has beneficial effects upon both the arterial and the venous sides of the circulation.

(Figure 1 on next page)




Figure 1. Simulated pressure waveforms during standard and IAC-CPR in a typical adult. Steady state pressures after 20 compression cycles are shown. Peak applied pressures: chest 60 mmHg, abdomen 110 mmHg. Compression rate 90/min. (a) Standard CPR. (b) IAC-CPR. Ao=thoracic aorta, rh=right heart, pump=thoracic pump (pulmonary arteries and veins), aa=abdominal aorta, ivc=inferior vena cava. CPP is coronary perfusion pressure in mmHg. Flow is total systemic blood flow in L/min. Data redrawn from reference¹¹

4. Animal studies (Level 6)

The prediction of such theoretical models that IAC could nearly double blood flow during CPR turned out to be quite congruent with results of animal studies, which are also considered as Level 6 evidence. Much of the research comparing interposed abdominal compression CPR with standard CPR has been done in animal models—typically anesthetized pigs and anesthetized dogs with electrically induced ventricular fibrillation. Generally such models, summarized in Table 1 have found indices of blood flow during CPR to increase by 50 to 100 percent with the addition of IAC to the resuscitation protocol. Examples include increased carotid artery flow^{22, 23, 30, 52}, regional blood flow^{16, 18, 19, 25}, systemic and coronary perfusion pressures^{14, 19, 25, 29}, cardiac output²⁵, and oxygen delivery²⁵.

Animal studies have also tracked resuscitation success and survival. Kern's initial study of groups of 10 animals showed no significant difference in immediate resuscitation success or 24 hour survival²⁷. However Lindner et al.²⁹, using groups of 14 ketamine anesthetized pigs, found return of spontaneous circulation in 0/14 animals with standard CPR vs. 14/14 with IAC-CPR. Tang³⁰, using the Lifestick device that permitted one rescuer to do alternating chest and abdominal compression, obtained similar positive results in pigs—10/10 vs. 0/10 survival at 48-hours for chest compression plus IAC vs. chest compression alone.

5. Preliminary clinical studies (Levels 3-5)

Preliminary human studies, short of randomized clinical trials, provided further evidence for hemodynamic effects of external abdominal compressions. Howard et al.³⁵ instrumented 14 patients in the emergency department for blood pressure measurements during alternate trials of standard versus IAC-CPR. They found that IAC and high force IAC increased diastolic and mean arterial pressures, but not necessarily arteriovenous pressure differences, and that the effects of IAC were greater in non-obese patients. Adams³⁶ studied hemodynamics of IAC during human CPR in patients that were unsalvageable by conventional ACLS. There were 13 responders with a mean change in coronary perfusion pressure of 9.5 mmHg and 7 non-responders with a mean change in coronary perfusion pressure of -2.6 mmHg. Berryman and Phillips³⁴ also studied IAC-CPR in six cardiac arrest patients after unsuccessful ACLS. Mean arterial pressure increased from 26 to 39 mmHg. Villa, Colombo, and coworkers⁴² reported one case of a successful 1 hour and 20 min long resuscitation of a patient with pulseless electrical activity using a combination of mechanical chest compression and manual IAC. The patient lived subsequently for 8 hours following cardiac arrest and resuscitation.

Table 4. Summary of randomized clinical trials of IAC vs. standard CPR

Lead Author	Year	n	Results
Mateer ⁵³	1985	291	40/145 (28%) resuscitated with pre-hospital protocol involving STD-CPR, paramedic IAC-CPR, then STD-CPR during transport. 45/146 (31%) resuscitated with STD-CPR throughout the pre-hospital phase. No evidence of abdominal injury.
Ward ³³	1989	33	6/16 (37%) resuscitated in the emergency department with initial 20 min of IAC-CPR vs. 3/17 (18%) with STD-CPR. End Tidal CO ₂ averaged 17.1 mmHg with IAC-CPR vs. 9.6 mmHg with STD-CPR, indicating improved blood flow. No evidence of abdominal injury.
Sack #1 ¹	1992	103	29/48 (60%) resuscitated from in-hospital cardiac arrest with IAC-CPR vs. 14/55 (25%) with STD-CPR. Survival to discharge was 12/48 (25%) for IAC-CPR vs. 4/55 (7%) for STD-CPR. No evidence of abdominal injury from IAC.
Sack #2 ⁴¹	1992	143	33/67 (49%) resuscitated from in-hospital asystole or EMD with IAC-CPR vs. 21/76 (28%) with STD-CPR. No long term survivors. No evidence of abdominal trauma from IAC.

IAC = interposed abdominal compression, STD=standard, EMD=electromechanical dissociation (pulseless electrical activity of the heart)

6. Randomized clinical trials (Levels 1-2)

6.1. Efficacy of IAC-CPR

Three randomized clinical trials of IAC-CPR for in-hospital cardiac arrest have shown statistically significant improvement of outcome measures^{1,41}. One randomized trial of pre-hospital IAC-CPR, combined when possible with standard CPR in the field, showed no difference in outcome or in complications⁵³. These clinical trials are summarized in Table 4. Pooled analysis of all available data for both pre-hospital and in-hospital resuscitations show improvement in the return of spontaneous circulation with IAC-CPR, compared to standard CPR (Table 5). When only the in-hospital studies are examined, the effect of IAC becomes much greater. Pooled data from the two studies that examined long term, neurologically intact survival following in-hospital resuscitations show a similar relative benefit of IAC-CPR compared with standard CPR. A formal meta-analysis of statistical significance is presented in section 6.3.

Table 5. Aggregate human ROSC and survival data for IAC vs. standard CPR

Outcome Measure	Studies	IAC-CPR	Standard CPR
Return of spontaneous circulation (ROSC) in or out-of-hospital	Mateer ⁵³	40/145 (28%)	45/146 (31%)
	Ward ³³	6/16 (38%)	3/17 (18%)
	Sack #1 ¹	29/48 (60%)	14/55 (25%)
	Sack#2 ⁴¹	33/67 (49%)	21/76 (28%)
	All 4 studies	108/276 (39%)	83/294 (28%)
Return of spontaneous circulation (ROSC) after in-hospital resuscitation	Ward ³³	6/16 (38%)	3/17 (18%)
	Sack #1 ¹	29/48 (60%)	14/55 (25%)
	Sack#2 ⁴¹	33/67 (49%)	21/76 (28%)
	All 3 studies	68/131 (52%)	38/148 (26%)
Survival to discharge, neurologically intact after in-hospital resuscitation	Ward ³³	1/16 (6%)	0/17 (0%)
	Sack #1 ¹	8/48 (17%)	3/55 (5%)
	Both studies	9/64 (14%)	3/72 (4%)

6.2. Complications of IAC-CPR

The safety of interposed abdominal compressions, reviewed previously⁵⁴, has been well documented in 426 humans, 151 dogs and 14 pigs. Only one isolated case report of traumatic pancreatitis in a child⁴⁵ describes local trauma from abdominal compression during CPR. These data compare favorably with the well-known and frequent incidence of rib fracture and pulmonary contusion from chest compression during CPR⁵⁵⁻⁵⁷. Increased emesis and aspiration from IAC have not been reported, and there is evidence that if

positive abdominal pressure is applied during ventilations from the beginning of an arrest, the rate of gastric inflation before endotracheal intubation is reduced³¹.

6.3. Meta-analysis

A more rigorous meta-analysis of IAC-CPR has recently been published⁵⁸ that is limited to human clinical trials in which the end point was either short term or long term survival. These include all available survival data from the three in-hospital trials and one and out-of-hospital trial^{1, 33, 40, 41}. That is, hemodynamic data were not considered. Meta-analysis refers to the quantitative synthesis of data from multiple clinical studies in order to minimize both Type I and Type II statistical errors. Inverse variance weighting of outcome data is typically used to account for the relative numbers of patients in the various studies. Results for such an analysis of IAC-CPR trials reported in references are shown in Figure 2. This figure shows cumulative meta-analysis plots that demonstrate historical trends with the publication of each successive clinical study. In each plot the top data point and its 95 percent confidence interval represent the historically first trial, the next a combination of the first two trials, the third a combination of the first three trials, etc. This format is similar to that introduced by Lau and coworkers⁵⁹ for cumulative meta-analysis in cardiovascular medicine.

The combined studies of IAC-CPR, including all available in-hospital and out-of-hospital data (Figure 2(A)), showed a significant treatment effect for short-term and probably for long-term survival. The difference in the proportion of survivors, $\overline{\Delta p}$, is 10.7 percent for return of spontaneous circulation and 8.7 percent for hospital discharge. The effect of IAC in the overall meta-analysis is highly significant for return of spontaneous circulation ($P = 0.006$) and of borderline statistical significance ($P = 0.06$) for discharge survival. These summary values can be immediately converted into the number-needed-to-treat, $NNT = 1/\overline{\Delta p}$, the number of patients that must be treated to obtain one additional survivor⁶⁰. For return of spontaneous circulation the NNT is 9. For discharge survival the corresponding NNT is 12. When only in-hospital studies are considered (Figure 2(B)) the statistical significance of the meta-analysis for return of spontaneous circulation is even greater ($P < 0.0001$, $NNT = 4$).




Figure 2. A cumulative meta-analysis for IAC-CPR in terms of the mean difference in proportion of survivors⁵⁸. Successive points from top to bottom represent a separate meta-analysis after the appearance of each study^{1, 33, 41, 53} in the series. Solid symbols refer to data for return of spontaneous circulation (ROSC), open symbols to data for neurologically intact discharge survival. Horizontal lines indicate 95 percent confidence intervals. (A) All studies in which survival was measured, including one out-of-hospital and three in-hospital studies. (B) In-hospital studies only^{1, 33, 41}.

7. Implementation issues

Practical implementation of a new method requires consideration of issues other than simply efficacy and safety. If a technique is extremely costly, difficult to teach or to learn, or likely only to benefit a small number of patients (with a large number-needed-to-treat) then it may not be suitable for widespread clinical implementation. Since IAC requires no new equipment or drug, its added cost is minimal, as long as adequate personnel are available in the hospital setting. However, as initially discovered by Mateer and coworkers⁴⁰, lack of trained personnel in out-of-hospital settings can obviate the use of IAC-CPR. Short reviews of the issues of complexity, cost vs. benefit, and pedagogy are as follows.

7.1. Complexity

Making resuscitation efforts more complicated, especially for lay rescuers, is a tricky business. More complicated techniques may be less effective in practice, if the added complexity is difficult to learn and easily forgotten^{61,62}. The limitation of IAC-CPR to trained health care providers is one way to minimize problems in this area. Fortunately, the technique of interposed abdominal compression itself is much less complicated than most steps of basic life support for either lay or professional rescuers (notably the recovery position, the Heimlich maneuver, and AED use). A detailed analysis of the complexity of the maneuver compared to other steps of basic life support is provided in Tables 6 and 7. Abdominal compression is clearly no more difficult than chest compression and arguably easier, since rescuers already know chest compression, and there is no varying chest wall stiffness to overcome in the abdomen.

(Table 6 on next page)

Table 6. Summary of complexity analysis of various basic life support maneuvers done by lay or professional rescuers

Number of Steps	9		Placing Victim in Recovery Position			
	8					
	7				Dealing With Choking	
	6	Starting Chest Compressions				
	5					Using an AED
	4			Starting Rescue Breathing		
	3					
	2		Opening airway OR Doing IAC			
	1					
		1	2	3	4	5
	Number of if—then branches					

Table 7: Details of complexity analysis of various basic life support maneuvers done by lay or professional rescuers

Action	Basic Steps	Variations
Open airway	<ol style="list-style-type: none"> 1. Tilt head 2. Lift chin or jaw thrust 	<ol style="list-style-type: none"> 1. Standard head tilt 2. Injured head or neck
Call 911	<ol style="list-style-type: none"> 1. Call 911 2. Get AED 	<ol style="list-style-type: none"> 1. Alone 2. Have help
Start breathing	<ol style="list-style-type: none"> 1. Check breathing (look, listen, feel) 2. Mouth seal 3. Two slow breaths 4. Watch chest rise 	<ol style="list-style-type: none"> 1. Mouth-to-mouth 2. Face shields 3. Face masks
Start circulation	<ol style="list-style-type: none"> 1. Maintain head tilt 2. Find signs of circulation 3. Check for signs of circulation 4. Find compression position 5. Position hands 6. Begin compressions 	<ol style="list-style-type: none"> 1. Lay or professional
Place victim in recovery position	<ol style="list-style-type: none"> 1. Kneel 2. Arm position waving goodbye 3. Other arm position 4. Grasp hand 5. Pull and roll 6. Adjust hand and knee 7. Tilt head 8. Check breathing 9. Verify 911 call 	<ol style="list-style-type: none"> 1. Victim keeps breathing 2. Victim stops breathing
Deal with choking	<ol style="list-style-type: none"> 1. Position victim 2. Check ability to speak and/or airway 3. Get in position for Heimlich 4. Do Heimlich 5. Check results 6. Do finger sweeps 7. Repeat as needed 	<ol style="list-style-type: none"> 1. Conscious, standard 2. Unconscious, standard 3. Conscious, obese or pregnant 4. Unconscious, obese or pregnant
Use AED	<ol style="list-style-type: none"> 1. Turn on machine 2. Apply electrodes 3. Analyze 4. Shock sequence 5. No shock indicated sequence 	<ol style="list-style-type: none"> 1. Water present 2. Pacemaker 3. Transdermal patch 4. Hairy chest 5. Children
Perform IAC	<ol style="list-style-type: none"> 1. Locate compression site 2. Compress on chest release 	<ol style="list-style-type: none"> 1. Standard 2. Pregnancy or distended abdomen

7.2. Cost versus benefit

Suppose that IAC-CPR were implemented for in-hospital resuscitations only. How many more long term survivors ($\Delta N_{\text{survivors}}$) of cardiac arrest might be expected? One simple calculation can be done as follows:

$$\Delta N_{\text{survivors}} = N_{\text{arrests}} \cdot P_{\text{in-hosp}} \cdot \Delta P_{\text{rosc}} \cdot P_{\text{long term}},$$

where,

N_{arrests} is the number of cardiac arrests in nation per year

$P_{\text{in-hosp}}$ is the proportion of in-hospital resuscitations

ΔP_{rosc} is the change in proportion of ROSC attributable to IAC

$P_{\text{long term}}$ is the proportion of initial saves who walk out of the hospital.

The research reviewed herein indicates that ΔP_{rosc} is 22% for in-hospital studies.

Suppose, for example, that $N_{\text{arrests}} = 250,000$, $P_{\text{in-hosp}} = 0.2$, $\Delta P_{\text{rosc}} = 0.2$, and $P_{\text{long term}} = 0.3$, then

$$\Delta N_{\text{survivors}} = 3000 \text{ extra survivors}$$

Although IAC-CPR itself is a simple manual technique, the costs of implementing it widely for professional rescuers in hospitals would be significant. Every professional rescuer course, adult and pediatric, available from the American Heart Association and other organizations internationally would require revision of instructor manuals, student manuals, videotapes, and slides. If such revisions were done as part of the normal cycle of guideline revisions, every 5 to 10 years, then the added cost of including IAC-CPR with other revisions may well be reasonable. Similarly, there would be the cost of retraining CPR instructors. If upgrades in manikins for training were required (see the following section 7.3) to include abdominal pressure monitors, then these costs would need to be included as well.

If use of IAC-CPR were expanded from in-hospital resuscitations by professional rescuers to out-of-hospital resuscitations by professional rescuers, the logistical limitation of adequate personnel comes into play. Most EMS systems now have only two professional rescuers per vehicle. This limitation would preclude IAC-CPR in those systems, either limiting benefit, or increasing cost dramatically if a third paid rescuer were included on ambulances. This limitation was first noted by Mateer et al.⁴⁰, who could not do IAC-CPR during transport of patients in their initial out-of-hospital study. For this reason it would seem prudent to limit initial clinical implementation of IAC-CPR to use by professional rescuers in hospitals.

7.3. Teaching IAC-CPR

A few issues arise regarding teaching IAC-CPR, since any technique, inadequately learned and performed, will not be effective. The following is a synopsis of important points, illustrating that teaching and learning the technique of IAC should not be difficult.

The recommended technique of abdominal compression is adapted from that described by Sack and coworkers¹, who had the most favorable clinical results. Use a two-handed, straight-armed technique similar to that for chest compressions. The abdominal compression point is in the midline, one-half hand width (5 cm) headward of the umbilicus. To ensure proper hand placement on the abdomen, the umbilicus should be visible below the rescuer's hands as a midline landmark. Apply pressure with heel of the hand straight down at the abdominal compression point, keeping fingers off the abdomen.

The amount of force applied should be sufficient to generate approximately normal mean arterial pressure (95 mmHg) within the abdomen near the abdominal aorta. This value is similar to that reported to give good results in animal studies and in theoretical models. It is also less than that required to produce objectionable discomfort in an awake, conscious volunteer⁶³. As has been previously described, this is the same amount of force required for palpation of the abdominal aortic pulse during an ordinary physical examination of the abdomen and is therefore not excessive⁶³.

It is quite easy to modify a standard whole body mannequin for the teaching and practice of three-rescuer IAC-CPR. One can add extra foam rubber to the lower thoracic compartment of the mannequin and also to the abdominal compartment of the mannequin to simulate soft tissues. The next step is to wrap a standard arm blood pressure cuff around a rolled towel and place it in the abdominal compartment beneath the abdominal compression point (5-cm headward of the umbilicus in the midline). Then bring the tubing, aneroid manometer gauge, and squeeze bulb out at belt-line so they are visible to the trainees. Tape target pressure markers on the manometer dial at 20 and 120 mmHg. Inflate the cuff to a resting pressure of 20 mmHg. During practice coach trainees to hit the target pressure of 120 during IAC and release to a target pressure of 20 during chest compression. Failure to release abdominal pressure can lead to liver injury during the subsequent chest compression.

One can initiate trainees to the rhythm and mechanics of manual abdominal compressions in the following sequence: Let two experienced rescuers perform two-rescuer CPR on the mannequin as usual. The third (abdominal) rescuer, who is the trainee, takes position on opposite side of victim from chest compressor at the level of the abdomen. The chest compressor says, "You press here whenever I release," pointing to the abdominal compression point. The chest compressor counts "one – AND – two – AND – three – AND ...". The abdominal rescuer applies pressure during "AND". Abdominal compression is also maintained during mouth-to-mouth or bag-valve-mask rescue breaths to prevent gastric inflation with air³¹. It is helpful to start with slow-motion practice and then gradually to increase toward a normal compression rate. After the rhythm is mastered

the trainee can focus on hitting the target pressure on the manometer. These details suggest that IAC-CPR will be easy to teach and easy to learn.

7.4. Protocols for implementation

In keeping with current guidelines for ACLS⁴ it is best not to use IAC-CPR as a last ditch effort on individuals who have already failed conventional resuscitation efforts. Such a strategy ignores the results of the best clinical research studies on the subject. Outcome can be improved if IAC-CPR is begun as first line therapy in a hospital¹. However, IAC is known to be relatively ineffective in the failed ACLS model⁶⁴. Another known failure mode is the delayed institution of IAC-CPR after the unknown and varying down times of out-of-hospital cardiac arrest and failed initial resuscitation by standard CPR⁵³. The published evidence does not support the use of IAC-CPR as second line therapy. At best the technique simply doubles blood flow during CPR. Three rescuer IAC-CPR is not magic, but rather a modest evolution and extension of conventional two-rescuer CPR, which can improve hemodynamics and outcome when performed by trained professionals in a hospital setting.

8. Summary

IAC-CPR is an orphan innovation for a major public health menace in the developed world—sudden cardiac death. Having been independently discovered in Japan, Israel, the United Kingdom, and the United States by diverse researchers primarily interested in other topics, it seems to have been poorly championed and poorly advertised in the scientific community. Mentioned as an alternative adjunct in American Heart Association guidelines since 1992, the method is still rarely utilized. A review of published evidence indicates that IAC-CPR has a sound physiologic basis, does produce improved hemodynamics, and at the very least improves short-term resuscitation success (return of spontaneous circulation) in human beings, when initiated early in the resuscitation protocol.

Analysis of all available randomized clinical trials, including pre-hospital and in-hospital resuscitations, shows IAC-CPR by trained health professionals improves the probability of return of spontaneous circulation compared with standard CPR (Figure 2(A), $P < 0.01$). When analysis is limited to in-hospital studies only, the improvement in return of spontaneous circulation is highly statistically significant (Figure 2(B), $P < 0.0001$). Hemodynamic studies in a variety of animal and mechanical models generally show 50 to 100 percent improvement in artificial circulation and coronary perfusion pressure with the addition of IAC. Complications of IAC have been negligible in over 400 patients studied. There remains some uncertainty about the practicality and efficacy of out-of-hospital use of IAC, pending further research. Given the issues of increased cost, decreased practicality, and reduced effectiveness in out-of-hospital or pre-hospital settings, it is reasonable at the present time to recommend more wide spread clinical use of IAC-CPR by trained professional rescuers in a hospital setting. Since the most favorable clinical results

have been obtained when IAC-CPR is applied from the beginning of resuscitation, early application of the technique is to be encouraged. Use of IAC-CPR as a last-ditch effort after prolonged, failed ACLS is not recommended and is only marginally effective.

References

1. Sack JB, Kesselbrenner MB, Bregman D. Survival from in-hospital cardiac arrest with interposed abdominal counterpulsation during cardiopulmonary resuscitation. *JAMA*. 1992;267:379-385
2. Cummins RO, Chamberlain D, Hazinski MF, Nadkarni V, Kloeck W, Kramer E, Becker L, Robertson C, Koster R, Zaritsky A, Ornato JP, Callanan V, Allen M, Steen P, Connolly B, Sanders A, Idris A, Cobbe S. Recommended guidelines for reviewing, reporting, and conducting research on in-hospital resuscitation: The in-hospital "utstein style". *Circulation*. 1997;95:2213-2239
3. Cummins RO. Introduction to the international guidelines 2000 for cpr and ecc. *Circulation*. 2000;102 (suppl I):I-1--I-11
4. Cummins RO. American heart association in collaboration with the international liaison committee on resuscitation. Guidelines 2000 for cardiopulmonary resuscitation and emergency cardiovascular care: International consensus on science. *Circulation*. 2000;102 (suppl I):I-1--I-384
5. Waldman PJ, Walters BL, Grunau CF. Pancreatic injury associated with interposed abdominal compressions in pediatric cardiopulmonary resuscitation. *Am J Emerg Med*. 1984;2:510-512
6. Babbs CF. Abdominal counterpulsation in cardiopulmonary resuscitation: Animal models and theoretical considerations. *American Journal of Emergency Medicine*. 1985;3:165-170
7. Babbs CF, Weaver JC, Ralston SH, Geddes LA. Cardiac, thoracic, and abdominal pump mechanisms in cpr: Studies in an electrical model of the circulation. *American Journal of Emergency Medicine*. 1984;2:299-308
8. Babbs CF, Ralston SH, Geddes LA. Theoretical advantages of abdominal counterpulsation in cpr as demonstrated in a simple electrical model of the circulation. *Annals of Emergency Medicine*. 1984;13:660-671
9. Beyar R, Kishon Y, Sideman S, Dinnar U. Computer studies of systemic and regional blood flow during cardiopulmonary resuscitation. *Medical & Biological Engineering and Computing*. 1984;22:499-506
10. Babbs CF, Thelander K. Theoretically optimal duty cycles for chest and abdominal compression during external cardiopulmonary resuscitation. *Acad Emerg Med*. 1995;2:698-707
11. Babbs CF. Cpr techniques that combine chest and abdominal compression and decompression: Hemodynamic insights from a spreadsheet model. *Circulation*. 1999;100:2146-2152
12. Watson JT, Platt MR, Rogers DE, Sugg WL, Willerson JT. Similarities in coronary flow between external counterpulsation and intra-aortic balloon pumping. *American Journal of Physiology*. 1976;230:1616-1621

13. Coletti RH, Kaskel PS, Cohen SR, Bregman D. Abdominal counterpulsation (ac) -- a new concept in circulatory assistance. *Trans Am Soc Artif Intern Organs*. 1982;28:563-566
14. Ralston SH, Babbs CF, Niebauer MJ. Cardiopulmonary resuscitation with interposed abdominal compression in dogs. *Anesthesia and Analgesia*. 1982;61:645-651
15. Voorhees WD, Ralston SH, Babbs CF. Regional blood flow during cardiopulmonary resuscitation with abdominal counterpulsation in dogs. *Am J Emerg Med*. 1984;2:123-128
16. Walker JW, Bruestle JC, White BC, Evans AT, Indreri R, Bialek H. Perfusion of the cerebral cortex using abdominal counterpulsation during cpr. *Am J Emerg Med*. 1984;2:391-393
17. Einagle V, Bertrand F, Wise RA, Roussos C, Madger S. Interposed abdominal compressions and carotid blood flow during cardiopulmonary resuscitation: Support for a thoracoabdominal unit. *Chest*. 1988;93:1206-1212
18. Hoekstra OS, van Lambalgen AA, Groeneveld AB, van den Bos GC, Thijs LG. Abdominal compressions increase vital organ perfusion during cpr in dogs: Relation with efficacy of thoracic compressions. *Ann Emerg Med*. 1995;25:375-385
19. Wenzel V, Lindner KH, Prengel AW, Strohmenger HU. Effect of phased chest and abdominal compression-decompression cardiopulmonary resuscitation on myocardial and cerebral blood flow in pigs. *Crit Care Med*. 2000;28:1107-1112
20. Babbs CF, Blevins WE. Abdominal binding and counterpulsation in cardiopulmonary resuscitation. *Critical Care Clinics*. 1986;2:319-332
21. Molokhia FA, Ponn RB, Robinson WJ, Asimacopoulos PJ, Norman JC. A method of augmenting coronary perfusion during internal cardiac massage. *Chest*. 1972;62:610-613
22. Ohomoto T, Miura I, Konno S. A new method of external cardiac massage to improve diastolic augmentation and prolong survival time. *Ann Thorac Surg*. 1976;21:284-290
23. Beyar R, Kishon Y, Kimmel E, Neufeld H, Dinnar U. Intrathoracic and abdominal pressure variations as an efficient method for cardiopulmonary resuscitation: Studies in dogs compared with computer model results. *Cardiovasc Res*. 1985;19:335-342
24. Christenson JM, Hamilton DR, Scott-Douglas NW, Tyberg JV, Powell DG. Abdominal compressions during cpr: Hemodynamic effects of altering timing and force. *J Emerg Med*. 1992;10:257-266
25. Voorhees WD, Niebauer MJ, Babbs CF. Improved oxygen delivery during cardiopulmonary resuscitation with interposed abdominal compressions. *Annals of Emergency Medicine*. 1983;12:128-135
26. Lippert AC, Evans AT, White BC, Eyster GE. The effect of resuscitation technique and pre-arrest state of oxygenation on blood-gas values during cardiopulmonary resuscitation in dogs. *Vet Surg*. 1988;17:283-290
27. Kern KB, Carter AB, Showen RL, Voorhees WD, Babbs CF, Tacker WA, Ewy GA. Twenty-four hour survival in a canine model of cardiac arrest comparing three methods of manual cardiopulmonary resuscitation. *J Am Coll Cardiol*. 1986;7:859-867

28. Kern KB, Carter AB, Showen RL, Voorhees WdD, Babbs CF, Tacker WA, Ewy GA. Cpr-induced trauma: Comparison of three manual methods in an experimental model. *Ann Emerg Med.* 1986;15:674-679
29. Lindner KH, Ahnefeld FW, Bowdler IM. Cardiopulmonary resuscitation with interposed abdominal compression after asphyxial or fibrillatory cardiac arrest in pigs. *Anesthesiology.* 1990;72:675-681
30. Tang W, Weil MH, Schock RB, Sato Y, Lucas J, Sun S, Bisera J. Phased chest and abdominal compression-decompression. A new option for cardiopulmonary resuscitation. *Circulation.* 1997;95:1335-1340
31. Babbs CF, Schoenlein WE, Lowe MW. Gastric insufflation during iac-cpr and standard cpr in a canine model. *American Journal of Emergency Medicine.* 1985;3:99-103
32. Rainer EH, Bullough J. Respiratory and cardiac arrest during anaesthesia in children. *British Medical Journal.* 1957:1024-1028
33. Ward KR, Sullivan RJ, Zelenak RR, Summer WR. A comparison of interposed abdominal compression cpr and standard cpr by monitoring end-tidal pco2. *Ann Emerg Med.* 1989;18:831-837
34. Berryman CR, Phillips GM. Interposed abdominal compression-cpr in human subjects. *Annals of Emergency Medicine.* 1984;13:226-229
35. Howard M, Carrubba C, Foss F, Janiak B, Hogan B, Guinness M. Interposed abdominal compression-cpr: Its effects on parameters of coronary perfusion in human subjects. *Ann Emerg Med.* 1987;16:253-259
36. Adams CP, Martin GB, Rivers EP, Ward KR, Smithline HA, Rady MY. Hemodynamics of interposed abdominal compression during human cardiopulmonary resuscitation. *Acad Emerg Med.* 1994;1:498-502
37. Barranco F, Lesmes A, Irls JA, Blasco J, Leal J, Rodriguez J, Leon C. Cardiopulmonary resuscitation with simultaneous chest and abdominal compression: Comparative study in humans. *Resuscitation.* 1990;20:67-77
38. Soroff HS, Ruiz U, Birtwell WC, Many M, Giron F., Deterling RA. Assisted circulation by external pressure variation. *Physiology and Metabolism.* 1969;5:506-513
39. McDonald JL. Effect of interposed abdominal compression during cpr on central arterial and venous pressures. *Am J Emerg Med.* 1985;3:156-159
40. Mateer JR, Stueven HA, Thompson BM, Aprahamian C, Darin JC. Pre-hospital iac-cpr versus standard cpr: Paramedic resuscitation of cardiac arrests. *Am J Emerg Med.* 1985;3:143-146
41. Sack JB, Kesselbrenner MB, Jarrad A. Interposed abdominal compression-cardiopulmonary resuscitation and resuscitation outcome during asystole and electromechanical dissociation. *Circulation.* 1992;86:1692-1700
42. Villa GF, Colombo S, Cabrini L, Scandroglio AM, Torri G. [interposed abdominal compression in mechanical cardiopulmonary resuscitation. Description of a clinical case]. *Minerva Anesthesiol.* 1998;64:415-418
43. Howard M, Carrubba C, Foss F, Janiak B, Hogan B, Guinness M. Interposed abdominal compression-cpr: Its effects on parameters of coronary perfusion in human subjects. *Annals of Emergency Medicine.* 1987;16:253-259
44. Hillman H. Abdominal pumping. *Acad Emerg Med.* 1994;1:478-481

45. Waldman PJ, Walters BL, Grunau C. Pancreatic injury associated with interposed abdominal compressions in pediatric cardiopulmonary resuscitation. *Am J Emerg Med.* 1984;2:510-512
46. Babbs CF. Interposed abdominal compression-cardiopulmonary resuscitation: Are we missing the mark in clinical trials? [editorial]. *American heart journal.* 1993;126:1035-1041
47. Biros MH, Fish SS, Lewis RJ. Implementing the food and drug administration's final rule for waiver of informed consent in certain emergency research circumstances. *Acad Emerg Med.* 1999;6:1272-1282
48. Biros MH, Runge JW, Lewis RJ, Doherty C. Emergency medicine and the development of the food and drug administration's final rule on informed consent and waiver of informed consent in emergency research circumstances. *Acad Emerg Med.* 1998;5:359-368
49. Kouwenhoven WB, Jude JR, Knickerbocker GG. Closed-chest cardiac massage. *JAMA.* 1960;173:1064-1067
50. Rosborough JP, Niemann JT, Criley JM, O'Bannon W, Rouse D. Lower abdominal compression with synchronized ventilation--a cpr modality. *Circulation.* 1981;64(IV):303
51. Criley JM, Blaufuss AH, Kissel GL. Cough-induced cardiac compression--a self-administered form of cardiopulmonary resuscitation. *JAMA.* 1976;236(11):1246-1250
52. Einagle V, Bertrand F, Wise RA, Roussos C, Magder S. Interposed abdominal compressions and carotid blood flow during cardiopulmonary resuscitation. Support for a thoracoabdominal unit. *Chest.* 1988;93:1206-1212
53. Mateer JR, Steuven HA, Thompson BM, Aprahamian C, Darin J. Pre-hospital iac-cpr versus standard cpr: Paramedic resuscitation of cardiac arrests. *American Journal of Emergency Medicine.* 1985;3:143-146
54. Sack JB, Kesselbrenner MB. Hemodynamics, survival benefits, and complications of interposed abdominal compression during cardiopulmonary resuscitation. *Acad Emerg Med.* 1994;1:490-497
55. Clark DT. Complications following closed-chest cardiac massage. *JAMA* 181(4). 1962:337-338
56. Nagel EL, Fine EG, Krischer JP, Davis JH. Complications of cpr. *Critical Care Medicine* 9. 1981:424
57. Silberberg B, Rachmaninoff N. Complications following external cardiac massage. *Surg Gynecol Obstet* 119. 1964:6-10
58. Babbs CF. Simplified meta-analysis of clinical trials in resuscitation. *Resuscitation.* 2003;In Press
59. Lau J, Antman EM, Jimenez-Silva J, Kupelnick B, Mosteller F, Chalmers TC. Cumulative meta-analysis of therapeutic trials for myocardial infarction. *N Engl J Med.* 1992;327:248-254
60. McQuay HJ, Moore RA. Using numerical results from systematic reviews in clinical practice. *Ann Intern Med.* 1997;126:712-720
61. Assar D, Chamberlain D, Colquhoun M, Donnelly P, Handley AJ, Leaves S, Kern KB. Randomised controlled trials of staged teaching for basic life support. 1. Skill acquisition at bronze stage. *Resuscitation.* 2000;45:7-15.

62. Chamberlain D, Smith A, Colquhoun M, Handley AJ, Kern KB, Wollard M. Randomized controlled trials of staged teaching for basic life support. 2. Comparison of cpr performance and skill retention using either staged instruction or conventional teaching. *Resuscitation*. 2001;50:27-37
63. Babbs CF. Iac-cpr: Are we missing the mark in clinical trials? *American heart journal*. 1993;126:1035-1041
64. Ward KR. Possible reasons for the variability of human responses to iac-cpr. *Acad Emerg Med*. 1994;1:482-489