Purdue University

Purdue e-Pubs

Historical Documents of the Purdue Cooperative Extension Service

Department of Agricultural Communication

2-1-1958

Amounts of Plant-Food Used by Different Crops and Proportion of Nitrogen Returned to the Land in Various Legumes

Purdue University Cooperative Extension Service

Follow this and additional works at: https://docs.lib.purdue.edu/agext Higher Crop Yields From Improved Soils

Purdue University Cooperative Extension Service, "Amounts of Plant-Food Used by Different Crops and Proportion of Nitrogen Returned to the Land in Various Legumes" (1958). *Historical Documents of the Purdue Cooperative Extension Service*. Paper 336.

https://docs.lib.purdue.edu/agext/336

For current publications, please contact the Education Store: https://mdc.itap.purdue.edu/
This document is provided for historical reference purposes only and should not be considered to be a practical
reference or to contain information reflective of current understanding. For additional information, please contact the
Department of Agricultural Communication at Purdue University, College of Agriculture: http://www.ag.purdue.edu/
agcomm

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Purdue University Agricultural Extension Service Mimeo AY-25a Revised Feb., 1958

AMOUNTS OF PLANT-FOOD USED BY DIFFERENT CROPS AND PROPORTION OF NITROGEN RETURNED TO THE LAND IN VARIOUS LEGUMES

		Plant-food	removed, (pounds per a	acre)*
	Yield		Phosphoric	
Kind of Crop	per acre	Nitrogen_	acid	Potash
Corn, ears	80 bus.	76	32	19
Corn stover	2.5 tons	45	10	58
<u>Total</u>		121	142	77
Wheat, grain	30 bus.	36	16	10
Wheat straw	1.2 ton	12	4	<u>18</u>
<u>Total</u>		48	20	28
Oats, grain	50 bus.	31	13	
Oats straw	1 ton	12	4	30
<u>Total</u>		43	17	39
Rye, grain	25 bus.	29	10	7
Rye straw	1.5 tons	1/4	<u> </u>	<u>27</u>
Total		43	15	34
Barley, grain	30 bus.	27	12	12
Barley straw	.8 tons			19
Total	20 1	36	15	31
Soybean grain	30 bus.	109	24	32
Soybean straw	1.2 ton	14	8	20
Total Alfalfa hay	1 + 000	123 200	32	<u>52</u> 128
Clover hay	4 tons 2 tons	82	33 16	65
Lespedeza hay	1.5 tons	61	9	34
neshedesa nah	To Comp	OT	7) 4
Soybean hay	2 tons	94	20	40
Timothy hay	1.5 tons	30 30	9	45
Sudan grass	3 tons	85	16	78
Second Stand	7 0011B			10
Brome grass (W/a)	f.) 2 tons	82	17	110
Orchard grass	2 tons	40	19	100
		et =.	-,	

^{*}Compiled from Morrison's "Feeds and Feeding", 21st Edition, and Ill. A.E.S. Bulletin 518.

In general, legumes, if well inoculated, may secure up to two-thirds of their nitrogen from the air. They differ in the proportion removed in the tops and left in the roots and stubble.

Deep rooted legumes like the clovers, alfalfa, and sweet clover leave more nitrogen in the roots and stubble than shallow rooted crops like soybeans and possibly lespedeza.

A thick stand of spring seeded sweet clover plowed under in April the following spring will add from 75 pounds of nitrogen per acre from a fair stand and growth to 200 pounds from a thick heavy growth.

Roots and stubble of a full stand of clover contain about 40 pounds of nitrogen per acre, of alfalfa 50 pounds, of lespedeza 30 pounds and of soybeans 20 pounds.

When all top growth of inoculated soybeans is removed for hay or seed, more nitrogen is taken from the soil than the amount left in stubble and roots. When combined and the straw left on the ground, slightly more nitrogen is left than is removed in the grain.

About 2/3 the nitrogen, 3/4 the phosphoric acid, and 3/5 the potash in the ration feed to livestock can be returned to the land in manure if properly handled and the liquid saved by absorption in bedding.

(3M)

Cooperative Extension Work in Agriculture and Home Economics
State of Indiana, Purdue University
and the United States Department of Agriculture Cooperating
L. E. Hoffman, Director, Lafayette, Indiana
Issued in furtherance of the Acts of May 8 and June 30, 1914.