FINAL REPORT

Wetland Replacement Practices and Procedures for Indiana Highway Projects

FHWA/IN/JTRP-2002/18

by

James E. Alleman, *Professor* [Purdue University]

Steve M. Graves *MS* [Purdue University]

Project No: C-36-36EE

File No: 6-14-31

SPR-2205

***Joint Transportation Research Program
Purdue University***

Prepared as Part of an Investigation Conducted by the
Indiana Department of Transportation
and the
Federal Highway Administration
U.S. Department of Transportation

The contents of this report reflect the views of the authors who are responsible for the facts and accuracy of the data presented herein. The contents do not necessarily reflect the official views of, or policies of any of the sponsors. Furthermore, this report does not constitute a standard, specification, or regulation.

***School of Civil Engineering
Purdue University
West Lafayette, Indiana 47907***

June 2002

INDOT Research

TECHNICAL *Summary*

Technology Transfer and Project Implementation Information

TRB Subject Code: 17-2 Ecological Systems
Publication No.: FHWA/IN/JTRP-2002/18, SPR-2205

June 2002
Final Report

Wetland Replacement Practices and Procedures for Indiana Highway Projects

Introduction

This project originated from the need to determine what has led to the loss of wetlands within the state of Indiana. In particular, INDOT has been blamed for contributing to wetlands destruction within the state in relation to their historical involvement with Highway construction. This accusation assumes, rightly or wrongly, that the construction of state and federal roads carried out by INDOT has adversely affected wetlands on both a quantitative and qualitative basis. Whether the accusation of a negative impact quantitatively is properly placed is the purpose of this research.

Aspects of wetlands that were examined include the extent of lands that were potentially historic wetlands, the relationship between these and the wetlands that currently exist within the state of Indiana, the relationship between road proximity and the prevalence of wetlands, and the relationship between farming intensity and wetlands and forest cover and wetlands. This research does not seek to determine any qualitative effects of road construction and proximity of roads on wetlands.

Findings

This research was carried out primarily with the use of geographic information systems (GIS). GIS are digitized computer mapping systems that allow precisely scaled digitized maps to be linked to data/spreadsheets about the components (shapes, lines, points) with the maps. Maps using the same scale, datum and data storage format can be brought seamlessly together into a project. Maps can be layers and maps and data can be edited.

For this project a commercial program was used, ArcView GIS 3.2, a Windows-based GIS system developed by Environmental Systems Research Institute, Inc. (ESRI). Digital and non-digital data available on the web was used from several arenas including the USDA National Forest Service Inventory Analysis (FIA), the U.S. Dept. of Interior U.S. Fish and Wildlife Service National Wetlands Inventory (NWI), the U.S. Dept. of Agriculture National Resource Conservation Service (NRCS) soils data, the U.S.

Dept. of Agriculture 1997 Census of Agriculture, and Purdue University.

The results obtained with this study demonstrated that while wetlands prevalence (i.e., percent wetlands land area relative to total land area) decreased slightly in land areas close to state and federal roads (i.e., within the adjacent 500 to 1,000 ft. land reach), this change did not explain the dramatic decrease (in excess of 80%) in wetlands that has occurred in Indiana. In fact, hydric soil prevalence, a common indicator parameter for wetland areas, decreased only to a slight extent in land areas close to federal and state roads.

Since soil data are not expected to change much over time, the latter fact (i.e., the nominal decrease in hydric soil lands immediately adjacent to highways) indicates that the slight decrease in wetlands near state and federal roads is not due to adverse effects by road construction itself. Instead, this decrease may be due to roads being

constructed in areas and paths intended to avoid wetlands and hydric soils, thus lowering construction time and costs.

While the data developed in this project do not clearly indicated whether state and federal road construction has had an obvious quantifiable negative impact on wetlands, these data do not indicate that if there was an impact, it was a very small impact in comparison to the dramatic decline in wetlands that has been estimated to have occurred within the state of Indiana. This

decline could be due to many factors and research in this project has shown a correlation between wetlands loss as represented by the wetlands to hydric soil ratio and farming intensity and an opposite correlation between wetland loss and forest cover. The greater the farming intensity within a county, the greater was the estimated loss of wetlands. Also, a greater the area of forest cover within a county correlated with a lower estimated wetland loss.

Implementation

The topic of relative wetlands loss in the state of Indiana, and particularly the correlation between wetlands loss and land use in a general context, is an area in which more research could be performed. Two other factors that have not been examined in this project are the effect of county roads upon wetlands and the effect of urban sprawl upon wetlands.

These latter two factors likely played significant roles in wetland destruction. A problem lies in separating the effects on wetlands of agriculture, urban sprawl, state and federal roads, and county roads from each other. All of these developments occurred simultaneously and often overlap each spatially.

The lack of a consistently strong correlation of wetlands lying on hydric soils is also troubling. One of the key scientific foundations for this project was the part of the

definition of a wetland which requires it to be characterized by hydric soils. This correlation was considered to be the only way possible to gauge an estimate of historic wetlands.

However, the discrepancy between the wetlands definition and the fact that many NWI wetlands, which are based upon the wetland hydrology part of the wetland definition do not lie within SSURGO hydric soils, introduces a certain amount of error in this process. Conversely, NWI data are the best wetlands data available and the USDA SSURGO soils data are the best soils data available. This correlation between wetland presence and hydric soil characterization should, therefore, be given further research in conjunction with the future implementation of this project's current research findings.

Contacts

For more information:

Indiana Department of Transportation

Division of Research
1205 Montgomery Street
P.O. Box 2279
West Lafayette, IN 47906
Phone: (765) 463-1521
Fax: (765) 497-1665

Purdue University

Joint Transportation Research Program
School of Civil Engineering
West Lafayette, IN 47907-1284
Phone: (765) 494-9310
Fax: (765) 496-7996
<http://www.purdue.edu/jtrp>

		TECHNICAL REPORT STANDARD TITLE	
1. Report No. FHWA/IN/JTRP 2002/18	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Wetland Replacement Practices and Procedures for Indiana Highway Projects		5. Report Date June 2002	
		6. Performing Organization Code	
7. Author(s) J.E. Alleman and S. M. Graves		8. Performing Organization Report No. FHWA/IN/JTRP-2002/18	
9. Performing Organization Name and Address Joint Transportation Research Project Civil Engineering Building Purdue University West Lafayette, Indiana 47907-1284		10. Work Unit No.	
		11. Contract or Grant No. SPR-2205	
12. Sponsoring Agency Name and Address Indiana Department of Transportation State Office Building 100 North Senate Avenue Indianapolis, IN 46204		13. Type of Report and Period Covered Final Report	
		14. Sponsoring Agency Code	
15. Supplementary Notes Prepared in cooperation with the Indiana Department of Highways and Federal Highway Administration.			
16. Abstract <p>It is estimated that Indiana has already lost more than 80% of its original, native wetlands, and concerns have been expressed that some fraction, and perhaps even a sizable fraction, of these losses can be historically linked to highway construction associated with INDOT-related activities. These wetlands are important for several reasons, in that they help keep ground water and surface water clean, serve as reservoirs for floodwaters, help recharge groundwater, provide habitats for many species of animals, and serve as recreational areas. This underlying concern behind this accusation, therefore, is that the construction of state and federal roads historically carried out by INDOT has adversely affected wetlands quantitatively and qualitatively. Whether the accusation of a negative impact quantitatively is properly placed is the purpose of this research.</p> <p>Aspects of wetlands that were examined include the extent of lands that were potentially historic wetlands, the relationship between these and the wetlands that currently exist within the state of Indiana, the relationship between road proximity and the prevalence of wetlands, and the relationship between farming intensity and wetlands and forest cover and wetlands. This research, however, did not attempt to determine any qualitative effects of road construction and proximity of roads in relation to wetland damage and/or loss.</p> <p>This research effort was carried out primarily with the use of geographic information systems (GIS) using map data derived from a variety of sources, including: 1) the USDA National Forest Service Inventory Analysis (FIA); 2) the U.S. Dept. of Interior U.S. Fish and Wildlife Service National Wetlands Inventory (NWI); 3) the U.S. Dept. of Agriculture National Resource Conservation Service (NRCS) soils data; 4) the U.S. Department of Agriculture 1007 Census of Agriculture; and 5) Purdue University.</p>			
17. Key Words Wetland, loss, land, highway, construction, GIS		18. Distribution Statement No restrictions. This document is available to the public through the National Technical Information Service, Virginia, 22161	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages: 39	22. Price

ACKNOWLEDGMENTS

The authors would like to acknowledge the following individuals...

- **Barry Partridge**, INDOT, Chief, INDOT Division of Research
- **David Ward**, INDOT Section Manager,
*[INDOT Applied and Environmental Research, Electrical Engineering and
Technical Support]*

...for their technical assistance, professional insight, and financial support

TABLE OF CONTENTS

ACKNOWLEDGMENTS.....	ii
LIST OF FIGURES.....	iv
LIST OF TABLES	vi
ACRONYMS AND ABBREVIATIONS	vii
IMPLEMENTATION REPORT.....	viii
EXECUTIVE SUMMARY	ix
1. INTRODUCTION.....	1
2. PROBLEM STATEMENT	2
3 OBJECTIVES	5
4. DATA SOURCES.....	6
5 METHODS FOR ANALYSIS OF DATA	11
6 RESULTS	25
7. CONCLUSIONS.....	36
8. RECOMMENDATIONS	38
9. REFERENCES	39

LIST OF FIGURES

Number	Title	Page
1	NWI progress on a national wetlands mapping project	7
2	SSURGO data available online	9
3	Indiana rivers and streams shapefile with Tippecanoe County boundary superimposed.....	12
4	Indiana roads shapefile with Tippecanoe County boundary superimposed.....	12
5	Danville wetlands shapefile with Tippecanoe County border superimposed	13
6	Tippecanoe County rivers and streams shapefile	14
7	Tippecanoe County roads shapefile	14
8	Tippecanoe County wetlands shapefile.....	14
9	Close-up of rivers, wetlands, and roads shapefiles	14
10	Query process for selection of SSURGO soils that are hydric	15
11	Tippecanoe County SSURGO soils shapefile.....	16
12	Tippecanoe County hydric soils shapefile	16
13	U.S. 52 shapefile after selection and creation	17
	from the roads shapefile.	
14	State and federal roads shapefile with the roads shapefile.....	17
15	500 ft. buffer shapefile around state and federal roads	17
16	1000 ft. buffer shapefile around state and federal roads	17
17	Wetlands within 1000 feet of state and federal roads	18
18	Hydric soils within 1000 feet of state and federal roads	18
19	Laporte County wetlands	19
20	Laporte County hydric soils	19
21	Roads & state and federal roads.....	19

22	Close up of the city of Laporte with wetlands layered..... 19 above hydric soils with roads	19
23	1000 foot wetlands for Laporte County	19
24	1000 foot hydric soils for Laporte County	19
25	Wetland shapefiles downloaded for Buena Vista County..... 20 and brought into ArcView	20
26	Wetland themes merged into a single wetland shapefile	20
27	Clipped wetlands shapefile for Buena Vista County layered..... 21 above the merged wetlands shapefile	21
28	Wetlands shapefile layered above the Buena Vista County shapefile	21
29	SSURGO shapefiles downloaded for Buena Vista County	22
30	SSURGO shapefiles merged into a single SSURGO shapefile	22
31	Clipped SSURGO shapefile for Buena Vista County, IA.....	23
32	Hydric soils shapefile for Buena Vista County, IA.....	23
33	Wetlands of Wright County Minnesota	24
34	Hydric soils of Wright County Minnesota	24
35	Graph of wetlands and their prevalence near and away from roads	27
36	Graph of hydric soils and their prevalence near and away from roads	28
38	Relationship between wetland loss and forest cover.....	32
39	Southeastern Tippecanoe County wetlands.....	34
40	Southeastern Tippecanoe County hydric soils	34

LIST OF TABLES

Number	Title	Page
1	Results from Indiana counties concerning the prevalence of wetlands near state and federal roads	25
2	Results from Indiana counties concerning the prevalence of hydric soils near state and federal roads	25
3	Ratio of current wetlands to hydric soils, an indicator of potential historic wetlands	26
4	Results from counties around the eastern U.S. concerning the relationship between farming intensity and wetlands prevalence	29
5	Results from counties around the eastern U.S. concerning the relationship between woodland prevalence and wetlands prevalence	31
6	Percent of wetlands lying within hydric soils for several counties	33

ACRONYMS AND ABBREVIATIONS

CAAGIS	Center for Advanced Applications in GIS
EPA	US Environmental Protection Agency
ESRI	Environmental Systems Research Institute
FIA	Forest Inventory Analysis (USDA)
FWS	US Fish and Wildlife Service
GIS	Geographical information system
INDOT	Indiana Department of Transportation
NAD	North American Datum
NRCS	National Resource Conservation Service
NWI	National Wetlands Inventory (US_Fish and Wildlife Service)
SSURGO	Soil Survey Geographic Database
USGS	US Geological Survey
USDA	US Department of Agriculture

IMPLEMENTATION REPORT

The topic of relative wetlands loss in the State of Indiana, and particularly the correlation between wetlands loss and land use in a general context, is an area in which more research certainly could, and should, be performed in conjunction with the future implementation of this existing project. Two such important factors that have not been examined in this project are the effect of county roads upon wetlands and the effect of urban sprawl upon wetlands.

These latter two factors likely played significant roles in wetlands destruction. A problem lies in separating the effects on wetlands of agriculture, urban sprawl, state and federal roads, and county roads from each other. All of these developments have occurred simultaneously and often overlap each other spatially.

The lack of a consistently strong correlation of wetlands lying on hydric soils is also troubling. One of the key scientific foundations for this project was the part of the definition of a wetland which requires it to be characterized by hydric soils. This correlation was considered to be the only way possible to gauge an estimate of historic wetlands.

However, the discrepancy between the wetlands definition and the fact that many NWI wetlands, which are based upon the wetland hydrology part of the wetland definition, do not lie within SSURGO hydric soils introduces a certain amount of error in this process. Conversely, NWI data is the best wetlands data available and the USDA SSURGO soils data is the best soils data available. This correlation between wetland presence and hydric soil characterization should, therefore, be given further research relative to the future implementation of this project's current research findings.

EXECUTIVE SUMMARY

In Indiana, it is reported that the majority of original wetland area (approximately 85%) has been lost. As defined by the Indiana Wetlands Conservation Plan (DNR 1996), wetlands are qualified on the following basis:

“Wetlands are lands transitional between terrestrial and aquatic systems where the water table is usually at or near the surface or the land is covered by shallow water.”

These wetlands are important for several reasons, in that they help keep ground water and surface water clean, serve as reservoirs for floodwaters, help recharge groundwater, provide habitats for many species of animals, and serve as recreational areas. Indeed, wetlands serve many important functions, both in human benefits such as maintaining water quality and flood control and environmental benefits such as providing a habitat for endangered plants and animals. As a result, on a national level it has been recognized that a critical need exists to retain native wetlands areas whenever possible...and to replace wetlands lost or damaged with suitable new man-made wetlands areas.

Unfortunately, though, concerns have been expressed that some fraction, and perhaps even a sizable fraction, of these wetlands losses (i.e., ~85%) can be linked to highway construction associated with INDOT-related activities. While these lands could well have been drained and/or filled mostly for agriculture production, it is also possible that some degree of loss has also resulted from highway construction.

The underlying concern behind the accusation regarding INDOT's adverse historical impact on wetlands, therefore, is that the construction of state and federal roads carried out by INDOT might have negatively affected wetlands on both a quantitative and qualitative basis. Whether the accusation of a negative impact quantitatively is properly placed, or not, is the purpose of this research.

Aspects of wetlands that were examined include the extent of lands that were potentially historic wetlands, the relationship between these and the wetlands that currently exist within the state of Indiana, the relationship between road proximity and the prevalence of wetlands, and the relationship between farming intensity and wetlands

and forest cover and wetlands. This research, however, did not attempt to determine any qualitative effects of road construction and proximity of roads in relation to wetland damage and/or loss.

This research effort was carried out primarily with the use of geographic information systems (GIS) using map data derived from a variety of sources, including: 1) the USDA National Forest Service Inventory Analysis (FIA), 2) the U.S. Dept. of Interior U.S. Fish and Wildlife Service National Wetlands Inventory (NWI), 3) the U.S. Dept. of Agriculture National Resource Conservation Service (NRCS) soils data, 4) the U.S. Dept. of Agriculture 1997 Census of Agriculture, and 5) Purdue University.

As mandated by Presidential Executive Order, the Indiana Department of Transportation (INDOT) is now required to replace wetlands which are removed or impacted during highway construction on an acre-for-acre basis such that there is *no net loss* of surficial wetland area. Existing wetlands negatively harmed during these construction activities will either have to be totally replaced (e.g., creating entirely new sites) or enlarged (e.g., expanding an existing site). In all cases, the ratio of new wetland area to impacted wetland area must be greater than 2:1. For certain critical wetland areas, the replacement ratio may be as large as 4:1. The goal in either case is to create a self-sustaining habitat for aquatic plants and animals which provides the same function as the impacted wetland.

Currently, INDOT is involved in 50 to 60 wetland delineations/yr., which result in about 10 wetland mitigations/yr. The primary obstacle currently faced by INDOT with regard to wetlands is a lack of wetlands inventory data for Indiana. Simply stated, INDOT has experienced difficulty dealing with wetlands because it does not have accurate information on wetland locations, areas, and types. Such information is needed to minimize the impact of new construction and, where necessary, to optimize the mitigation actions for impacted wetlands.

The current available data sources (USGS quadrangle maps, aerial photos, U. S. Fish and Wildlife National Wetlands Inventory maps) are typically out of date or unconfirmed, and as a result are reliable only for larger wetlands (greater than 20 acres). Thus, the process of highway planning, design, and construction may be delayed until

time-consuming field reconnaissance is completed. INDOT is one of many organizations that are affected by the lack of reliable wetlands inventory information.

For this reason, the Indiana Task Force on Wetlands has been created and includes members from INDOT (Steve Cecil), Indiana Department of Environmental Management (IDEM), the Indiana Department of Natural Resources (IDNR), industry, conservation groups, and community groups. The Indiana Task Force on Wetlands has identified the need for updated wetland inventory information as a top priority. Likewise, one of the objectives of the Indiana Wetlands Conservation Plan (p. 44) is to *"have an inventory system capable of quantitatively identifying and monitoring Indiana's wetlands in place by May 1998"*.

In some areas of the U. S., wetland replacement is practiced on an in-kind basis in which the vegetation of the replacement area must be the same as that lost to a project. This approach is straightforward, but it assumes that wetland functions correlate with vegetation structure, which may not in actuality be the case. In-kind replacement also may not take into consideration the importance of site location in replicating the functions of the wetland. Thus, the current trend is away from in-kind vegetation replacement and towards the concept of wetland functional replacement. This involves a more detailed consideration of site selection and design, the type and condition of underlying soils, and surface water and groundwater quality at these sites (FHWA 1990).

Public concern regarding the preservation of remaining wetlands has resulted in the enactment of federal, state, and local laws. Examples of important federal mandates include sections of the Clean Water Act, the Coastal Zone Management Act, and the National Environmental Policy Act. In addition, a wide range of policies and regulations have been enacted at the state level. Wetland laws, regulations, and policies vary from state to state. Many have specific mitigation policies relating to wetland replacement, including the required replacement ratio, the location of the mitigation site (off-site vs. on-site), as well as the type of mitigation allowed (in-kind vs. out-of-kind).

1. INTRODUCTION

This project originated from the need to determine what has led to the loss of wetlands within the state of Indiana. In particular INDOT, the Indiana Department of Transportation, is an organization that has been blamed for some of the wetlands destruction within the state. This accusation assumes that the construction of state and federal roads carried out by INDOT has adversely affected wetlands quantitatively and qualitatively. Whether the accusation of a negative impact quantitatively is properly placed or not is the purpose of this research. Aspects of wetlands that were examined include the extent of lands that were potentially historic wetlands, the relationship between these and the wetlands that currently exist within the state of Indiana, the relationship between road proximity and the prevalence of wetlands, and the relationship between farming intensity and wetlands and forest cover and wetlands. This research does not seek to determine any qualitative effects of road construction and proximity of roads on wetlands.

This research was carried out primarily with the use of geographic information systems, GIS. GIS are digitized computer mapping systems that allow precisely scaled digitized maps to be linked to data/spreadsheets about the components (shapes, lines, points) within the maps. Maps using the same scale, datum and data storage format can be brought seamlessly together into a project. Maps can be layered and maps and data can be edited. For this project a commercial program was used, ArcView GIS 3.2, a windows based GIS system developed by ESRI, Environmental Systems Research Institute, Inc. Digital and non-digital data available on the web was used from several arenas including the USDA National Forest Service Inventory Analysis (FIA), the U.S. Dept. of Interior U.S. Fish and Wildlife Service National Wetlands Inventory (NWI), the U.S. Dept. of Agriculture National Resource Conservation Service (NRCS) soils data, the U.S. Dept. of Agriculture 1997 Census of Agriculture, and Purdue University.

2. PROBLEM STATEMENT

It is estimated that Indiana has already lost about 87% of its original wetlands. These wetlands are important for several reasons, they help keep ground water and surface water clean, serve as resevoirs for floodwaters, help recharge groundwater, provide habitats for many species of animals, and serve as recreational areas (Miller and MacGowan). Originally the idea behind this project was to develop an historical wetlands map of a couple of Indiana counties, including Tippecanoe county, in a time period approximately 50 years ago with the use of aerial photographs from the 1940's or 1950's and a 1940's Map of the Soil Associations in Indiana put out by the Soil Conservation Service. This historical wetlands map would then be used in comparison against a current wetlands map and a map of state and federal roads. Several difficulties occurred with this strategy.

First the 'Map of the Soil Associations in Indiana' had no classification system to precisely identify wetlands. This is partially due to wetland preservation and identification not becoming important in this country until the 1970's and later. The best approximation using the Soil Association map was several classifications identified by a common theme, muck, but this classification was never determined to have a direct correlation to the present definition and classification scheme for wetlands. Another problem stemmed from the Soil Association map not being digitized and with the level of detail existing in this map the process to digitize that map would be extremely difficult and time consuming, far beyond the scope for this project.

A second problem stemmed from the aerial photographs. It was believed that these might be of some use in identifying wetlands as they existed 50 years ago but examination of them led to the conclusion that it was not possible to distinguish many wetlands from forests. "...the most common wetland types in Indiana are forested wetlands and small temporarily flooded wetlands in farmed fields." (Miller and MacGowan). So a new approach was needed. One note of interest that the aerial photographs did reveal was that many of the state and federal roads in Tippecanoe County were already constructed by the 40's and 50's. Thus the value of these aerial

photographs is further limited since they would not indicate wetland extent before state and federal road development.

The current technical and federal government definition of wetlands is "...areas that have a predominance of hydric soils and that are inundated or saturated by surface or ground water at a frequency and duration sufficient to support a prevalence of hydrophytic vegetation typically adapted for life in saturated soil conditions." (NRCS Distribution of Wetlands).

"Wetlands have three essential characteristics: hydric soils, hydrophytic vegetation, and wetland hydrology. Hydrophytic vegetation is plant life growing in water or in soils that are periodically deficient in oxygen as a result of saturation of the soil with water, such as cattails, sedges, and willows. Wetland hydrology refers to periodic inundation (flooding or ponding) or saturation to the soil surface, usually for a week or more. Hydric soils are those that are saturated, flooded, or ponded long enough during the growing season to develop anaerobic conditions in the upper part." (Franzmeier and Kladvko)

This definition of hydric soils is confirmed on the USDA – NRCS web page entitled Hydric Soils of the United States located at the following Web site location:

<http://www.statlab/iastate/edu/soils/hydric/intro.htm>.

The Army Corps of Engineers requires all three parts of the definitions of a wetland be used in the determination and delineation of wetlands. The first part of the wetland definition, hydrophytic vegetation, while a useful current delineation and classification tool, posed a problem with acquiring historical vegetation data. No digitized historical vegetation data exists that could be found. Nor was any digitized data on historical wetlands hydrology found. Interestingly the current National Wetlands Inventory uses wetland hydrology as its primary tool for identifying current wetlands (Santos).

Hydric soils were determined as the best tool for mapping historical wetlands. As it turns out hydric soils are readily obtained from USDA SSURGO digitized data that is available for download from the web. Using the USDA digital soils map then the best initial estimate of an historical wetlands map would be to select all the hydric soils, one part of the federal definition of a wetland (Harbor). Approximately 26% of Indiana soils are hydric (Franzmeier and Kladvko).

“Before any drainage systems were installed, practically all of these soils would have qualified as wetlands by the current definition. Recent observations in undrained areas, however, show that hydrophytic vegetation does not extend quite as high in the landscape as do hydric soils. Thus at the time of European settlement, about 25% of Indiana would have qualified as wetlands according to the current technical definition.” (Franzmeier and Kladivko).

While not a precise gauge of historical wetlands, hydric soils was the best method found for approximating the wetlands that existed prior to settlement of and development of the lands in Indiana. Use of hydric soils for this purpose is consistent with the definition of wetlands.

Using hydric soils is not a good measure of wetlands in the 1940's because many wetlands had already been drained by that time as Indiana had been settled for over 100 years. Also, as has been pointed out previously, state and federal road development had already been active prior to the 1940's. Hydric soils are a better measure of the effect of road development on wetlands as they existed prior to any road or agricultural development, which significantly predates the 1940's. There are many factors that could explain the loss of wetlands besides state and federal road construction, such as county road construction, urban sprawl, deforestation, and agricultural development. Since all of these factors occurred concurrently it is difficult to separate them from one another but this project has attempted to develop some correlation between state and federal road construction, agricultural prevalence, and the extent of forest cover to the decline in wetlands.

3. OBJECTIVES

The following list summarizes the primary objectives for this research effort:

- to develop a methodology for accurately quantifying the location and area of existing wetlands in selected state regions using current spatial imaging resources (e.g., GIS analysis of National Wetlands Inventory maps, etc.),
- to devise a system for determining a historical correlation between highway location and current historical wetland area shifts,
- to test the accuracy of the methodology and system by actual field verification of selected wetland sites in two specific counties, and
- To investigate and characterize the accusation regarding INDOT's adverse historical impact on wetlands, whereby the construction of state and federal roads carried out by INDOT might have negatively affected wetlands on both a quantitative and qualitative basis. Whether the accusation of a negative impact quantitatively is properly placed, or not, is the purpose of this research.

4. DATA SOURCES

The digital wetlands data used in this project originated from the National Wetlands Inventory, NWI, whose web site is located at <http://www.nwi.fws.gov/>. NWI, a division of the U.S. Fish and Wildlife Service, FWS, lies within the U.S. Department of the Interior, and "... produces information on the characteristics, extent, and status of the Nation's wetlands and deepwater habitats." (NWI homepage.) In 1986 the U.S. Congress passed the Emergency Wetland Resources Act of 1986 which instructed the Fish and Wildlife Service to develop maps and a digital database of the wetlands of the United States. To date the NWI has mapped 89% and digitized 39% of the wetlands of the lower 48 states. "This information is used by Federal, State, and local agencies, academic institutions, the U.S. Congress, and the private sector." (NWI Overview.)

"The source material used to produce the National Wetlands Inventory (NWI) digital data for these maps was prepared primarily by stereoscopic analysis of high altitude photographs based on vegetation, visible hydrology, and geography in accordance with the Classification of Wetlands and Deepwater Habitats of the United States." (Santos). NWI wetlands maps for Indiana were developed using "...color infrared photography from the early to mid 1980's, and some field verification. Soil surveys were used as collateral data only." (Santos). Wetlands that have been mapped and those that have been digitized and are available for downloading are presented on the following map located at <http://www.ftw.nrcs.usda.gov/jpg/archived.jpg>.

To download NWI data these series of sites were used. From the NWI site at <http://www.nwi.fws.gov/> a link titled Wetlands Interactive Mapper was used. This links to http://wetlands.fws.gov/mapper_tool.htm where the user must choose data from either the eastern half or the western half of the nation. Only states east of Kansas City were used for this project. Thus the eastern half of the nation is selected from the St. Petersburg, FL link. This web site, http://wetlands2.nwi.fws.gov/nwi_mapplet/, brings up a form where a county name must be input. Upon selecting the proper county state combination (there are often several counties with the same name but in different states) the submit button is depressed and a wetlands map of that county and surrounding areas is accessed. County, wetland, road, and river data can be downloaded from this site in two file formats. The easiest to use is the ESRI format files which can brought directly into ArcView GIS as a theme or shapefile. But first these downloaded files, which are zip files, must be unzipped. Files throughout this project were unzipped using WinZip 7.0.

Digital soil data was acquired from the National Resources Conservation Service, NRCS, which is an agency within the U.S. Department of Agriculture, USDA. The need for the NRCS, originally the Soil Conservation Service, grew out of the Dust Bowl era of the 1930's when soil erosion suddenly became a serious concern. The mission of the NRCS is to facilitate a partnership between federal agencies and local communities to encourage conservation of national resources such as land, soil, air, water, plants and animals. The NRCS encourages responsible stewardship of these resources by private landowners, such as farmers and ranchers, and provides technical assistance to these landowners. (NRCS at a Glance).

The NRCS heads the National Cooperative Soil Survey, NCSS, which maintains three soil geographic databases "...for collecting, storing, maintaining, and distributing soil survey information for privately owned lands in the United States." (NRCS SSURGO Data Users Guide). These three databases are the Soil Survey Geographic database, SSURGO, the State Soil Geographic database, STATSGO, and the National Soil Geographic database, NATSGO. NATSGO data is intended for use in national resource planning, while STATSGO data is intended for state resource planning. SSURGO data provides the most detailed information and was designed for local planning such as counties, townships, farmers, ranchers, and other landowners. SSURGO data was chosen for this project due to its level of detail and its identification of soils as being hydric or non-hydric. SSURGO is the only soil database with a sufficient level of detail for use with this project.

The counties in the US where digital SSURGO data is available for downloading is presented in the following map located at <http://www.ftw.nrcs.usda.gov/jpg/archived.jpg>.

Figure 2: SSURGO data available online.

SSURGO digitized data is available from these web pages. <http://www.nrcs.usda.gov/TechRes.html> is the technical resources page for the NRCS. From here soils data is accessed via the “Soils Data” link, <http://www.statlab.iastate.edu/soils/nsdaf/> which is a web page that links to various different soil formats. One of these is SSURGO data, accessed via the “SSURGO” link http://www.ftw.nrcs.usda.gov/ssur_data.html. On this page several links are available, the detailed PDF format ‘User’s Guide’ description of SSURGO data, the national archived county map (pictured above), and a link to download SSURGO Data. At this point an account with NRCS is required, which is free, and then SSURGO data is available for download by county. There are close to 1000 counties currently in the database, with more being added all the time, which is arranged alphabetically by state abbreviation then county.

A convenient source of data to download for the state of Indiana is through the Center for Advanced Applications in GIS, CAAGIS, of Purdue University located at this web site, <http://danpatch.ecn.purdue.edu/~caagis/ftp/gisdata/data.html>. Here SSURGO data can be downloaded for six of the eleven counties in Indiana that have had their soils

data digitized by the NRCS. Other useful data available for download at this site include roads, railroads, rivers and streams, county boundaries, and others. Wetlands data acquired from NWI can also be downloaded from this site.

Early on in this project the advantage of using this site is that all downloadable data can be acquired in NAD 83 format. All CAAGIS data is available in both NAD 83 and NAD 27, except for wetlands, which like the NWI download site, is only available in NAD 83. SSURGO data available from the NRCS download site was found early in this project to be in NAD 27, but more recent downloads have resulted in files in NAD 83 format. NAD stands for North American Datum and is a reference against which the maps are constructed. NAD 27 is the old standard developed in 1927. NAD 83 is the current U.S. standard developed in 1983 and is GPS compatible. For Indiana, if NAD 27 and NAD 83 soils data for the same county are brought together into a map and layered on top of each other, an approximate 100 m north south shift results between the two data sets. This separation of the data means that map layers are not precisely placed on top of one another and limits the level of editing and calculations that can be performed on the maps.

There are methods to convert a data file from one datum to another datum, but the most reliable method is in a program called ArcInfo, which was not learned for this project. The newest version of ArcView contains a tool for this conversion but it is reported to be unreliable and was also not learned for this project.

Another source of data used in this project was the USDA 1997 Census of Agriculture located at <http://www.nass.usda.gov/census/census97/volume1/vol1pubs.htm>. At this site a wealth of statistical data is available on state and county agriculture. For this project data gathered included county area, percent of county that is farmland, and farm acreage within the county.

Also used was the USDA National Forest Service, Forest Inventory and Analysis (FIA), located at <http://fia.fs.fed.us>. From here the state where data is desired is selected and the FIA table generator is used to pull up the information desired. For this project the data gathered was the forest cover for each county analyzed.

5. METHODS FOR ANALYSIS OF DATA

Two methods were used to develop useful maps and data for this project. The first entailed using CAAGIS data to develop detailed GIS maps of several counties in Indiana. Each final map included the county itself and within the county state and federal roads, railroads, all county roads, rivers and streams, total hydric soils, hydric soils within 1000' of state and federal roads, hydric soils within 500' of state and federal roads, total wetlands, wetlands within 1000' of state and federal roads, wetlands within 500' of state and federal roads, and areas common to both hydric soils and wetlands. This method was used primarily to examine the relationship between loss of wetlands and the proximity to state and federal roads. Unfortunately much of this work had to be repeated using data from NWI and NRCS due to some inconsistencies of the CAAGIS data.

The second method employed entailed using NWI data, NRCS data, FIA data, and USDA Census data to develop GIS maps of other counties around the nation. Each final map included the county itself, total wetlands, total hydric soils, and the areas common to both wetlands and hydric soils. This data was used to examine the relationship between current and estimated historical wetlands to the level of agricultural activity within the county and the forest cover of each county. This method was initially used to look at only counties outside of Indiana, but the later revised versions of the Indiana counties examined in detail were constructed using this same method. Thus this second method proved to be the method of choice mainly due to the format of the NRCS and NWI data being more reliable, i.e. in a consistent NAD format. In these revisions CAAGIS roads and streams data was still used, but county boundaries, wetlands, and hydric soil data was acquired elsewhere. A detailed description of the steps involved in performing both methods, creating both types of GIS maps, follows.

The path for constructing detailed maps of each Indiana county was the same except for the data sets used. The steps in developing the Tippecanoe County map data will be presented. First the data must be downloaded from the CAAGIS web site, <http://danpatch.ecn.purdue.edu/~caagis/ftp/gisdata/data.html>. Files that were downloaded included counties, roads, railroads, rivers and streams, wetlands from the Danville section

of the Indiana Map since Tippecanoe county is located in that section, and Tippecanoe county NAD 83 SSURGO soils. Each file is a zip file and must be unzipped.

Arcview GIS uses shapefiles to represent area shapes in a map. Each shapefile can contain thousands of shapes and acts as a separate layer or theme in an ArcView project. The rivers and streams shapefile and the roads shapefile downloaded for this project encompass the entire state of Indiana. The Wetlands shapefile encompassed a large section of central Indiana and central Illinois.

Figure 5: Danville wetlands shapefile with Tippecanoe county border superimposed.

ArcView GIS uses what are called extensions to perform various functions that the basic program cannot do. An extension must be activated or brought into the ArcView project in order to be able to use it. Geoprocessing Wizard is an extension that was used extensively in this project. This extension allows themes to be merged, intersected, unioned, clipped, or dissolved with each other. The Tippecanoe county shapefile was used to clip the desired areas of all three themes pictured above. This clipping resulted in streams, roads, and wetland shapefiles for just the Tippecanoe county area. These three shapefiles are pictured below.

Figure 6: Tippecanoe county rivers and streams shapefile.

Figure 7: Tippecanoe county roads shapefile.

Figure 8: Tippecanoe county wetlands shapefile.

Figure 9: Closeup of rivers, wetlands, and roads shapefiles of West Lafayette and Lafayette.

Every shapefile or theme in ArcView has a table or spreadsheet of data associated with it. Every shape within the shapefile constitutes one row in the table. The columns

are made up of attributes such as area, perimeter, etc. The SSURGO soils shapefile was used to develop a map of hydric soils because the SSURGO table has an attribute called 'hydric' which indicated whether a particular shape (corresponding to an area of land) in the soils shapefile is hydric or not. This was accomplished by using an ArcView GIS function called Query Builder which allows the search of a shapefile's attributes. In this case all shapes whose hydric attribute was "Y" was selected (highlighted in yellow) as can be seen in the figure below. This group of selected shapes was then converted into a new shapefile which is the hydric soil shapefile pictured below.

Figure 10: Query process for selection of SSURGO soils that are hydric.

<p>Figure 11: Tippecanoe county SSURGO soils shapefile.</p>	<p>Figure 12: Tippecanoe county hydric soils shapefile.</p>

The next step was to create a shapefile of the state and federal roads within Tippecanoe County. The Tippecanoe county roads shapefile contains all roads within the county with no way of easily separating state and federal roads from county roads. Thus the only way to identify state and federal roads is to go in and individually identify and select each shape, a line segment in this case, that is part of a state or federal road. For a reference a Tippecanoe county fold out map, published by UniversalMAP and available in local gas stations and convenient stores, was used. Each road consists of many line segments and the level of detail requires extensive use of the zooming features of ArcView GIS to be able to see clearly. After an entire stretch of road is highlighted using the selection feature in ArcView GIS, it is converted to a new shapefile. After all the state and federal road shapefiles have been created then they are unioned into one single shapefile. The figures below show parts of this time consuming process.

Figure 13: U.S. 52 shapefile in red after selection and creation from the roads shapefile in gray.

Figure 14: State and federal roads shapefile in red with the roads shapefile in gray.

Next 500 feet and 1000 feet buffer shapefiles were created around state and federal roads. This was accomplished using the create buffers function of ArcView GIS.

Figure 15: 500 ft. buffer shapefile around state and federal roads.

Figure 16: 1000 ft. buffer shapefile around state and federal roads.

The purpose of the buffer shapefiles is to gather information about the state of wetlands within 500 feet and 1000 feet of state and federal roads. This was accomplished by clipping the wetlands and hydric soil shapefiles with each buffer and creating new shapefiles.

After this process is completed, several useful shapefiles exist for Tippecanoe County. These include wetlands, wetlands within 500 feet of state and federal roads, wetlands within 1000 feet of state and federal roads, hydric soils, hydric soils within 500 feet of state and federal roads, hydric soils within 1000 feet of state and federal roads, and the county shapefile. Another ArcView GIS extension was used at this point, Xtools extension – meters/acres version. Xtools contains a valuable tool that allows the area of each shape within a shapefile to be calculated. For example the area of each wetland in the wetlands shapefile can be calculated with Xtools. The area of these buffer shapefiles were determined by first clipping the buffers with the county border and then calculating the area of the buffer using Xtools.

The next step is to quantify the area data in a spreadsheet. Using Microsoft Excel the table associated with each shapefile, a .dbf file, can be opened and converted to excel files if desired. A basic sum operation can be run on the column of data in the table that contains the area for each shape in the shapefile. For example the area of all of the

wetlands within Tippecanoe County can be found in this fashion. This completes the first method for data analysis.

This same process was carried out for Laporte County and other counties in Indiana. The following images show some results for Laporte County.

The second method entails acquiring wetland and hydric soil data for counties around the nation. CAAGIS data could not be used for this purpose because CAAGIS only covers Indiana counties. Thus NWI wetlands and NRCS SSURGO data was used. For demonstration purposes, the development of data for Buena Vista County in Iowa will be discussed as a representative case.

After proceeding through the NWI web pages mentioned above in the source section of this paper, selecting the proper county, wetlands and county ESRI files were downloaded in a zip file and unzipped. The unzipped files are then brought directly into the ArcView project for Buena Vista County by adding new themes. The wetland files must then be merged into one theme using Geoprocessing Wizard.

Figure 25: Wetland shapefiles downloaded for Buena Vista county and brought into ArcView.

Figure 26: Wetland themes merged into a single wetland shapefile.

Next the Buena Vista County shapefile is used to clip the merged wetlands shapefile creating a new shapefile of the wetlands within Buena Vista County. This clipped wetlands shapefile contains shapes for both wetlands and uplands which are non-wetlands. Using the Query Builder all of the shapes that are uplands, where attribute equals "U", are selected. The reverse selection tool was then used so that all other shapes, the wetlands, are selected and all uplands are unselected. These selected wetlands are then converted to a new shapefile, which contains only wetlands for Buena Vista County.

Figure 27: Clipped wetlands shapefile (orange) for Buena Vista county layered above the merged wetlands shapefile (purple).

Figure 28: Wetlands shapefile (green) layered above the Buena Vista county shapefile (gray).

An hydric soils shapefile was developed next. The page for locating the download button for data from Buena Vista County was found using the steps mentioned earlier in the source section of this paper. The zip files downloaded were cov.zip, an ArcInfo coverage file, which is basically the digitized map, and tab.zip, which is a table or spreadsheet of data about shapes, lines, and points, in the coverage file. These files were then unzipped.

The coverage file contains many different files, but three basic files exist. Point shapefiles end in a 'p', line shapefiles end in a 'l', and area shapefiles end in an 'a'. It is the area shapefiles that were important for this project and these were added to the Buena Vista ArcView project as new themes/shapefiles. These shapefiles must be merged using the Geoprocessing Wizard.

The county shapefile acquired earlier with the NWI data was then used to clip the SSURGO data using the ArcView extension Geoprocessing Wizard. This produces a shapefile of the soils within Buena Vista County.

Unfortunately the table files associated with the SSURGO shapefiles have limited data. They contain no information on whether the soil is hydric or not. This is why the tab.zip file was also downloaded for Buena Vista County. This tab file, short for table, contains several tables related to the SSURGO soil shapefiles, but they must first be brought into ArcView GIS and joined to the limited table the SSURGO soil shapefile already possesses. To accomplish this the table titled “comp”, which contains hydric information, is opened in Microsoft Excel. This table is then saved as comp.dbf so it can be opened in ArcView. After the comp.dbf table is opened in ArcView it must be joined to the merged SSURGO shapefile’s table. This process is too complicated to describe here, but once the two tables are joined the merged SSURGO shapefile contains a single larger table which now contains the hydric information.

Using Query Builder on the new table, those soils where the hydric attribute equals “Y” for yes or “U” for unknown is selected. This highlighted data was converted to a new shapefile, the hydric soils for Buena Vista County.

A note of some importance is discussed here. The “U” data was selected even though it is not hydric because examination of SSURGO shapefiles indicated that this value was used to represent large bodies of water, mainly lakes. Why these soils are not classified as hydric was not determined, however they are also not classified as “N”, non-hydric. In addition these large bodies of water are classified as wetlands in the NWI data. Since the point of this method is to compare areas of hydric soils to that of wetlands it was determined that leaving out large areas that are certainly hydric by the nature of lying underwater would skew the results. Thus these soils were added to the hydric soils shapefile.

<p>Figure 31: Clipped SSURGO shapefile for Buena Vista county, IA.</p>	<p>Figure 32: Hydric soils shapefile for Buena Vista county, IA.</p>

Next the Xtools extension was used to calculate the areas of the county shapefile, wetlands shapefile, and hydric soils shapefile. Microsoft Excel was then used to access the .dbf table files associated with these shapefiles and total areas of all wetlands and all hydric soils could be calculated.

This method was carried out on thirteen other counties in various states around the eastern half of the nation. The results for another county, Wright County in Minnesota, is pictured below.

Due to some discrepancies with some of the CAAGIS data, the second method described above was used in the revisions of the Indiana counties examined in detail, previously performed using method one. County and wetland themes were created using NWI data. Soils and hydric soils data was created using NRCS SSURGO data. CAAGIS data that was still used included, roads, streams and railroads data. Also the process for creating themes of state and federal roads, 500' and 1000' buffers of these roads, and wetlands and hydric soils within these buffers remains the same as described above in the first method.

6. RESULTS and DISCUSSION

The results for the Indiana counties examined in detail are listed below:

COUNTY	County Area (acres)	Area of 500' Buffer Around Roads (acres)	Area of 1000' Buffer Around Roads (acres)	Wetlands w/in 500' Buffer (acres)	Wetlands within 500' Buffer/ Area of 500' Buffer (%)	Wetlands w/in 1000' Buffer (acres)	Wetlands within 1000' Buffer/ Area of 1000' Buffer (%)	Total Wetlands (acres)	Total Wetlands/ County Area (%)
Carroll	239850	12246	24152	265	2.2%	700	2.9%	8626	3.6%
Fountain	254465	11362	22298	251	2.2%	579	2.6%	10474	4.1%
Elkhart	299246	23820	46466	806	3.4%	2322	5.0%	17131	5.7%
Laporte	386443	28676	55879	1125	3.9%	3006	5.4%	27821	7.2%
Scott	123284	9360	18150	302	3.2%	794	4.4%	8514	6.9%
Tippecanoe	321815	20667	40158	538	2.6%	1370	3.4%	13155	4.1%
Vermillion	166232	11362	22298	251	2.2%	579	2.6%	6793	4.1%
Wayne	258756	16948	33088	356	2.1%	884.5	2.7%	7393	2.9%

Table 1: Results from Indiana counties concerning the prevalence of wetlands near state and federal roads.

COUNTY	County Area (acres)	Area of 500' Buffer Around Roads (acres)	Area of 1000' Buffer Around Roads (acres)	Hydric Soils w/in 500' Buffer (acres)	Hydric soils within 500' Buffer/ Area of 500' Buffer (%)	Hydric Soils w/in 1000' Buffer (acres)	Hydric soils within 1000' Buffer/ Area of 1000' Buffer (%)	Total Hydric Soils (acres)	Total Hydric soils/ County Area (%)
Carroll	239850	12246	24152	3168	25.9%	6498	26.9%	67851	28.3%
Fountain	254465	11362	22298	1894	16.7%	3789	17.0%	57889	22.7%
Elkhart	299246	23820	46466	5011	21.0%	10651	22.9%	73055	24.4%
Laporte	386443	28676	55879	7528	26.3%	15647	28.0%	134684	34.8%
Scott	123284	9360	18150	4447	47.5%	8565	47.2%	57569	46.7%
Tippecanoe	321815	20667	40158	4261	20.6%	8726	24.7%	81230	25.2%
Vermillion	166232	11362	22298	1894	16.7%	3789	17.0%	30671	18.5%
Wayne	258756	16948	33088	1528.7	9.0%	3227	9.8%	28866	11.2%

Table 2: Results from Indiana counties concerning the prevalence of hydric soils near state and federal roads.

COUNTY	Ratio of Wetlands within 500' to Hydric Soils Within 500' of state & fed roads	Ratio of Wetlands within 1000' to Hydric Soils Within 1000' of state & fed roads	Ratio of total county wetlands to total county hydric soils
	(%)	(%)	(%)
Carroll	8%	11%	13%
Fountain	13%	15%	18%
Elkhart	16%	22%	23%
Laporte	15%	19%	21%
Scott	7%	9%	15%
Tippecanoe	13%	16%	16%
Vermillion	13%	15%	22%
Wayne	23%	27%	26%

Table 3: Ratio of Current Wetlands to Hydric Soils, an indicator of potential historic Wetlands.

With hydric soils representing the potential for wetlands to have existed in the past, the wetlands to hydric soil ratio provides an indicator of what percentage of wetlands exist today as compared to what may have existed prior to European settlement. In the counties examined the wetlands to hydric ratios are small ranging from 26% to 13%. This represents an approximate 80% decline in the amount of wetlands that may have existed at one time.

When only those wetlands and hydric soils near state and federal roads, those INDOT is responsible for, are looked at, the results are similar, but with a further decline in the ratio as proximity to the road increases. This could be do to a couple of factors. One possibility is that road construction has had a negative impact on the prevalence of wetlands. Where road construction went through a wetland, then that land had to be drained and made into a suitable roadbed. But since hydric soils prevalence as well as wetlands prevalence decreases with the proximity of roads, this could also indicate that road construction was undertaken through areas with fewer hydric soils, and thus fewer wetlands, thereby minimizing construction cost and effort required. Whether a soil is an hydric or not should not change over time. The fact that hydric soil prevalence does decrease as proximity to roads increases supports the idea that state and federal roads were, to some degree, constructed to avoid hydric soils/historic wetlands. The following two figures summarize the prevalence of wetlands and hydric soils as they near state and federal roads.

But in any case, the data indicates that the majority of wetland loss occurred separate from state and federal road construction. Countywide the potential losses range from 74% to 87%, a dramatic decline. Proximity to roads only increases this loss only another few percent, a figure that pales in comparison to the huge loss already indicated.

The next analysis which centered on examining agricultural intensity within a county and comparing that to the amount of wetlands remaining in that county from what potentially existed at one time is summarized in the figure below.

State	County	area (acres)	wetlands (%)	farm (acres)	county area (including lakes) (acres)	hydric (acres)	wetlands (acres)	wetlands (%)	hydric (%)	(hyd/total) (%)	(wet/total) (%)
Illinois	Edgar	399095	88.3	352,401	399,198	148736	3689	2.5	37.3	0.9	
	Champaign	638205	89	567697	638,072	357219	6396	1.8	56.0	1.0	
	McHenry	386654	62.7	242484	390,947	108374	54266	50.1	27.7	13.9	
	Franklin	179588	68.1	263743	276,067	69517	38157	54.9	25.2	13.8	
Iowa	Buena Vista	367885	97	356751	371,206	133170	6128	4.6	35.9	1.7	
	Humboldt	278037	92.6	257411	278,706	158068	4600	2.9	56.7	1.7	
	Van Buren	310561	82.8	257227	313,760	83685	9031	10.8	26.7	2.9	
Minnesota	Sherburne	279403	37.6	117701	288,429	70594	60984	86.4	24.5	21.1	
	Wright	422934	59.5	251832	456,901	170574	141385	82.9	37.3	30.9	
Missouri	Wright	436701	71.5	312388	436,911	6496.5	2756	42.4	1.5	0.6	
New Jersey	Hunterdon	275261	38.2	105230	279,940	17481	10100	57.8	6.2	3.6	
New York	St. Lawrence	1718863	23.1	396406	1,383,314	292279	222107	76.0	21.1	16.1	
North Carolina	Hyde	392208	24.3	95327	443,965	444079	289879	65.3	100.0	65.3	
	Johnston	506868	41.6	211011	509,656	141103	68566	48.6	27.7	13.5	
Indiana	Laporte	382897	64.7	247756	386,443	134685	27821	20.7	34.9	7.2	
	Tippecanoe	319896	75.5	241532	321,707	81230	13155	16.2	25.2	4.1	
	Elkhart	296856	61.6	182771	299,246	73055	17131	23.4	24.4	5.7	
	Carroll	238251	91.6	218170	239,849	67851	8626	12.7	28.3	3.6	
	Delaware	251720	68.9	176443	253,304	94217	7329	7.8	37.2	2.9	
	Fountain	253264	80.8	204554	254,465	57889	10474	18.1	22.7	4.1	
	Scott	121856	47.1	57372	123,284	57569	8514	14.8	46.7	6.9	
	Vermillion	164418	71.8	118065	166,232	30671	6793	22.1	18.5	4.1	
	Wayne	258288	66.9	172860	258,756	28866	7393	25.6	11.2	2.9	
	Perry	244104	34.5	84251	247,066	6184	5506	89.0	2.5	2.2	
	Harrison	310561	52	161378	311,399	3023	4832	159.8	1.0	1.6	
	Owen	246538	43.5	107265	248,014	6306	8412	133.4	2.5	3.4	

Table 4: Results from counties around the eastern U.S. concerning the relationship between farming intensity and wetlands prevalence.

There are two chief figures, both highlighted, to observe in the above spreadsheet. The first is the percent farms which is a measure of how much of the land in that county is used as farmland or in other words a measure of the agricultural intensity within that county. The second figure to note is the wetlands to hydric soils ratio, which indicates the percent of current wetlands to that of the potential historic wetlands.

When these two factors are graphed against one another, a trend is seen. As the amount of farmland decreases, the amount of potential historic wetlands still in existence increases. While this relationship is not precisely linear, it would not be expected to be since there are many factors potentially affecting wetland losses.

The next analysis which centered on examining forest cover within a county and comparing that to the amount of wetlands remaining in that county from what potentially existed at one time is summarized in the figure below.

State	County	area (acres)	forest cover (acres)	forest cover (%)	county area including lake (acres)	hydric (acres)	wetlands (acres)	(wet/hydric) (%)	(hyd/total) (%)	(wet/total) (%)
Illinois	Edgar	399095	8,500	2.1	399,198	148736	3689	2.5	37.3	0.9
	Champaign	638205	0	0.0	638,072	357219	6396	1.8	56.0	1.0
	McHenry	386654	29100	7.4	390,947	108374	54266	50.1	27.7	13.9
	Franklin	179588	67200	24.3	276,067	69517	38157	54.9	25.2	13.8
Iowa	Buena Vista	367885	7900	2.1	371,206	133170	6128	4.6	35.9	1.7
	Humboldt	278037	2600	0.9	278,706	158068	4600	2.9	56.7	1.7
	Van Buren	310561	52000	16.6	313,760	83685	9031	10.8	26.7	2.9
Minnesota	Sherburne	279403	50700	17.6	288,429	70594	60984	86.4	24.5	21.1
	Wright	422934	43300	9.5	456,901	170574	141385	82.9	37.3	30.9
Missouri	Wright	436701	154800	35.4	436,911	6496.5	2756	42.4	1.5	0.6
New Jersey	Hunterdon	275261	96300	34.4	279,940	17481	10100	57.8	6.2	3.6
New York	St. Lawrence	1718863	1106300	80.0	1,383,314	292279	222107	76.0	21.1	16.1
North Carolina	Hyde	392208	235100	53.0	443,965	444079	289879	65.3	100.0	65.3
	Johnston	506868	240000	47.1	509,656	141103	68566	48.6	27.7	13.5
Indiana	Laporte	382897	59500	15.4	386,443	134685	27821	20.7	34.9	7.2
	Tippecanoe	319896	32700	10.2	321,707	81230	13155	16.2	25.2	4.1
	Elkhart	296856	28600	9.6	299,246	73055	17131	23.4	24.4	5.7
	Carroll	238251	31400	13.1	239,849	67851	8626	12.7	28.3	3.6
	Delaware	251720	9500	3.8	253,304	94217	7329	7.8	37.2	2.9
	Fountain	253264	42700	16.8	254,465	57889	10474	18.1	22.7	4.1
	Scott	121856	47000	38.1	123,284	57569	8514	14.8	46.7	6.9
	Vermillion	164418	36100	21.7	166,232	30671	6793	22.1	18.5	4.1
	Wayne	258288	21500	8.3	258,756	28866	7393	25.6	11.2	2.9
	Perry	244104	165100	66.8	247,066	6184	5506	89.0	2.5	2.2
	Harrison	310561	118300	38.0	311,399	3023	4832	159.8	1.0	1.6
	Owen	246538	106900	43.1	248,014	6306	8412	133.4	2.5	3.4

Table 5: Results from counties around the eastern U.S. concerning the relationship between woodland prevalence and wetlands prevalence.

There are two chief figures, both highlighted, to observe in the above spreadsheet. The first is the percent forest cover, which is a measure of how much of the land is covered with forest. The second figure to note is the wetlands to hydric soils ratio, which, as mentioned above, indicates the percent of current wetlands to that of the potential historic wetlands.

A trend can be seen here also, although once again not precisely linear. But as the forest cover, the natural state for most of these counties, increases the amount of potential historic wetlands remaining increases. Increased forest cover would indicate that the land has probably not been disturbed from its original state as much as a county where most of the forest cover has been removed. Minnesota and Iowa were left out of this data set due to these counties residing in predominantly prairie areas.

These two data sets indicate that a significant relationship exists between farming intensity and historic wetlands remaining and forest cover and historic wetlands remaining. Both are a measure of how much the land has been altered, though they approach the problem from different directions. However there are probably other factors at work. A couple of these could be urban sprawl and other road development,

meaning county roads. County roads make up many more miles of roads than do state and federal roads in most counties.

One other factor was looked at in this project. Since part of the definition of a wetland is that it contains a hydric soil it was decided to see how well this played out with wetlands data versus hydric soil data. Using the Geoprocessing Wizard of ArcView an intersection of the wetlands shapefile and the hydric soils shapefile was performed on the counties analyzed. This intersection results in a shapefile with land areas common to both wetlands and hydric soils. If the definition of a wetland holds up then most of the wetlands should exist on hydric soils. Unfortunately this experiment produced mixed results as can be seen in the spreadsheet below.

State	County	county area area (acres)	including lakes (acres)	hydric (acres)	wetlands (acres)	(wet/hydric) (%)	(hyd/total) (%)	(wet/total) (%)	hydric wetlands intersection (acres)	wetlands that are hydric (%)
Illinois	Edgar	399095	399,198	148736	3689	2.5	37.3	0.9	124	3.4
	Champaign	638205	638,072	357219	6396	1.8	56.0	1.0	5543	86.7
	McHenry	386654	390,947	108374	54266	50.1	27.7	13.9	49309	90.9
	Franklin	179588	276,067	69517	38157	54.9	25.2	13.8	25713	67.4
Iowa	Buena Vista	367885	371,206	133170	6128	4.6	35.9	1.7	5068	82.7
	Humboldt	278037	278,706	158068	4600	2.9	56.7	1.7	1816	39.5
	Van Buren	310561	313,760	83685	9031	10.8	26.7	2.9	5412	59.9
Minnesota	Sherburne	279403	288,429	70594	60984	86.4	24.5	21.1	50511	82.8
	Wright	422934	456,901	170574	141385	82.9	37.3	30.9	127709	90.3
Missouri	Wright	436701	436,911	6496.5	2756	42.4	1.5	0.6	238	8.6
New Jersey	Hunterdon	275261	279,940	17481	10100	57.8	6.2	3.6	6350	62.9
New York	St. Lawrence	1718863	1,383,314	292279	222107	76.0	21.1	16.1	105049	47.3
North Carolina	Hyde	392208	443,965	444079	289879	65.3	100.0	65.3	284148	98.0
	Johnston	506868	509,656	141103	68566	48.6	27.7	13.5	50095	73.1
Indiana	Laporte	382897	386,443	134685	27821	20.7	34.9	7.2	19862.7	71.4
	Tippecanoe	319896	321,707	81230	13155	16.2	25.2	4.1	6015	45.7
	Elkhart	296856	299,246	73055	17131	23.4	24.4	5.7	13964	81.5
	Carroll	238251	239,849	67851	8626	12.7	28.3	3.6	4929	57.1
	Delaware	251720	253,304	94217	7329	7.8	37.2	2.9	5014	68.4
	Fountain	253264	254,465	57889	10474	18.1	22.7	4.1	3259	31.1
	Scott	121856	123,284	57569	8514	14.8	46.7	6.9	4836	56.8
	Vermillion	164418	166,232	30671	6793	22.1	18.5	4.1	2430	35.8
	Wayne	258288	258,756	28866	7393	25.6	11.2	2.9	1482	20.0
	Perry	244104	247,066	6184	5506	89.0	2.5	2.2	208	3.8
	Harrison	310561	311,399	3023	4832	159.8	1.0	1.6	1205	24.9
	Owen	246538	248,014	6306	8412	133.4	2.5	3.4	2555	30.4

Table 6: Percent of wetlands lying within hydric soils for several counties.

Several counties produced favorable results with percentages in the 70's, 80's, and 90's. However some were quite low. Tippecanoe County only produced a value of 44%. It was thought that large rivers systems might be throwing off the data. An investigation was made for Tippecanoe county to see what would happen if a shapefile of all wetlands not touching a river or stream was created. This shapefile was then intersected with the hydric soil shapefile. This intersection however only produced a

value of 58% of wetlands lying on hydric soils. While this was larger than 44% it was still low and indicated either a problem with the definition of a wetland or the data sets available from the government.

A suspicion of the accuracy of NWI wetlands data had occurred earlier. An excursion to southeastern Tippecanoe County to observe some of these wetlands in person produced surprising results. Several NWI wetlands did not appear to be wetlands at all, but rather forests. In addition several of these wetlands/forests had houses built in the middle of them. A conversation with one resident indicated that ponding of water was never a problem. Examination of the hydric soil maps for these wetlands showed that several of these areas did not contain hydric soils. The two figures below show the wetlands and hydric soils of this area. The two circled areas indicate two of the wetlands/forests that were observed. Note that hydric soils are not found in the same area in which the wetlands are found.

NWI wetland hardcopy maps are qualified with these and other words. “The aerial photographs typically reflect conditions during the specific year and season when they were taken. In addition, there is a margin of error inherent in the use of the aerial photographs (Santos).” There are other factors that could have affected the quality of the data such as poor quality photography or forest stands on a particular day could have had

a very dark signature that would lead to it being mistaken for a wetland. Quality control steps, like limited field surveys and use of collateral data like soil surveys, are used to try to catch these problems. Also NWI data is derived from photos taken in the early to mid 80's, so this data does not reflect any changes since that time (Santos). It should also be noted that while wetland delineation work by the U.S. Army Corps of Engineers requires all three definitions of a wetland; hydric soil, wetland hydrology, and hydrophytic vegetation, be present, NWI mapping requires only the wetland hydrology criteria always be met (Santos).

The result is that NWI wetland data requires only the wetland hydrology criteria be met, this project used only the hydric soil criteria for an estimation of historic wetlands, and for true delineation of wetlands all three criteria need to be met. Thus some error is going to creep into this process. Unfortunately there was not found any source, hard copy or digital, that has delineated and archived the wetlands of the U.S. or Indiana. Delineation, according to the U.S. Army Corps of Engineers, proceeds on a case by case basis. When an area of land is developed or transformed and wetlands may be affected, then a delineation needs to be carried out. But there has been no national effort to delineate wetlands nationwide. The NWI data comes closest and is a very useful tool, but is not a substitute for the delineation process. Local, state, and federal regulatory agencies still have jurisdiction over wetlands.

7. CONCLUSIONS and RECOMMENDATIONS

The results obtained in this study demonstrated that while wetlands prevalence decreased slightly in land areas close to state and federal roads, this did not explain the dramatic decrease (in excess of 80%) in wetlands that has occurred in Indiana. In fact, hydric soil prevalence (i.e., a common indicator parameter for wetland areas) decreased only to a slight extent in land areas close to federal and state roads.

Since soil data is not expected to change much over time, the latter fact (i.e., the nominal decrease in hydric soil lands immediately adjacent to highways) indicates that the slight decrease in wetlands near state and federal roads is not due to adverse effects by road construction itself. Instead this decrease may be due to roads being constructed in areas and paths intended to avoid wetlands and hydric soils, thus lowering construction time and costs.

While the data developed in this project does not clearly indicate whether state and federal road construction has had an obvious quantifiable negative impact on wetlands, this data does indicate that if there was an impact it was a very small impact in comparison to the dramatic decline in wetlands that has been estimated to have occurred within the state of Indiana. This decline could be due to many factors and research in this project has shown a correlation between wetlands loss as represented by the wetlands to hydric soil ratio and farming intensity and an opposite correlation between wetland loss and forest cover. The greater the farming intensity within a county the greater the estimated loss of wetlands. Also the greater the area of forest cover within a county correlated with a lower estimated wetland loss.

This is an area in which more research could be performed. Two other factors that have not been examined in this project are the effect of county roads upon wetlands and the effect of urban sprawl upon wetlands. These are two factors that may have played significant roles in wetlands destruction. A problem lies in separating the effects on wetlands of agriculture, urban sprawl, state and federal roads, and county roads from each other. All of these developments have occurred simultaneously and often overlap each other spatially.

The lack of a consistently strong correlation of wetlands lying on hydric soils is troubling. The foundation for this project is the part of the definition of a wetland that requires it to be characterized by hydric soils. That was the only way found to gauge an estimate of historic wetlands. The discrepancy between the wetlands definition and the fact that many NWI wetlands, which are based upon the wetland hydrology part of the wetland definition, do not lie within SSURGO hydric soils introduces a certain amount of error in this process. However NWI data is the best wetlands data available and the USDA SSURGO soils data is the best soils data available.

A final note of importance is that this project is entirely based upon quantifiable data. Qualitative evaluations of state and federal road development effects upon wetlands is not the focus of this research, but has been touched upon in the research of others. Such research has sought to determine the effects of human land development and road construction on wetlands with the effect being a loss of biodiversity. A decline in biodiversity results from road construction due to wetland fragmentation, restricted movement between wetlands and species populations, roadkill, and increased human access to these ecosystems. Research has shown that much of this effect is not immediate but takes place over many years after the initial disrupting event (Findlay and Bourdages). The negative effects land development, human populations, and urban sprawl have on the biodiversity of wetlands is the result of wetland fragmentation. Minimum wetland densities are required to sustain healthy populations in these ecosystems. Even though these wetlands are not continuous systems, they are inter-dependent systems because their proximities to one another affect the ability of populations to genetically mix (Gibbs).

8. RECOMMENDATIONS

Based on the findings developed during this research effort, the following recommendations are proposed for future investigation of this topic:

- 1) Examine the effect of county roads upon wetlands using the procedures developed and documented with this project, and
- 2) Examine the effect of urban sprawl on wetlands, in relation to historical highway routing and construction.

9. REFERENCES

Findlay, C. S. and Bourdages, J. (2000). "*Response Time of Wetland Biodiversity to Road Construction on Adjacent Lands.*" Conservation Biology. Vol 14 Feb 2000, pp. 86-91.

Franzmeier, D. P. and Kladvko, E. J. (2000). Personal Communication, Department of Agronomy, Purdue University, West Lafayette, Indiana.

Findlay, C. S. and Bourdages, J. (2000). "*Response Time of Wetland Biodiversity to Road Construction on Adjacent Lands.*" Conservation Biology. Vol 14 Feb 2000, pp. 86-91.

Gibbs, James P. (2000). "*Wetland Loss and Biodiversity Conservation.*" Conservation Biology v 14 Feb 2000, pp. 314-317.

Harbor, J. (2001). Personal Communication. Department of Earth and Atmospheric Sciences, Purdue University, West Lafayette, Indiana.

Miller, B. K. and MacGowan, B. J. (2001). Personal Communication. Department of Forestry and Natural Resources, Purdue University Cooperative Extension Service, West Lafayette, Indiana.

NRCS at a Glance. <http://www.nrcs.usda.gov/NRCSgInc.html>

NRCS Distribution of Wetlands. <http://www.nhq.nrcs.usda.gov/land/meta/m2702.html>

NRCS SSURGO Data Users Guide (via PDF file format)
http://www.ftw.nrcs.usda.gov/ssur_data.html

NWI homepage. <http://www.nwi.fws.gov/>

NWI Overview. <http://www.nwi.fws.gov/overview.htm>

U.S. Army Corps of Engineers Wetlands Delineation Manual. Wetlands Research Program Technical Report Y-87-1. <http://www.wes.army.mil/el/wetlands/wetlands.html>

U.S. Map of SSURGO Digitized County Soil Data.
<http://www.ftw.nrcs.usda.gov/jpg/archived.jpg>

U.S. Map of Digitized NWI Wetlands Data. <http://wetlands.fws.gov/images/webstat.gif>

Santos, K. (2000). Personal Communication. Regional Wetlands Coordinator for Region 3 (Midwest_Region), U.S. Fish and Wildlife Service.