

Physics

Physics Research Publications

Purdue University

Year 2008

New open string solutions in AdS(5)

R. Ishizeki

M. Kruczenski

A. Tirziu

This paper is posted at Purdue e-Pubs.

http://docs.lib.purdue.edu/physics_articles/808

New open string solutions in AdS₅R. Ishizeki,^{*} M. Kruczenski,⁺ and A. Tirziu[‡]*Department of Physics, Purdue University, 525 Northwestern Avenue, W. Lafayette, Indiana 47907-2036, USA*

(Received 9 May 2008; published 23 June 2008)

We describe new solutions for open string moving in AdS₅ and ending in the boundary, namely, dual to Wilson loops in $\mathcal{N} = 4$ SYM theory. First we introduce an ansatz for Euclidean curves whose shape contains an arbitrary function. They are Bogomol'nyi-Prasad-Sommerfield monopoles (BPS) and the dual surfaces can be found exactly. After an inversion they become closed Wilson loops whose expectation value is $W = \exp(-\sqrt{\lambda})$. After that we consider several Wilson loops for $\mathcal{N} = 4$ SYM in a pp -wave metric and find the dual surfaces in an AdS₅ pp -wave background. Using the fact that the pp -wave is conformally flat, we apply a conformal transformation to obtain novel surfaces describing strings moving in AdS space in Poincare coordinates and dual to Wilson loops for $\mathcal{N} = 4$ SYM in flat space.

DOI: 10.1103/PhysRevD.77.126018

PACS numbers: 11.25.-w, 11.25.Tq

I. INTRODUCTION

The understanding of the AdS/CFT correspondence between $\mathcal{N} = 4$ SYM theory and the dual type IIB string theory on AdS₅ \times S⁵ has seen steady progress in recent years. Recently, the study of the correspondence between open string solutions and the dual Wilson loops received renewed attention. Following earlier discussions [1,2] on the cusp anomaly, in [3] the authors proposed that the planar four gluon scattering amplitude is described, at strong coupling, by an open string solution ending on a sequence of four lightlike lines at the boundary of AdS₅ space in T -dual coordinates. This strong coupling string calculation is in agreement with the gauge theory conjecture of [4]. A series of papers studied various aspects of this proposal [5–8].¹ The proposal put forward in [3] was further analyzed in [10]. There, a prescription of how to compute amplitudes for processes that involve local operators and gluon final states was also proposed. While the string ends on lightlike lines at the horizon in ordinary AdS₅ space, the operator insertion leads to string solutions going to the boundary of ordinary AdS₅ space, and which, at the same time are stretched in the direction in which the momentum of the operator is nontrivial.

Motivated by these recent developments in understanding various gauge theory scattering processes at strong coupling through a string description, we find in this paper several new open string solutions in AdS₅ that end at the boundary on various Wilson lines.

First we find a class of solutions ending on spacelike Wilson loops. These solutions extend to the horizon and end at the boundary of AdS₅ on a large class of open lines. The (regularized) classical area of the corresponding world-sheet is zero since these solutions are BPS. Per-

forming an inversion transformation we obtain a class of solutions that end on a closed spacelike Wilson loop at the boundary. In this case their expectation values turn out to be $\langle W \rangle = e^{-\sqrt{\lambda}}$. This result is in agreement with the conjecture in [11] that considers two Euclidean Wilson $W_{1,2}$ loops which are related by an inversion, with one of them W_1 extending to infinity and the other W_2 given by a closed contour. In that case the conjecture is that $\langle W_2 \rangle = e^{-\sqrt{\lambda}} \langle W_1 \rangle$ at the leading order in strong coupling λ . The usual example is the straight-line $\langle W_1 \rangle = 1$ and the circle $\langle W_2 \rangle = e^{-\sqrt{\lambda}}$. It would be interesting to obtain the relationship between the Wilson loops $W_{1,2}$ also at subleading orders in strong coupling.

In the second part of this paper we exploit the relation between the AdS₅ pp -wave and pure AdS₅ to find new open string solutions in the latter space. We extend the solution obtained in [12] in this context to other, more complex solutions. We find solutions ending on two straight lightlike lines in the x_+ direction on the boundary in the pp -wave AdS₅ space, which when translated in the pure AdS₅ look like hyperbolas in a (t, x) plane. Another new solution that we find is one that ends on a timelike straight line in the time direction. In pure AdS₅ coordinates it looks like a tangent shaped line in a $[x_{\pm} = \frac{1}{\sqrt{2}}(x \pm t)]$ plane. The function $z = z(x)$ for fixed t is a line with one end on the boundary, and the other at the horizon. We suggest that this type of solution can describe the scattering between a quark at the boundary moving at a speed less than the speed of light and a gluon coming out from infinity. However, it corresponds to forward scattering. It should be desirable to look for other solutions in which the incoming gluon changes direction. We find also solutions that end on two parallel lines in the time direction in the pp -wave, which are mapped into two tangent type Wilson lines on the boundary in pure AdS₅ space.

The organization of this paper is as follows: In the next section we described the new Wilson loops solutions found using an ansatz in light-cone coordinates. In Sec. III we

^{*}rishizek@purdue.edu⁺markru@purdue.edu[‡]atirziu@purdue.edu¹See also [9] and the references therein.

describe the solutions found by conformally mapping the pp -wave onto flat space. Finally we give our conclusions in the last section.

II. STRING SOLUTIONS ENDING ON OPEN/ CLOSED SPACE-LIKE WILSON LOOPS

The Nambu-Goto action for a string moving in AdS_5 in Poincaré coordinates is given by

$$\begin{aligned} S &= i \frac{\sqrt{\lambda}}{2\pi} \int d\sigma d\tau \mathcal{L} \\ &= i \frac{\sqrt{\lambda}}{2\pi} \int d\sigma d\tau \frac{1}{z^2} \sqrt{(\partial_\sigma X)^2 (\partial_\tau X)^2 - (\partial_\sigma X \cdot \partial_\tau X)^2}. \end{aligned} \quad (2.1)$$

The space-time metric is set as (here $x_\pm = \frac{1}{\sqrt{2}}(x \pm t)$)

$$ds^2 = \frac{1}{z^2} (2dx_+ dx_- + dx_1^2 + dx_2^2 + dz^2). \quad (2.2)$$

We consider the following ansatz

$$\begin{aligned} x_+ &= u(\sigma, \tau), & x_- &= 0, & x_1 &= \tau, \\ x_2 &= a, & z &= \sigma, \end{aligned} \quad (2.3)$$

where a is constant which we can set to zero (but we leave it here for later use). We also have that $z = \sigma \geq 0$, while τ varies over the real line. The surface ends at the boundary $z = \sigma = 0$ in the (spacelike) line

$$x_+ = u(\sigma = 0, \tau), \quad x_- = 0, \quad x_1 = \tau, \quad x_2 = a. \quad (2.4)$$

Our objective is to find $u(\sigma, \tau)$ given an arbitrary function $u(\sigma = 0, \tau)$.

Solving the equation of motion for x_- , we obtain,

$$u''(\sigma, \tau) - \frac{2}{\sigma} u'(\sigma, \tau) + \ddot{u}(\sigma, \tau) = 0. \quad (2.5)$$

We also find that the equations for all the other coordinates are satisfied as long as Eq. (2.5) is true. A simple check is that $u = 0$ is a solution corresponding to a straight line along x_2 . The main point of this ansatz is that the resulting Eq. (2.5) is linear and therefore can be solved using a variety of standard methods for any initial condition $u(\sigma = 0, \tau)$. Indeed, taking the Fourier transform (where we use the fact that $u(\sigma, \tau)$ is real):

$$u(\sigma, \tau) = \int_0^\infty d\omega [e^{i\omega\tau} u(\sigma, \omega) + e^{-i\omega\tau} u^*(\sigma, \omega)] \quad (2.6)$$

in (2.5) leads to

$$u''(\sigma, \omega) - \frac{2}{\sigma} u'(\sigma, \omega) - \omega^2 u(\sigma, \omega) = 0. \quad (2.7)$$

The general solution of this equation is

$$u(\sigma, \omega) = C_1 e^{\omega\sigma} (\omega\sigma - 1) + C_2 e^{-\omega\sigma} (\omega\sigma + 1). \quad (2.8)$$

Since $\omega > 0$ we keep the second solution which does not diverge as $\sigma \rightarrow \infty$. After that, the most general solution of (2.5) is

$$\begin{aligned} u(\sigma, \tau) &= \int_0^\infty d\omega e^{-\omega\sigma} (\omega\sigma + 1) \\ &\quad \times [u(0, \omega) e^{i\omega\tau} + u^*(0, \omega) e^{-i\omega\tau}] \end{aligned} \quad (2.9)$$

which is the main result of this section, allowing to compute $u(\sigma, \tau)$ given $u(0, \tau)$. A simpler expression can be obtained if we define the positive frequency part of $u(\sigma, \tau)$ as

$$\begin{aligned} u(\sigma, \tau) &= \chi_+(\sigma, \tau) + \chi_+^*(\sigma, \tau), \\ \chi_+(\sigma, \tau) &= \int_0^\infty d\omega e^{i\omega\tau} u(\sigma, \omega). \end{aligned} \quad (2.10)$$

Now we should find $\chi_+(\sigma, \tau)$ given $\chi_+(0, \tau)$. Using the previous calculations we find

$$\chi_+(\sigma, \tau) = \int_0^\infty d\omega e^{i\omega\tau - \omega\sigma} (\omega\sigma + 1) u(0, \omega) \quad (2.11)$$

$$= (1 - \partial_\xi) \int_0^\infty d\omega e^{i\omega\tau - \xi\omega\sigma} u(0, \omega) |_{\xi=1} \quad (2.12)$$

$$= \chi_+(\tau + i\sigma) - i\sigma \chi_+'(\tau + i\sigma) \quad (2.13)$$

where $\chi_+'(\tau) = \partial_\tau \chi_+(\tau)$. In fact it is straightforward to check that the function

$$\chi_+(\sigma, \tau) = \chi_+(\tau + i\sigma) - i\sigma \chi_+'(\tau + i\sigma), \quad (2.14)$$

satisfies Eq. (2.5). As an example consider a bell-shaped Wilson loop

$$x_+ = \frac{1}{\tau^2 + a^2}, \quad x_2 = \tau, \quad x_- = 0. \quad (2.15)$$

Doing the Fourier transform we obtain (assuming $a > 0$):

$$\begin{aligned} \frac{1}{\tau^2 + a^2} &= \frac{1}{2a} \int_0^\infty e^{-a\omega + i\omega\tau} d\omega + \text{c.c.}, \\ &\Rightarrow \chi_+(0, \tau) \\ &= \frac{1}{2a} \frac{1}{a - i\tau} \end{aligned} \quad (2.16)$$

which implies

$$\begin{aligned} \chi_+(\sigma, \tau) &= \chi_+(\tau + i\sigma) - i\sigma \chi_+'(\tau + i\sigma) \\ &= \frac{1}{2a} \frac{a + 2\sigma - i\tau}{(a - i\tau + \sigma)^2}. \end{aligned} \quad (2.17)$$

From here, using Eq. (2.10), we find the solution

$$u(\sigma, \tau) = \frac{(a + \sigma)^2 (a + 2\sigma) + a\tau^2}{a[(a + \sigma)^2 + \tau^2]^2}. \quad (2.18)$$

This is just an example to check the validity of the method. Given any shape we can write the solution in term of Fourier integrals using Eq. (2.9) or (2.14).

Going back to the general solution (2.3), the action S is imaginary and proportional to the area of the world sheet A :

$$S = iA \frac{\sqrt{\lambda}}{2\pi}. \quad (2.19)$$

The action is

$$-iS = \frac{\sqrt{\lambda}}{2\pi} \int d\sigma d\tau \mathcal{L} = \frac{\sqrt{\lambda}}{2\pi} \int \frac{d\sigma d\tau}{\sigma^2} = \frac{\sqrt{\lambda}}{2\pi} \frac{T}{\epsilon}, \quad (2.20)$$

where we chose the τ interval $-\frac{T}{2} \leq \tau \leq \frac{T}{2}$ and T is large. Also, we introduced a cutoff, ϵ , in z to account for the divergency near the boundary. The solution (2.3) at the boundary $z = 0$ is an open line as τ spans the real line. While the action is divergent, one should recall [13] that in fact the physical result is obtained by adding a boundary term which precisely cancels the linear divergency $\frac{1}{\epsilon}$, so the physical action is actually zero. This suggests that this solution is BPS as we now check. Note also that the result is independent of the function $u(\tau, \sigma)$.

A. Supersymmetry of the Wilson loop

The Wilson loop we considered in the previous subsection is BPS as we now proceed to prove. The arguments are well known, here we follow [14]. Given the Wilson loop

$$W = \frac{1}{N} \text{Tr} \hat{P} \exp \int d\tau (iA^\mu(x) \dot{x}_\mu + \Phi_1(x) |\dot{x}|), \quad (2.21)$$

where $x_\mu(\tau)$ parametrizes the path and $\dot{x} = \partial_\tau x$. Also, Φ_1 is a scalar field of the $\mathcal{N} = 4$ SYM theory. Supersymmetry is preserved if we can find spinor solutions ϵ to satisfy

$$(i\Gamma^\mu \dot{x}_\mu + \tilde{\Gamma}_1 |\dot{x}|) \epsilon = 0, \quad (2.22)$$

where $\Gamma^{\mu=\pm,1,2}$, $\tilde{\Gamma}^{i=1 \dots 6}$ are ten-dimensional gamma matrices. In the case of the path analyzed in the previous subsection, this boils down to

$$(i\Gamma^+ \dot{x}_+ + i\Gamma^1 + \tilde{\Gamma}^1) \epsilon = 0. \quad (2.23)$$

Since we are taking $x_+(\tau)$ as an arbitrary function we require (in special cases such as $x_+ = 0$ or $x_+ = \tau$ the Wilson loop is $\frac{1}{2}$ BPS):

$$\Gamma^+ \epsilon = 0, \quad (i\Gamma^1 + \tilde{\Gamma}^1) \epsilon = 0. \quad (2.24)$$

Using, for example, the representation of gamma matrices in terms of creation and annihilation operators (see [15]) it is easily seen that only four independent solutions exist, namely, the Wilson loop preserve one quarter of the 16 supercharges. In other words these Wilson loops are $\frac{1}{4}$ BPS. Their expectation value should be $W = 1$ independent of the coupling. Notice that in [14] it was stated that Wilson loops involving just one of the scalar fields (here Φ_1) can be BPS only if they are straight lines. However the analysis

was done in Euclidean space, in the Minkowski case this is no longer true as we have just shown.

B. Inversion and closed Wilson loop

To construct a closed loop at the boundary of AdS₅ we consider the inversion transformation $\tilde{x}_\mu = \frac{x_\mu}{|x|^2}$ which leaves the metric (2.2) invariant. After this transformation the solution (2.3) becomes

$$\tilde{x}_+ = \frac{u(\sigma, \tau)}{\tau^2 + \sigma^2 + a^2} \quad (2.25)$$

$$\begin{aligned} \tilde{x}_- &= 0, & \tilde{x}_1 &= \frac{\tau}{\tau^2 + \sigma^2 + a^2}, \\ \tilde{x}_2 &= \frac{a}{\tau^2 + \sigma^2 + a^2}, & \tilde{z} &= \frac{\sigma}{\tau^2 + \sigma^2 + a^2}. \end{aligned} \quad (2.26)$$

Starting with the Nambu-Goto action in the new coordinates with metric

$$ds^2 = \frac{1}{\tilde{z}^2} (2d\tilde{x}_+ d\tilde{x}_- + d\tilde{x}_1^2 + d\tilde{x}_2^2 + d\tilde{z}^2) \quad (2.27)$$

one can easily see that indeed (2.26) is a solution with u satisfying again (2.5). Before doing the inversion transformation, the radial coordinate z goes all the way to the horizon $0 \leq z < \infty$. After the inversion \tilde{z} has a maximum attained at $\tau = 0$, and $\sigma = a$, so $0 \leq \tilde{z} \leq \frac{1}{2a}$. Thus after the inversion the surfaces closes up in the bulk (here we assume $a \neq 0$). More precisely, the solution (2.26) projected over the subspace $x_+ = x_- = 0$ defines a sphere

$$\left(\tilde{x}_2 - \frac{1}{2a}\right)^2 + \tilde{x}_1^2 + \tilde{z}^2 = \frac{1}{4a^2} \quad (2.28)$$

which is the same as the circular solution in Euclidean space considered in [11,16,17]. Over that sphere x_+ varies according to the function $x_+ = u(\sigma, \tau)$.

At the boundary, $\tilde{z} = 0$, i.e. $\sigma = 0$, we obtain

$$\begin{aligned} \tilde{x}_+ &= \frac{u(\sigma=0, \tau)}{\tau^2 + a^2}, & \tilde{x}_- &= 0, \\ \tilde{x}_1 &= \frac{\tau}{\tau^2 + a^2} & \tilde{x}_2 &= \frac{a}{\tau^2 + a^2}. \end{aligned} \quad (2.29)$$

Projected onto the subspace $x_-, x_+ = 0$ this curve is a circle with radius, $\frac{1}{2a}$,

$$\tilde{x}_1^2 + \left(\tilde{x}_2 - \frac{1}{2a}\right)^2 = \frac{1}{4a^2}. \quad (2.30)$$

Therefore, we can parametrize the solution at the boundary as follows

$$\begin{aligned} \tilde{x}_+ &= \frac{(1 + \cos\theta)u(\theta)}{2a^2}, & \tilde{x}_1 &= \frac{1}{2a} \sin\theta, \\ \tilde{x}_2 &= \frac{1}{2a} + \frac{1}{2a} \cos\theta, \end{aligned} \quad (2.31)$$

where $u(\theta) = u(\sigma = 0, \tau)$, $\tau = \tau(\theta) = a \tan \frac{\theta}{2}$, and $0 \leq \theta < 2\pi$. As a result of the inversion transformation the Wilson lines closes up. At least as long as we take $x_+ = u(0, \tau)$ as a bounded function.

Let us compute the action for the solution (2.26)

$$-iS = \frac{\sqrt{\lambda}}{2\pi} \int d\sigma d\tau \mathcal{L} = \frac{\sqrt{\lambda}}{2\pi} \int \frac{d\sigma d\tau}{\sigma^2}. \quad (2.32)$$

As in the original coordinates, in the inverted coordinates we again introduce a cutoff near the boundary of AdS₅. This cutoff is now in \tilde{z} . This leads to the condition

$$\tilde{z} = \frac{\sigma}{\tau^2 + \sigma^2 + a^2} > \epsilon. \quad (2.33)$$

This equation is equivalent to

$$\tau^2 + \left(\sigma - \frac{1}{2\epsilon}\right)^2 < \frac{1}{4\epsilon^2} - a^2. \quad (2.34)$$

Now we can parametrize such region with

$$\tau = \rho \sin \phi \quad (2.35)$$

$$\sigma = \rho \cos \phi + \frac{1}{2\epsilon}, \quad (2.36)$$

where $0 \leq \rho < R = \sqrt{\frac{1}{4\epsilon^2} - a^2}$, and $0 \leq \phi < 2\pi$.

Therefore the action (2.32) can be computed using this parametrization as

$$-iS = \frac{\sqrt{\lambda}}{2\pi} \int \frac{\rho d\rho d\phi}{(\rho \cos \phi + \frac{1}{2\epsilon})^2} \quad (2.37)$$

$$= \frac{\sqrt{\lambda}}{2a\epsilon} - \sqrt{\lambda} + O(\epsilon^2). \quad (2.38)$$

As in the case of the open Wilson line, the linear divergent part is proportional to the length of the string in the (\tilde{y}, \tilde{w}) plane at the boundary. When the lengths in the two cases are equal, i.e. $T = \frac{\pi}{a}$, it makes sense to subtract from (2.38) the open line result so that the linear divergent term cancels. In contrast to the action (or world sheet area) of the open loop in (2.20), the action for the closed loop in (2.38) has a nontrivial finite part. This relationship is the same as the one from open/closed Wilson loops analyzed in Euclidean space in [11]. One can also compute string 1-loop corrections to these solutions following the procedure developed in [18]. Let us again observe that the result in (2.38) is independent of the function $u(\tau, \sigma)$. This Wilson loop is not BPS but should be invariant under a superconformal charge. Presumably the expectation value of this class of Wilson loops can be computed exactly by using the methods of [19] (generalized to Minkowski signature). Summing rainbow diagrams also leads to the same result as in the case of the circular Wilson loop. Finally, let us also mention that it would be interesting to

see whether there exists any relationship between the class of solutions considered here and the ones discussed in [20].

C. General procedure

In this subsection we show that the same procedure can in principle be applied to other solutions with Euclidean world-sheet metric. Indeed, suppose we have a metric of the form

$$ds^2 = 2g_{+-}(x_i, x_{\pm})dx_+dx_- + g_{ij}(x_i)dx^i dx^j \quad (2.39)$$

and a (conformal gauge) solution $x_i = \bar{x}_i(\sigma, \tau)$ with $x_{\pm} = 0$. Now let us look for a new solution with the ansatz $x_+ = u(\sigma, \tau)$, $x_- = 0$, and $x_i = \bar{x}_i(\sigma, \tau)$. The constraints (recalling that we are looking at Euclidean solutions)

$$g_{\mu\nu}(\dot{x}^\mu \dot{x}^\nu - x'^\mu x'^\nu) = 0, \quad g_{\mu\nu} \dot{x}^\mu x'^\nu = 0 \quad (2.40)$$

are automatically satisfied (if they were satisfied for the original solution). The equations of motion

$$\partial_\tau(g_{\mu\nu} \dot{x}^\nu) + \partial_\sigma(g_{\mu\nu} x'^\nu) = \partial_\mu g_{\alpha\beta} (\dot{x}^\alpha \dot{x}^\beta + x'^\alpha x'^\beta) \quad (2.41)$$

are satisfied when $\mu = i$, and, since $\partial_\pm g_{ij} = 0$, also for $\mu = +$. The equation with $\mu = -$ gives

$$\partial_\tau(g_{-+} \dot{x}^+) + \partial_\sigma(g_{-+} x'^+) = 0. \quad (2.42)$$

After replacing for the solution $x_i = \bar{x}_i(\sigma, \tau)$, this equation of motion is linear in x_+ if $\partial_+ g_{+-} = 0$ as was the case in our previous discussion. If not, the equation is in general nonlinear. In both cases this method can provide new solutions given old Euclidean ones. Here we looked at the straight string and the circular Wilson loop. It might be interesting to analyze other cases as well.

III. OPEN STRING SOLUTIONS OBTAINED THROUGH THE MAP TO pp -WAVE BACKGROUND

One can find new open string solutions in AdS₅ by exploiting the map between pure AdS₅ and the AdS₅ pp -wave backgrounds. These solutions correspond to Wilson loops in Minkowski space in the dual gauge theory. The relationship between the AdS₅ metric (here $x_{\pm} = \frac{x \pm t}{\sqrt{2}}$, $i = 1, 2$, and we use the notation $x_+ \equiv x^+$, $x_- \equiv x^-$)

$$ds^2 = \frac{1}{\tilde{z}^2} (2d\tilde{x}_+ d\tilde{x}_- + d\tilde{x}_i d\tilde{x}_i + d\tilde{z} d\tilde{z}) \quad (3.1)$$

and pp -wave AdS₅ metric

$$ds^2 = \frac{1}{z^2} \left(2dx_+ dx_- - \frac{1}{2}(x_i^2 + z^2)F(x_+) dx_+^2 + dx_i^2 + dz^2 \right) \quad (3.2)$$

is given by [21]

$$\begin{aligned}\tilde{x}_+ &= f(x_+), & \tilde{x}_- &= x_- - \frac{1}{4}(x_i^2 + z^2)\frac{f''}{f'}, \\ \tilde{x}_i &= x_i\sqrt{f'}, & \tilde{z} &= z\sqrt{f'},\end{aligned}\quad (3.3)$$

where F is given by the Schwarzian derivative

$$F(x_+) = \{f, x_+\} = \frac{f'''}{f'} - \frac{3}{2}\left(\frac{f''}{f'}\right)^2. \quad (3.4)$$

Mapping back simple solutions in the pp -wave AdS₅ background usually produce new and potentially useful solutions in pure AdS₅.

Although in principle one can consider various functions $F(x_+)$ to obtain new Wilson loops, in this paper we just consider the simplest nontrivial case corresponding to $F(x_+) = a$ where a is a constant. In that case, the solution to (3.4) has the form (up to shifts² in x_+)

$$f(x_+) = A\sqrt{\frac{2}{a}}\tan\sqrt{\frac{a}{2}}x_+, \quad (3.5)$$

where A is a constant. Notice that in the limit $a \rightarrow 0$, AdS₅ is mapped onto itself through

$$f(x_+) = \frac{A}{x_+}. \quad (3.6)$$

The transformation of coordinates in this case is given by

$$\begin{aligned}\tilde{x}_+ &= \frac{A}{x_+}, & \tilde{x}_- &= x_- + \frac{1}{2}\frac{x_i^2 + z^2}{x_+}, \\ \tilde{x}_i &= x_i\frac{\sqrt{-A}}{x_+}, & \tilde{z} &= z\frac{\sqrt{-A}}{x_+},\end{aligned}\quad (3.7)$$

where one needs to choose $A < 0$.

In what follows we will be concerned with the situation when $F(x_+)$ is constant but nonzero. As in [12] let us choose $f(x_+)$ of the form $f(x_+) = \frac{1}{\mu}\tan\mu x_+$. Therefore, the AdS₅ pp -wave metric and the transformation are

$$ds^2 = \frac{1}{z^2}(2dx_+dx_- - \mu^2(x_i^2 + z^2)dx_+^2 + dx_i^2 + dz^2) \quad (3.8)$$

and

$$\begin{aligned}\tilde{x}_+ &= \frac{1}{\mu}\tan\mu x_+, & \tilde{x}_- &= x_- - \frac{1}{2}\mu(x_i^2 + z^2)\tan\mu x_+, \\ \tilde{x}_i &= \frac{x_i}{\cos\mu x_+}, & \tilde{z} &= \frac{z}{\cos\mu x_+}.\end{aligned}\quad (3.9)$$

The parameter μ can be scaled away by a rescaling $x_+ \rightarrow \frac{x_+}{\mu}$, $x_- \rightarrow \mu x_-$, so it can be set to 1. However, below we will keep μ explicitly in order to be able to make the connection to pure AdS₅ which corresponds to the limit

²Note that a shift in x_+ is an isometry if $F(x_+)$ is constant.

$\mu \rightarrow 0$. Using the map (3.9) we find below new open string solutions. Let us finish this section by pointing out that the map (3.9) is singular at $\mu x_+ = (2k+1)\frac{\pi}{2}$, $k \in \mathbb{Z}$. This means that this map takes part of the pp -wave AdS₅ space into the full AdS₅ space.

A. Straight lines along the x_+ direction

1. 1-line

We start by reviewing the solution in the AdS₅ pp -wave that ends on a single lightlike solution, and was found in [12]

$$x_+ = \tau, \quad x_- = 0, \quad x_i = 0, \quad z = \sigma. \quad (3.10)$$

On the boundary $\sigma = 0$, this solution may be interpreted as the world line of a massless quark at $x_- = 0$. Let us point out that this is also a solution in pure AdS₅ whose world sheet is just a half-plane ending on the x_+ line at the boundary. One may wonder whether there exists a direct extension of this solution to a timelike line with $x_i \neq 0$ but constant, as for example $x_i = b$. It turns out that there are no such solutions, i.e. $b = 0$. The charges in the AdS₅ pp -wave background have been computed in [12], where it was found that their divergent parts are related by

$$P_+ \approx \frac{\lambda}{4\pi} \ln|P_-|. \quad (3.11)$$

Let us use the transformation (3.9) and write the solution (3.10) in pure AdS₅

$$\begin{aligned}\tilde{x}_+ &= \frac{1}{\mu}\tan\mu\tau, & \tilde{x}_- &= -\frac{\mu}{2}\sigma^2\tan\mu\tau, \\ \tilde{z} &= \frac{\sigma}{\cos\mu\tau}, & \tilde{x}_i &= 0.\end{aligned}\quad (3.12)$$

The world-sheet surface is given by

$$\tilde{z} = \sqrt{-\frac{2\tilde{x}_-}{\mu^2\tilde{x}_+}(1 + \mu^2\tilde{x}_+^2)}. \quad (3.13)$$

On the boundary this solution again ends on a lightlike line but now in contrast with (3.10) the string world sheet is more complicated. For convenience let us parametrize the surface as

$$\tilde{z}(\tilde{x}, \tilde{t}) = \sqrt{-\frac{2}{\mu^2}\frac{\tilde{x} - \tilde{t}}{\tilde{x} + \tilde{t}}(1 + \frac{\mu^2}{2}(\tilde{x} + \tilde{t})^2)}. \quad (3.14)$$

The plot of \tilde{z} as a function of \tilde{x} for fixed $\tilde{t} < 0$ is shown in Fig. 1, while for $\tilde{t} > 0$ in Fig. 2. For $\tilde{t} < 0$ the rounded spike is to the left of the vertical straight line and it moves to the left as \tilde{t} increases. At $\tilde{t} = 0$ the curve is a straight vertical line. As \tilde{t} becomes positive the spike moves further to the left being now to the right of the vertical line. The position of the maximum of the semicircular part of the solution depends on \tilde{t} and it is at zero in the $|\tilde{t}| \rightarrow \infty$ limit.




FIG. 1 (color online). Plot of \tilde{z} versus \tilde{x} for $\mu = 1$, and $\tilde{t} = -200, -100, -50, -10$.

As already pointed out, the position of the rounded spike is close to $-\tilde{t}$ for $\tilde{t} < 0$ but a bit to the left of the vertical straight line (see Fig. 1). This shift depends on \tilde{t} and in the large \tilde{t} limit we obtain $\tilde{x}_s = -\tilde{t} - \delta_1$, with $\delta_1 = \frac{\sqrt{2}}{\mu}$. For $\tilde{t} > 0$ the position of the spike is a little bit to the right of the vertical line (Fig. 2). In the limit $\tilde{t} \rightarrow \infty$, we now have $\tilde{x}_s = -\tilde{t} + \delta_1$. This means that there is a time delay in this process when moving from $-\infty$ to ∞ in \tilde{t} . In the limit $|\tilde{t}| \rightarrow \infty$ the height of the spike goes to the horizon as $z_s \sim 2^{5/4} \sqrt{\frac{|\tilde{t}|}{\mu}}$. The plots in Fig. 1 and 2, suggest that we can think of this solution as corresponding to a scattering process in the $1+1$ dimensional world-sheet with \tilde{t}, \tilde{x} being the $2d$ time and space coordinates. Because of the symmetry of the solution under $(\tilde{x}, \tilde{t}) \rightarrow (-\tilde{x}, -\tilde{t})$ this is an elastic process, which is characteristic to integrable models in $1+1$ dimensions.

Regarding the space-time/gauge theory interpretation of this solution, it was suggested in [12] that one can identify the end of the string attached to the boundary as a quark. In addition there is the spike coming out from the horizon,




FIG. 2 (color online). Plot of \tilde{z} versus \tilde{x} for $\mu = 1$, and $\tilde{t} = 200, 100, 50, 10$.

which may be identified with a gluon. Then what this solution appears to describe is a quark-gluon scattering process, but it is not clear how to make this suggestion more precise. It would be easier if one could find solutions describing scattering in which the gluon changes direction.³

For completeness it is interesting to compute the energy and momentum of the solution. Since one is forcing the quark to move along a particular trajectory, they are not conserved. In fact, the result determines the force necessary to push the quark. For $\tilde{t} > 0$ the energy is $E = \frac{\sqrt{\Lambda}}{2\pi} \mathcal{E}$

$$\begin{aligned} \mathcal{E}(\tilde{t}) &= \int_{-\tilde{t}}^{\tilde{t}} d\tilde{x} \frac{\partial L}{\partial \tilde{t}} = \int_{-\tilde{t}+\epsilon_0}^{\tilde{t}-\epsilon_0} d\tilde{x} \frac{1 + \tilde{z}'^2}{\tilde{z}^2 \sqrt{1 - \tilde{z}^2 + \tilde{z}'^2}} \\ &= \frac{\mu}{4\sqrt{2}} \left[-1 + \frac{4}{2 + \mu^2 \epsilon_0^2} + \frac{4}{2 + \mu^2 (\epsilon_0 - 2\tilde{t})^2} \right. \\ &\quad \left. + \frac{\epsilon_0}{\epsilon_0 - 2\tilde{t}} + \frac{2\tilde{t}}{\epsilon_0} + 2 \log \frac{2\tilde{t} - \epsilon_0}{\epsilon_0} \right] \\ &= \frac{\mu \tilde{t}}{2\sqrt{2} \epsilon_0} + \frac{\mu}{4\sqrt{2}} \left[-3 + \frac{2}{1 + 2\mu^2 \tilde{t}^2} - \log \frac{\epsilon_0^2}{4\tilde{t}^2} \right], \end{aligned} \quad (3.15)$$

where dot represents derivative in respect to \tilde{t} , and prime is derivative in respect to \tilde{x} . We have introduced a small cutoff ϵ_0 , and expanded in the last line in small ϵ_0 . The cutoff ϵ_0 can be related to the physical cutoff ϵ in \tilde{z} by $\epsilon = \sqrt{2\tilde{t} + \frac{1}{\mu^2 \tilde{t}}} \epsilon_0$. For $\tilde{t} < 0$ one obtains a similar expression

$$\begin{aligned} \mathcal{E}(\tilde{t}) &= -\frac{\mu \tilde{t}}{2\sqrt{2} \epsilon_0} + \frac{\mu}{4\sqrt{2}} \left[-3 + \frac{2}{1 + 2\mu^2 \tilde{t}^2} - \log \frac{\epsilon_0^2}{4\tilde{t}^2} \right] \\ &\quad + O(\epsilon_0). \end{aligned} \quad (3.16)$$

Let us compute the momentum for $\tilde{t} > 0$, $P = \frac{\sqrt{\Lambda}}{2\pi} \mathcal{P}$

$$\begin{aligned} \mathcal{P}(\tilde{t}) &= \int_{-\tilde{t}}^{\tilde{t}} d\tilde{x} \frac{\partial L}{\partial \tilde{x}} = \int_{-\tilde{t}+\epsilon_0}^{\tilde{t}-\epsilon_0} d\tilde{x} \frac{\tilde{z} \tilde{z}'}{\tilde{z}^2 \sqrt{1 - \tilde{z}^2 + \tilde{z}'^2}} \\ &= -\frac{\mu \tilde{t}}{2\sqrt{2} \epsilon_0} + \frac{\mu}{4\sqrt{2}} \left[-1 + \frac{2}{1 + 2\mu^2 \tilde{t}^2} - \log \frac{\epsilon_0^2}{4\tilde{t}^2} \right] \\ &\quad + O(\epsilon_0). \end{aligned} \quad (3.17)$$

For $\tilde{t} < 0$ we obtain

$$\begin{aligned} \mathcal{P}(\tilde{t}) &= \frac{\mu \tilde{t}}{2\sqrt{2} \epsilon_0} + \frac{\mu}{4\sqrt{2}} \left[-1 + \frac{2}{1 + 2\mu^2 \tilde{t}^2} - \log \frac{\epsilon_0^2}{4\tilde{t}^2} \right] \\ &\quad + O(\epsilon_0). \end{aligned} \quad (3.18)$$

At $\tilde{t} = \pm \infty$ the energy of the string is the same, while at intermediate times the energy increases. The same happens with the momentum of the string. Thus when the gluon and

³In that respect, see [22] for some examples of how to construct solutions involving spikes.

quark get closer together the energy of the system increases. While both the energy and momentum of string depend on time \tilde{t} , their difference does not (here we ignore the linear divergent terms)

$$E = P - \frac{\sqrt{\lambda}}{4\sqrt{2}\pi} \mu. \quad (3.19)$$

As we already pointed out, there is a time delay when the rounded spike moves from $\tilde{t} \rightarrow -\infty$ to $\tilde{t} \rightarrow \infty$ given by $\Delta\tilde{t} = 2\delta_1 = \frac{2\sqrt{2}}{\mu}$. In 1 + 1 dimensional theories this time delay translates into a phase-shift which semiclassically is related to the time delay as [23]

$$\frac{d\delta(E)}{dE} = \frac{1}{2} \Delta\tilde{t}(E). \quad (3.20)$$

To compute this phase shift we need first to get the energy of the incoming gluon. We do not know how to define the energy of the incoming gluon but, since the only scale of the problem is μ we can assume that $E \sim \mu$. This gives then a phase shift $\delta \sim \ln E$ for this scattering.

2. 2-parallel lines

Let us look now for an open string solution which at the boundary ends on two timelike parallel lines extended in the x_+ direction as (we assume $b_1 < b_2$)

$$x_+ = \tau, \quad x_- = 0, \quad x_1 = b_1, \quad x_2 = 0 \quad (3.21)$$

$$x_+ = \tau, \quad x_- = 0, \quad x_1 = b_2, \quad x_2 = 0. \quad (3.22)$$

The full ansatz for such a solution is

$$\begin{aligned} x_+ = \tau, \quad x_- = 0, \quad x_1 = \sigma, \\ z = z(\sigma), \quad x_2 = 0. \end{aligned} \quad (3.23)$$

We want to have the condition at the boundary $z(b_1) = z(b_2) = 0$. The Nambu-Goto Lagrangian is explicitly dependent on σ

$$S = \frac{\sqrt{\lambda}}{2\pi} \int d\tau d\sigma \frac{\mu}{z^2} \sqrt{(\sigma^2 + z^2)(1 + z'^2)}. \quad (3.24)$$

The equation of motion for z is

$$z''z(z^2 + \sigma^2) + (1 + z'^2)(z^2 + \sigma z z' + 2\sigma^2) = 0. \quad (3.25)$$

To solve this differential equation let us first consider the change in function $z = \sigma f(\sigma)$, and then a change in variable $u = \ln \sigma$. Then the differential equation for f is

$$\begin{aligned} (f' + f'')f(f^2 + 1) \\ + [1 + (f + f')^2][f^2 + f(f + f') + 2] = 0, \end{aligned} \quad (3.26)$$

where here derivatives are with respect to the variable u . With the above change of variables we restricted σ to positive values, but to account also for possible solutions extended to negative σ we consider gluing different branches of solutions. Note that since $z \geq 0$, the physical

solutions are those with $f \geq 0$. To proceed we consider the differential equation for $u(f)$ instead of $f(u)$ as above

$$\begin{aligned} \ddot{u} = \frac{1}{f(f^2 + 1)} [f + 2(1 + 2f^2)\dot{u} + 6f(1 + f^2)\dot{u}^2 \\ + 2(1 + f^2)^2\dot{u}^3]. \end{aligned} \quad (3.27)$$

Setting $y = \dot{u}$, we integrate the equation for $y(f)$ and obtain the solution

$$y(f) = -\frac{f}{1 + f^2} \pm \frac{f^2}{(1 + f^2)\sqrt{1 + 2f^2 + c(1 + f^2)^2}}, \quad (3.28)$$

where c is an integration constant. For $c \geq 0$ the expression under the square root is always positive. For $c < 0$ we need $c > -1$, and f in the interval $0 \leq f \leq \sqrt{-\frac{1+c+\sqrt{1+c}}{c}}$ in order for the square root to be real.

The expression in (3.28) can be further integrated and we obtain a solution for $u = u(f)$ in terms of Elliptic integrals. The boundary condition for the last integration is $u(0) = \ln b_2$ with $b_2 > 0$. After the integration the result can be expressed as a transcendental equation for $z = z(\sigma)$ as $\sigma = e^{u(z/\sigma)}$. However, the other boundary condition that we want is that the other end of the string ends also on the boundary. Therefore we want to have $z = f e^{u(f)} \rightarrow 0$ for some f . It turns out that this is only possible when the constant of integration $c = 0$ and only for the minus sign solution in (3.28). Below we see that we can have solutions for arbitrary $c > 0$, which are extended to the negative σ region. For $c < 0$ there are no solutions possible even if one tries to glue solutions because the two branches in (3.28) behave differently.


Let us analyze first the solution with $c = 0$. In this case in the limit $f \rightarrow \infty$, indeed we have $z \rightarrow 0$, and this happens for $\sigma = b_1 = 0$. For $c = 0$ the transcendental equation for $f = f(\sigma)$ simplifies to

$$\begin{aligned} \sigma = \frac{b_2}{\sqrt{1 + f^2}} \left(\frac{\sqrt{1 + 2f^2} - f}{\sqrt{1 + 2f^2} + f} \right)^{-(1/2)} (\sqrt{2}f \\ + \sqrt{1 + 2f^2})^{-(1/\sqrt{2})}. \end{aligned} \quad (3.29)$$

We plot numerically the solution $z = z(\sigma)$ in Fig. 3. This solution is independent of μ . The constant b_2 can be expressed as

$$b_2 = \int_0^{b_2} d\sigma = - \int_0^\infty df \sigma', \quad (3.30)$$

where $\sigma' = \frac{d\sigma}{df}$ and one needs to use (3.29) and integrate this expression. The integral cannot be done exactly, but numerically one can show that relation (3.30) is satisfied. Note that b_2 drops out from (3.30) therefore b_2 is not fixed. The value of f when z is maximum can be obtained from


 FIG. 3 (color online). Plot of z versus σ for $b_2 = 1$.

$$\frac{dz}{d\sigma} = -\frac{1}{f} - f + \sqrt{1+2f^2} = 0 \Rightarrow f_0 = \sqrt{\frac{1+\sqrt{5}}{2}} \quad (3.31)$$

so that the position of the maximum in σ is

$$\begin{aligned} \sigma_0 &= b_2 \sqrt{\frac{1+\sqrt{5}}{2}} [\sqrt{1+\sqrt{5}} + \sqrt{2+\sqrt{5}}]^{-(1/\sqrt{2})} \\ &\approx 0.4897b_2. \end{aligned} \quad (3.32)$$

Notice that the position of the maximum z is not at half of the σ interval.

This solution ends on the boundary at $\sigma = 0$ on a lightlike line and at $\sigma = b_2$ on a timelike curve. Let us see how these lines look in the pure AdS₅ metric. The line at $\sigma = b_2$ maps into

$$\begin{aligned} \tilde{x}_+ &= \frac{1}{\mu} \tan \mu \tau, & \tilde{x}_- &= -\frac{1}{2} \mu b_2^2 \tan \mu \tau, \\ \tilde{x}_1 &= \frac{b_2}{\cos \mu \tau}, & \tilde{x}_2 &= 0, \end{aligned} \quad (3.33)$$

while the line at $\sigma = 0$ maps into

$$\tilde{x}_+ = \frac{1}{\mu} \tan \mu \tau, \quad \tilde{x}_- = 0, \quad \tilde{x}_1 = 0, \quad \tilde{x}_2 = 0. \quad (3.34)$$

The last null line is rather simple. The former line may be expressed as

$$\tilde{x}_1 = b_2 \sqrt{1 + \mu^2 \tilde{x}_+^2}, \quad \tilde{x}_- = -\frac{1}{2} b_2^2 \mu^2 \tilde{x}_+, \quad (3.35)$$

While in the pp -wave AdS₅, in the (x_1, x_+) plane the Wilson loop were straight parallel lines, in the pure AdS₅ one remains straight line but the other is a hyperbola in the $(\tilde{x}_1, \tilde{x}_+)$ plane. As these Wilson lines might correspond to physical particles moving on the boundary let us consider plotting \tilde{t} versus \tilde{x}_1 . Let us choose $b_2^2 = 2$ in (3.33). Then after a boost in the \tilde{x}_+, \tilde{x}_- directions one can get rid of the factors of μ in front, so the hyperbola line is

$$\begin{aligned} \tilde{x}_+ &= \tan \mu \tau, & \tilde{x}_- &= -\tan \mu \tau, \\ \tilde{x}_1 &= \frac{\sqrt{2}}{\cos \mu \tau}, & \tilde{x}_2 &= 0. \end{aligned} \quad (3.36)$$

Therefore one obtains

$$\tilde{t} = \sqrt{2} \tan \mu \tau, \quad \tilde{x} = 0, \quad \tilde{x}_1 = \sqrt{2(1 + \tilde{t}^2)}, \quad (3.37)$$

while for the lightlike line

$$\tilde{t} = \tilde{x} = \frac{1}{\sqrt{2}} \tan \mu \tau, \quad \tilde{x}_1 = 0. \quad (3.38)$$


The hyperbola $\tilde{t} = \pm \sqrt{\frac{\tilde{x}_1^2}{2} - 1}$ corresponding to the timelike line (3.37) is represented in Fig. 4.

The conserved charges in the pp -wave AdS₅ background can be expressed as

$$\begin{aligned} P_+ &= \frac{\sqrt{\lambda}}{2\pi} \mu \int_0^{b_2} d\sigma \frac{\sqrt{(\sigma^2 + z^2)(1 + z'^2)}}{z^2} \\ &= -\frac{\sqrt{\lambda}}{2\pi} \mu \int_0^\infty df \frac{\sigma'}{\sigma f^2} \sqrt{(1 + f^2) \left(1 + \left(f + \frac{\sigma}{\sigma'}\right)^2\right)} \end{aligned} \quad (3.39)$$

$$\begin{aligned} P_- &= -\frac{\sqrt{\lambda}}{2\pi\mu} \int_0^{b_2} d\sigma \frac{1}{z^2} \sqrt{1 + z'^2} \\ &= \frac{\sqrt{\lambda}}{2\pi\mu} \int_0^\infty df \frac{\sigma'}{\sigma^3 f^2} \sqrt{\frac{1 + \left(f + \frac{\sigma}{\sigma'}\right)^2}{1 + f^2}}. \end{aligned} \quad (3.40)$$

Near the boundary of AdS₅ the integrals diverge, therefore we introduce a cutoff ϵ in z . While we cannot perform the above integrals exactly we can extract the divergent parts. Cutoff in z translates into cutoffs in f . Recall that $z = 0$ both for $f = 0$ and for $f \rightarrow \infty$, thus there is a ϵ dependent large small cutoff in f , f_{\min} as well as a large one f_{\max} . Using (3.29) and let us expand $z = \sigma f$ in the limits


 FIG. 4 (color online). Plot of \tilde{t} versus \tilde{x}_1 for $b_2 = \sqrt{2}$ at the boundary $z = 0$.

$$z = \sigma f = b_2 f + O(f^2) = \epsilon \Rightarrow f_{\min} = \frac{\epsilon}{b_2} \quad (3.41)$$

$$\begin{aligned} z = \sigma f &= f^{-(1/\sqrt{2})} \left[b_2 (1 + \sqrt{2}) 2^{-(3/2\sqrt{2})} + O\left(\frac{1}{f^2}\right) \right] \\ &= \epsilon \Rightarrow f_{\max} = \left(\frac{b_2}{\epsilon}\right)^{\sqrt{2}} \frac{(1 + \sqrt{2})^{\sqrt{2}}}{2\sqrt{2}}. \end{aligned} \quad (3.42)$$

To compute the charges (3.39) and (3.40) we expand the integrands in the small and large f limits and extract only the values of the integral near f_{\min} and f_{\max} . More precisely (3.39) becomes

$$\begin{aligned} P_+ &= -\frac{\sqrt{\lambda}\mu}{2\pi} \left[\int_{f_{\min}} df \left(-\frac{1}{f^2}\right) + \int^{f_{\max}} \left(-\frac{1}{\sqrt{2}f}\right) \right] \\ &= \frac{\sqrt{\lambda}\mu}{2\pi} \left(\frac{b_2}{\epsilon} + \log \frac{b_2}{\epsilon} + O(\epsilon^0) \right). \end{aligned} \quad (3.43)$$

For the other charge (3.40) we obtain

$$\begin{aligned} P_- &= \frac{\sqrt{\lambda}}{2\pi\mu} \left[\int_{f_{\min}} df \left(-\frac{1}{b_2^2 f^2}\right) \right. \\ &\quad \left. + \int^{f_{\max}} df \frac{(2\sqrt{2} - 3) 2^{(3/\sqrt{2}) - (1/2)}}{b_2^2} f^{\sqrt{2}-1} \right] \\ &= -\frac{\sqrt{\lambda}}{2\pi\mu} \left[\frac{1}{2\epsilon^2} + \frac{1}{b_2 \epsilon} + O(\epsilon^0) \right]. \end{aligned} \quad (3.44)$$

Keeping the leading orders in ϵ in these charges, we eliminate ϵ and obtain the relationship between the diver-

gent parts of the charges

$$P_- \approx -\frac{\pi}{\sqrt{\lambda}\mu^3 b_2^2} P_+^2. \quad (3.45)$$

Let us finish the discussion about the charges by pointing out that the action (3.24) on the solution is real, and up to the length of the τ interval it is the same as P_+ .

Let us return to the differential equation (3.28) and consider the situation $c > 0$. The two branches in Eq. (3.28) behave in the same way. They both satisfy $\sigma \rightarrow 0$ as $f \rightarrow \infty$, but there is no point other than $\sigma = b$ for which $z = 0$. We take the negative branch, denoted z_- , to depend on c and $b_2 > 0$, while the positive one, z_+ on c and $b_1 > 0$. For fixed b_1 and b_2 we want to find a solution for $c = c(b_1, b_2)$ so that

$$z_+ - z_-|_{f \rightarrow \infty} = 0, \quad \frac{dz_+}{d\sigma} + \frac{dz_-}{d\sigma} \Big|_{f \rightarrow \infty} = 0. \quad (3.46)$$

For such solutions we use the positive branch for $\sigma < 0$, and therefore this branch smoothly continues to the negative branch to negative values of σ until it reaches the boundary at $\sigma = -b_1$. Let us find the constant c satisfying (3.46). The solutions of (3.28) for $c > 0$ is (recall that $\sigma = e^u$)

$$u_{\pm}(f) = \mp G(f, c) + \ln \frac{b}{\sqrt{1+f^2}}, \quad (3.47)$$

where

$$\begin{aligned} G(f, c) &= \frac{i}{\sqrt{1+c} - \sqrt{1+c}} \left(F \left[i \sinh^{-1} \left[f \sqrt{1 - \frac{1}{\sqrt{1+c}}} \right], \frac{2+c+2\sqrt{1+c}}{c} \right] \right. \\ &\quad \left. - \Pi \left[\frac{1+c+\sqrt{1+c}}{c}, i \sinh^{-1} \left[f \sqrt{1 - \frac{1}{\sqrt{1+c}}} \right], \frac{2+c+2\sqrt{1+c}}{c} \right] \right), \end{aligned} \quad (3.48)$$

where $F[x, m]$ and $\Pi[n, x, m]$ are the incomplete elliptic functions defined as

$$\begin{aligned} F[x, m] &= \int_0^x \frac{d\theta}{\sqrt{1 - m \sin^2 \theta}}, \\ \Pi[n, x, m] &= \int_0^x \frac{d\theta}{(1 - n \sin^2 \theta) \sqrt{1 - m \sin^2 \theta}}. \end{aligned} \quad (3.49)$$

The conditions (3.46) become

$$\begin{aligned} \frac{1}{2} \ln \frac{b_1}{b_2} &= G(f, c)|_{f \rightarrow \infty} \\ &= \sqrt{\frac{1+c+\sqrt{1+c}}{c}} \left[-K \left[-\frac{2(1+\sqrt{1+c})}{c} \right] \right. \\ &\quad \left. + \Pi \left[\frac{1}{1-\sqrt{1+c}}, -\frac{2(1+\sqrt{1+c})}{c} \right] \right] \end{aligned} \quad (3.50)$$

and

$$2f + \frac{1}{u'_+(f)} + \frac{1}{u'_-(f)} \Big|_{f \rightarrow \infty} = -\frac{1+c}{c} \frac{2}{f} + O\left(\frac{1}{f^2}\right) = 0, \quad (3.51)$$

where $K[x] = F[\frac{\pi}{2}, x]$ and $\Pi[x, y] = \Pi[x, \frac{\pi}{2}, y]$ are elliptic integrals. The condition (3.51) is satisfied in the limit $f \rightarrow \infty$. The right-hand side in (3.50) is negative and satisfies $\lim_{c \rightarrow 0} G(\infty, c) = -\infty$, and $\lim_{c \rightarrow \infty} G(\infty, c) = 0$. The fact that $G(\infty, c)$ is negative is in accord with our initial assumption $b_1 < b_2$. Also, the limit $b_1 \rightarrow 0$ gives $c \rightarrow 0$ in accord to the previous $c = 0$ analysis. For a fixed ratio $\frac{b_1}{b_2} < 1$, Eq. (3.50) gives a solution for $c = c(b_1, b_2)$. While it is not possible to invert (3.50) to get $c = c(b_1, b_2)$ one can solve it numerically. For example for $b_2 = 1$, $b_1 = 0.5$ we obtain $c \simeq 4.77$. Gluing together the two solutions

as explained above we obtain the solution shown in Fig. 5. On the boundary ($z = 0$) this solution ends on two timelike lines at $x_1 = b_2$ and $x_1 = -b_1$. They can be mapped to pure AdS₅ coordinates as in (3.33) where the lines look like hyperbolas.

Let us compute the charges (3.39) and (3.40) in this case with $c > 0$. Again the integrals cannot be computed exactly but we can extract the divergent terms coming from the region near the boundary $z = 0$. We need to expand the two branches given in (3.47) near $f = 0$ using the negative branch near $\sigma = b_2$ and the positive branch near $\sigma = b_1$. This leads to two cutoffs in f near the boundary of AdS₅

$$z_+ = f\sigma = b_1 f + O(f^2) = \epsilon \Rightarrow f_1 = \frac{\epsilon}{b_1}, \quad (3.52)$$

and

$$z_- = f\sigma = b_2 f + O(f^2) = \epsilon \Rightarrow f_2 = \frac{\epsilon}{b_2}. \quad (3.53)$$

The charge (3.39) can be written as

$$P_+ = -\frac{\sqrt{\lambda}}{2\pi} \mu \left[\int_{f_1}^{\infty} df \frac{\sigma'_+}{\sigma_+ f^2} \sqrt{(1+f^2) \left(1 + \left(f + \frac{\sigma_+}{\sigma'_+}\right)^2\right)} + \int_{f_2}^{\infty} df \frac{\sigma'_-}{\sigma_- f^2} \sqrt{(1+f^2) \left(1 + \left(f + \frac{\sigma_-}{\sigma'_-}\right)^2\right)} \right]. \quad (3.54)$$

Expanding the integrands near $f = 0$ we obtain

$$P_+ = -\frac{\sqrt{\lambda}}{2\pi} \mu \left[\int_{f_1} df \left(-\frac{1}{f^2}\right) + \int_{f_2} df \left(-\frac{1}{f^2}\right) \right] = \frac{\sqrt{\lambda}}{2\pi} \mu \frac{b_1 + b_2}{\epsilon} + O(\epsilon^0). \quad (3.55)$$

The other conserved charge (3.40) can be computed in a similar way and we obtain




FIG. 5 (color online). Plot of z versus σ for $b_2 = 1$, $b_1 = 0.5$, $c = 4.77$.

$$P_- = -\frac{\sqrt{\lambda}}{2\pi\mu} \frac{b_1 + b_2}{b_1 b_2 \epsilon} + O(\epsilon^0) \quad (3.56)$$

Let us observe that the divergent part of the charges do not depend explicitly on c . Eliminating ϵ we obtain the following relationship between the divergent parts of the charges

$$\frac{P_+}{P_-} \approx -\mu^2 b_1 b_2. \quad (3.57)$$

B. Straight lines along the time direction t

1. 1-line

Let us start with the metric (3.8) restoring the time coordinate in Poincare patch ($t = x_0$)

$$ds^2 = \frac{1}{z^2} \left(-dt^2 + dx^2 - \frac{\mu^2}{2} (x_i^2 + z^2) (dx + dt)^2 + dx_i^2 + dz^2 \right). \quad (3.58)$$

We look for a solution which at the boundary ends on a timelike line along the t direction. A solution of this form is

$$t = \tau, \quad x = 0, \quad z = \sigma, \quad x_i = 0. \quad (3.59)$$

Here $\sigma > 0$. In the pp -wave AdS₅ space we compute the conserved charges and obtain

$$E = \frac{\sqrt{\lambda}}{2\pi} \left[\frac{1}{\epsilon} + \frac{\mu}{\sqrt{2}} \log R + O(R^0, \epsilon) \right], \quad (3.60)$$

where we have introduced a cutoff ϵ near the boundary and a cutoff R near the horizon. Here cutoffs in z are the same as cutoffs in σ . The other conserved charge corresponding to the isometry along x is

$$P = \frac{\sqrt{\lambda}}{2\pi} \left[\frac{\mu}{\sqrt{2}} \log R + O(R^0) \right]. \quad (3.61)$$

Computing the action on this solution we obtain the same expression as the energy up to a factor of T , where we take $-\frac{T}{2} \leq \tau \leq \frac{T}{2}$ with T large

$$S = \frac{\sqrt{\lambda}}{2\pi} T \int_{\epsilon}^R d\sigma \frac{1}{\sigma^2} \sqrt{1 + \frac{\mu^2}{2} \sigma^2} = \frac{\sqrt{\lambda}}{2\pi} T \left[\frac{1}{\epsilon} + \frac{\mu}{\sqrt{2}} \log R + O(R^0, \epsilon) \right]. \quad (3.62)$$

We observe that the only divergency in ϵ is linear; like in the previous cases this divergency should be canceled by a boundary term. In the limit $\mu \rightarrow 0$ the pp -wave space is the same as pure AdS₅, and we see that in this limit (3.62) reduces to the well known straight string solution [24].

Let us now map this solution to pure AdS₅. Under the map (3.9) this solution is mapped into

$$\begin{aligned}\tilde{x}_+ &= \frac{1}{\mu} \tan \frac{\mu\tau}{\sqrt{2}}, & \tilde{x}_- &= -\frac{\tau}{\sqrt{2}} - \frac{\mu}{2} \sigma^2 \tan \frac{\mu\tau}{\sqrt{2}}, \\ \tilde{z} &= \frac{\sigma}{\cos \frac{\mu\tau}{\sqrt{2}}}, & \tilde{x}_i &= 0.\end{aligned}\quad (3.63)$$

On the boundary the straight line in pp -wave coordinates is mapped into a line with tangent shape in pure AdS₅ as shown in Fig. 6. The world-sheet surface is described by the equation

$$\tilde{z} = \sqrt{-\frac{2}{\mu^2 \tilde{x}_+} \left(\tilde{x}_- + \frac{\arctan \mu \tilde{x}_+}{\mu} \right) (1 + \mu^2 \tilde{x}_+^2)}. \quad (3.64)$$

This looks similar to the surface in (3.13) as one can see in Fig. 7 for $\tilde{t} < 0$. The difference with the case in (3.13) is that now the point where the line ends at the boundary is shifted compared to the situation in (3.13). It is obtained from the solution of equation $\tilde{x}_- + \frac{\arctan \mu \tilde{x}_+}{\mu} = 0$. This shows that in the present case the end point on the boundary moves with a speed less than the speed of light. The position of the vertical line is obtained when $\tilde{x}_+ = 0$, i.e. $\tilde{x} = -\tilde{t}$. On the world sheet, the process can again be viewed as an elastic scattering process, while in the spacetime/gauge theory it can be viewed as the scattering between a gluon and a quark moving at a speed less than the speed of light.

Following what was done for the line in the x_+ direction in (3.15), we would like to compute the energy and momentum of string. However, in this case the computation cannot be done exactly because of the complicated form of the solution (3.64), and the fact that the equation $\tilde{x}_- + \frac{\arctan \mu \tilde{x}_+}{\mu} = 0$ cannot be solved explicitly for $\tilde{x}_0 = \tilde{x}_0(\tilde{t})$. This solution gives the limit of integration in the computation of energy which for $\tilde{t} > 0$ is $-\tilde{t} \leq \tilde{x} \leq \tilde{x}_0$. As in the case of the line in the x_+ direction, here we again expect that both energy and momentum of the string depend on time. The time delay for the rounded spike as it




FIG. 6 (color online). Parametric plot of \tilde{z} versus \tilde{x} , for $\mu = \sqrt{2}$ at the boundary $z = 0$.




FIG. 7 (color online). Parametric plot of \tilde{z} versus \tilde{x} , for $\mu = 1$, and $\tilde{t} = -100, -50, -10, -3$.

moves from one site (at $\tilde{t} = -\infty$) to the other (at $\tilde{t} = \infty$) of the vertical line can be computed and we obtain $\Delta\tilde{t} = 2\delta_1 = \frac{2\sqrt{2}}{\mu}$. This is the same as what we had in the case of the solution in the x_+ direction in Sec. III A 1. Since μ is the only physical scale available we expect that asymptotically at $|\tilde{t}| \rightarrow \infty$, the string energy will be again proportional to μ . Upon eliminating μ we expect that in this case the resulting phase shift is again of the form $\delta(E) \sim \ln E$.

It is also interesting to consider a solution obtained from a particle which, in the pp -wave, moves at constant velocity. It interpolates between the one at rest, considered here and the one moving at the speed of light considered in the previous subsection. The solution in flat AdS₅ space can be obtained simply by using the symmetry:

$$\tilde{x}_+ \rightarrow \tilde{x}_+, \quad \tilde{x}_- \rightarrow \frac{1}{\xi^2} \tilde{x}_-, \quad \tilde{x}_i \rightarrow \frac{1}{\xi} \tilde{x}_i, \quad \tilde{z} \rightarrow \frac{1}{\xi} \tilde{z}, \quad (3.65)$$

which gives, from (3.64), the new solution:

$$\tilde{z} = \sqrt{-\frac{2}{\mu^2 \tilde{x}_+} \left(\tilde{x}_- + \xi^2 \frac{\arctan \mu \tilde{x}_+}{\mu} \right) (1 + \mu^2 \tilde{x}_+^2)} \quad (3.66)$$

interpolating between Eq. (3.64) for $\xi = 1$ and Eq. (3.13) for $\xi = 0$. The boundary Wilson loop is described by the equation

$$\tilde{x}_- = -\xi^2 \frac{\arctan \mu \tilde{x}_+}{\mu}. \quad (3.67)$$

It represents a particle coming from infinity at the speed of light which slows down, attaining its minimum speed

$$v = \frac{1 + \partial_+ \tilde{x}_-}{1 - \partial_+ \tilde{x}_-} \Big|_{\tilde{x}_+ = 0} = \frac{1 - \xi^2}{1 + \xi^2} \quad (3.68)$$

at $\tilde{x} = \tilde{t} = 0$. We see that the particle is at rest at the origin when $\xi = 1$ and moves at the speed of light when $\xi = 0$.

2. 2-parallel lines

We extend the above construction and look for a solution which at the boundary $z = 0$ consists of two timelike parallel lines along the t direction (let us assume $b > 0$)

$$t = \tau, \quad x = \pm b, \quad x_i = 0. \quad (3.69)$$

The ansatz for such a solution is

$$t = \tau, \quad x = \sigma, \quad z = z(\sigma), \quad x_i = 0. \quad (3.70)$$

The Nambu action becomes

$$S = \frac{\sqrt{\lambda}}{2\pi} \int d\tau d\sigma \frac{1}{z^2} \sqrt{1 + \left(1 + \frac{\mu^2}{2} z^2\right) z'^2}. \quad (3.71)$$

We want the solution for z to obey the boundary condition $z(\pm b) = 0$. Since the action does not depend explicitly on σ the conserved charge is

$$\frac{1}{z^2} \frac{1}{\sqrt{1 + \left(1 + \frac{\mu^2}{2} z^2\right) z'^2}} = d^2, \quad (3.72)$$

where d is a constant. The differential equation obeyed by z is

$$z' = \pm \frac{1}{z^2 d^2} \sqrt{1 - d^4 z^4}. \quad (3.73)$$

The world-sheet surface reaches a maximum value in the radial direction of AdS₅, i.e. $0 \leq z \leq \frac{1}{d}$, where we assume $d > 0$. The distance between the Wilson lines at the boundary can be expressed as

$$2b = \int_{-b}^b d\sigma = 2d^2 \int_0^{1/d} dz z^2 \sqrt{\frac{1 + \frac{\mu^2}{2} z^2}{1 - d^4 z^4}}. \quad (3.74)$$

Let us note that in the limit $\mu \rightarrow 0$ the equation of motion reduces, as it should, to the equation corresponding to the usual two parallel lines [24] solution

$$z' = \pm \frac{1}{z^2 d^2} \sqrt{1 - d^4 z^4}. \quad (3.75)$$

It is possible to express the integrals of the equation of motion (3.73) and condition (3.74) in general for arbitrary values of μ , d , in terms of the elliptic integrals but the expressions are not very useful, so we do not write them down. Only two of the parameters b , μ , d are independent because of the condition (3.74). Let us see the form of the solution after considering the transformation (3.9) that brings the metric back into pure AdS₅. The explicit solution is complicated in the general case, but we can look at how the Wilson lines at the boundary ($z = 0$) transform

$$\begin{aligned} \tilde{x}_+ &= \frac{1}{\mu} \tan \frac{\mu}{\sqrt{2}} (\pm b + \tau), & \tilde{x}_- &= x_- = \frac{1}{\sqrt{2}} (\pm b - \tau), \\ \tilde{x}_i &= 0. \end{aligned} \quad (3.76)$$

In the pure AdS₅ coordinates, \tilde{t} , \tilde{x} become

$$\begin{aligned} \tilde{t} &= \frac{1}{\sqrt{2}\mu} \tan \frac{\mu}{\sqrt{2}} (\pm b + \tau) - \frac{1}{2} (\pm b - \tau), \\ \tilde{x} &= \frac{1}{\sqrt{2}\mu} \tan \frac{\mu}{\sqrt{2}} (\pm b + \tau) + \frac{1}{2} (\pm b - \tau). \end{aligned} \quad (3.77)$$

While in the pp -wave AdS₅ space, in the plane t, x we had two straight lines in the t direction, in the pure AdS₅ in the plane \tilde{t}, \tilde{x} we have more complicated curves. The curves at $x = \pm b$ are plotted in the new coordinates in Fig. 8. As expected, of course, the curves in the pure AdS₅ coordinates are again timelike.

Let us compute the conserved charges associated to translations in t and x in the pp -wave AdS₅ background. The energy of the string is

$$E = \frac{\sqrt{\lambda}}{2\pi} \int d\sigma \frac{\partial L}{\partial(\partial_\tau t)} = \frac{\sqrt{\lambda}}{2\pi} 2 \int_\epsilon^{1/d} dz \frac{1}{z^2} \sqrt{\frac{1 + \frac{\mu^2}{2} z^2}{1 - d^4 z^4}}, \quad (3.78)$$

where we introduced a cutoff at small z . Let us recall that in fact μ can be set to 1 by rescaling of x_+ and x_- . For simplicity let us set $\mu = 1$. The integral in (3.78) can be done and, expanding the result in small ϵ we obtain




FIG. 8 (color online). Parametric plot of \tilde{t} versus \tilde{x} , for $\mu = \sqrt{2}$ and $b = 1$ at the boundary $z = 0$. The upper curve is the image in the tilde coordinates of the $x = -b$ straightline, and the lower curve corresponds to the $x = b$ straightline.

$$E = \frac{\sqrt{\lambda}}{\pi} \frac{1}{\epsilon} + \frac{\sqrt{\lambda}}{2\sqrt{2}\pi} \frac{1}{d} \left[(1 + 2d^2) K \left[\frac{2d^2 - 1}{4d^2} \right] - 4d^2 E \left[\frac{2d^2 - 1}{4d^2} \right] \right] + O(\epsilon), \quad (3.79)$$

where $E[x]$, $K[x]$ are elliptic integrals. The other conserved charge is

$$P = \frac{\sqrt{\lambda}}{2\pi} \int d\sigma \frac{\partial L}{\partial(\partial_\tau x)} = \frac{\sqrt{\lambda}}{2\pi} \frac{\mu^2}{2} 2 \int_0^{1/d} dz \sqrt{\frac{1 - d^4 z^4}{1 + \frac{\mu^2}{2} z^2}}. \quad (3.80)$$

This integral can be expressed in terms of elliptic integrals but the exact expression is not very illuminating. Let us mention that the integral is convergent so here we do not need a cutoff in z . The charge P should be dependent on the length of the string in the x direction, thus providing a relation between the parameters μ , d , b . Of course, we already exploited such a relation in (3.74).

We observe that for a particular value of d , i.e. $d = \frac{1}{\sqrt{2}}$ (or $d = \frac{\mu}{\sqrt{2}}$ for arbitrary μ) the integration in (3.73) is simpler and the solution, in this case, can be written as a transcendental equation for z

$$-\frac{z}{\sqrt{2}} \sqrt{1 - \frac{z^2}{2}} + \arcsin \frac{z}{\sqrt{2}} + c_1 = \pm \sqrt{2} \sigma d. \quad (3.81)$$

Constant c_1 can be obtained from the boundary conditions that $z(\pm b) = 0$. This gives $c_1 = -\sqrt{2}b$. The negative branch is to be used for positive values of σ . The condition (3.74) can also be easily integrated and it gives an expression for b as $b = \frac{\pi\sqrt{2}}{4}$ (or $b = \frac{\pi\sqrt{2}}{4\mu}$ for arbitrary μ). For this particular case the conserved charges reduce to

$$E = \frac{\sqrt{\lambda}}{\pi} \frac{\sqrt{1 - d^2 \epsilon^2}}{\epsilon} = \frac{\sqrt{\lambda}}{\pi} \frac{1}{\epsilon} + O(\epsilon), \quad P = \sqrt{\lambda} \frac{d}{4}. \quad (3.82)$$

The computation of the action on this solution gives

$$S = \frac{\sqrt{\lambda}}{2\pi} 2T \int_\epsilon^{1/d} dz \frac{1}{z^2} \sqrt{\frac{1 + \frac{\mu^2}{2} z^2}{1 - d^4 z^4}}, \quad (3.83)$$

where we have chosen the τ interval $-\frac{T}{2} \leq \tau \leq \frac{T}{2}$, with T large. Also, we introduced a small cutoff in z to account for the expected divergency near the boundary of AdS₅. Let us recall that because of the condition (3.74) only two parameters are independent, so the action can be expressed in general as $S = S(b, \mu)$. For simplicity let us choose again $\mu = 1$. Doing the integral and expanding in small ϵ we obtain the action

$$S = \frac{\sqrt{\lambda}}{\pi} \frac{T}{\epsilon} + \frac{\sqrt{\lambda}}{2\sqrt{2}\pi} \frac{T}{d} \left[(1 + 2d^2) K \left[\frac{2d^2 - 1}{4d^2} \right] - 4d^2 E \left[\frac{2d^2 - 1}{4d^2} \right] \right] + O(\epsilon). \quad (3.84)$$

The function of d at order $O(\epsilon^0)$ is decreasing. It is interesting to note that for the particular value $d = \frac{1}{\sqrt{2}}$ the $O(\epsilon^0)$ term vanishes. As in [13] one needs to regularize the area by adding a boundary term that cancels the linear divergent term $\frac{1}{\epsilon}$.

So far we discussed the solution in Minkowski space. It turns out that this solution can also be found in the Euclidean space-time and world sheet. The same was found in [25] in the case of other simple solutions that are nontrivial only in the AdS₅ part of AdS₅ × S⁵. One can go from Minkowski to Euclidean by taking $t \rightarrow it$, $\tau \rightarrow i\tau$. The solution (3.70) is also solution in Euclidean space. The equation of motion is again (3.73). In Euclidean space it makes sense to relate the action of a string solution that ends on the boundary on a Wilson loop to the expectation value of the Wilson loop.

IV. CONCLUSIONS

In this paper we have analyzed new methods to find solutions describing open strings moving in AdS₅ and ending in the boundary. According to the AdS/CFT correspondence they are dual to Wilson loops in $\mathcal{N} = 4$ SYM theory. The first solutions were found by using an ansatz leading to linear (and unconstrained) equations of motion. They describe $\frac{1}{4}$ BPS Euclidean Wilson loops with the shape $x_+ = x_+(x_1)$, $x_- = x_2 = 0$ in usual Minkowski light-cone coordinates x_\pm , $x_{1,2}$. Using an inversion we mapped them to closed Wilson loops whose expectation value is $\langle W \rangle = e^{-\sqrt{\lambda}}$. Other, different, solutions were found by considering Wilson loops for $\mathcal{N} = 4$ SYM living in a four dimensional pp -wave. By doing a conformal transformation we can obtain Wilson loops in usual flat space. In this way new interesting solutions were found. They seem to describe gluons interacting with quarks, the gluons being described by spikes coming out from the horizon. We expect these solutions to give a basis for further generalizations and a deeper understanding of Wilson loops in superconformal theories.

ACKNOWLEDGMENTS

We are grateful to P. Argyres, S. Das, R. Leigh, C. Sommerfield and C. Thorn for useful discussions. We are grateful to A. Tseytlin for discussions and comments on the paper. M. K. and A. T. were supported in part by NSF under grant PHY-0653357. The work of R. I. was supported in part by the Purdue Research Foundation.

- [1] M. Kruczenski, J. High Energy Phys. 12 (2002) 024.
- [2] Y. Makeenko, J. High Energy Phys. 01 (2003) 007.
- [3] L. F. Alday and J. M. Maldacena, J. High Energy Phys. 06 (2007) 064.
- [4] Z. Bern, L. J. Dixon, and V. A. Smirnov, Phys. Rev. D **72**, 085001 (2005).
- [5] M. Kruczenski, R. Roiban, A. Tirziu, and A. A. Tseytlin, Nucl. Phys. **B791**, 93 (2008).
- [6] E. I. Buchbinder, Phys. Lett. B **654**, 46 (2007).
- [7] Z. Komargodski and S. S. Razamat, J. High Energy Phys. 01, 044 (2008).
- [8] J. McGreevy and A. Sever, J. High Energy Phys. 02 (2008) 015.
- [9] L. F. Alday, arXiv:0804.0951.
- [10] L. F. Alday and J. Maldacena, J. High Energy Phys. 11 (2007) 068.
- [11] N. Drukker and D. J. Gross, J. Math. Phys. (N.Y.) **42**, 2896 (2001).
- [12] M. Kruczenski and A. A. Tseytlin, Phys. Rev. D **77**, 126005 (2008).
- [13] N. Drukker, D. J. Gross, and H. Ooguri, Phys. Rev. D **60**, 125006 (1999).
- [14] K. Zarembo, Nucl. Phys. **B643**, 157 (2002).
- [15] J. Polchinski, *String Theory. Vol. 2: Superstring Theory and Beyond* (University Press, Cambridge, UK, 1998), p. 531 (Appendix B).
- [16] D. E. Berenstein, R. Corrado, W. Fischler, and J. M. Maldacena, Phys. Rev. D **59**, 105023 (1999).
- [17] J. K. Erickson, G. W. Semenoff, and K. Zarembo, Nucl. Phys. **B582**, 155 (2000).
- [18] M. Kruczenski, and A. Tirziu, J. High Energy Phys. 05 (2008) 064.
- [19] V. Pestun, arXiv:0712.2824.
- [20] A. Mikhailov, arXiv:hep-th/0305196.
- [21] D. Brecher, A. Chamblin, and H. S. Reall, Nucl. Phys. **B607**, 155 (2001).
- [22] M. Kruczenski, J. High Energy Phys. 08 (2005) 014; M. Kruczenski, J. Russo, and A. A. Tseytlin, J. High Energy Phys. 10 (2006) 002; R. Ishizeki and M. Kruczenski, Phys. Rev. D **76**, 126006 (2007); A. E. Mosaffa and B. Safarzadeh, J. High Energy Phys. 08 (2007) 017; S. Ryang, J. High Energy Phys. 12 (2006) 043; R. Ishizeki, M. Kruczenski, M. Spradlin, and A. Volovich, J. High Energy Phys. 02 (2008) 009.
- [23] R. Jackiw and G. Woo, Phys. Rev. D **12**, 1643 (1975).
- [24] N. Drukker, D. J. Gross, and A. A. Tseytlin, J. High Energy Phys. 04 (2000) 021.
- [25] A. A. Tseytlin and K. Zarembo, Phys. Rev. D **66**, 125010 (2002).