

*Physics*

*Physics Research Publications*

---

*Purdue University*

*Year 2008*

---

Study of the decays  $D^0 \rightarrow \pi^{\pm} e^{\mp} \nu(e)$ ,  $D^0 \rightarrow K^{\pm} e^{\mp} \nu(e)$ ,  
 $D^+ \rightarrow \pi^0 e^+ \nu(e)$ , and  $D^+ \rightarrow \bar{K}^0 e^+ \nu(e)$

D. Cronin-Hennessy, K. Y. Gao, D. T. Gong, J. Hietala, Y. Kubota, T. Klein, B. W. Lang, R. Poling, A. W. Scott, A. Smith, P. Zweber, S. Dobbs, Z. Metreveli, K. K. Seth, A. Tomaradze, J. Ernst, H. Severini, S. A. Dytman, W. Love, V. Savinov, O. Aquines, Z. Li, A. Lopez, S. Mehrabyan, H. Mendez, J. Ramirez, G. S. Huang, D. H. Miller, V. Pavlunin, B. Sanghi, I. P. J. Shipsey, B. Xin, G. S. Adams, M. Anderson, J. P. Cummings, I. Danko, J. Napolitano, Q. He, J. Insler, H. Muramatsu, C. S. Park, E. H. Thorndike, F. Yang, T. E. Coan, Y. S. Gao, F. Liu, M. Artuso, S. Blusk, J. Butt, J. Li, N. Menea, R. Mountain, S. Nisar, K. Randrianarivony, R. Redjimi, R. Sia, T. Skwarnicki, S. Stone, J. C. Wang, K. Zhang, S. E. Csorna, G. Bonvicini, D. Cinabro, M. Dubrovin, A. Lincoln, D. M. Asner, K. W. Edwards, R. A. Briere, I. Brock, J. Chen, T. Ferguson, G. Tatishvili, H. Vogel, M. E. Watkins, J. L. Rosner, N. E. Adam, J. P. Alexander, K. Berkelman, D. G. Cassel, J. E. Duboscq, K. M. Ecklund, R. Ehrlich, L. Fields, L. Gibbons, R. Gray, S. W. Gray, D. L. Hartill, B. K. Heltsley, D. Hertz, C. D. Jones, J. Kandaswamy, D. L. Kreinick, V. E. Kuznetsov, H. Mahlke-Kruger, P. U. E. Onyisi, J. R. Patterson, D. Peterson, J. Pivarski, D. Riley, A. Ryd, A. J. Sadoff, H. Schwarthoff, X. Shi, S. Stroiney, W. M. Sun, T. Wilksen, M. Weinberger, S. B. Athar, R. Patel, V. Potlia, J. Yelton, P. Rubin, C. Cawfield, B. I. Eisenstein, I. Karliner, D. Kim, N. Lowrey, P. Naik, C. Sedlack, M. Selen, E. J. White, J. Wiss, M. R. Shepherd, D. Besson, and T. K. Pedlar

This paper is posted at Purdue e-Pubs.

[http://docs.lib.purdue.edu/physics\\_articles/867](http://docs.lib.purdue.edu/physics_articles/867)

## Study of the Decays $D^0 \rightarrow \pi^- e^+ \nu_e$ , $D^0 \rightarrow K^- e^+ \nu_e$ , $D^+ \rightarrow \pi^0 e^+ \nu_e$ , and $D^+ \rightarrow \bar{K}^0 e^+ \nu_e$

D. Cronin-Hennessy,<sup>1</sup> K. Y. Gao,<sup>1</sup> D. T. Gong,<sup>1</sup> J. Hietala,<sup>1</sup> Y. Kubota,<sup>1</sup> T. Klein,<sup>1</sup> B. W. Lang,<sup>1</sup> R. Poling,<sup>1</sup> A. W. Scott,<sup>1</sup> A. Smith,<sup>1</sup> P. Zweber,<sup>1</sup> S. Dobbs,<sup>2</sup> Z. Metreveli,<sup>2</sup> K. K. Seth,<sup>2</sup> A. Tomaradze,<sup>2</sup> J. Ernst,<sup>3</sup> H. Severini,<sup>4</sup> S. A. Dytman,<sup>5</sup> W. Love,<sup>5</sup> V. Savinov,<sup>5</sup> O. Aquines,<sup>6</sup> Z. Li,<sup>6</sup> A. Lopez,<sup>6</sup> S. Mehrabyan,<sup>6</sup> H. Mendez,<sup>6</sup> J. Ramirez,<sup>6</sup> G. S. Huang,<sup>7</sup> D. H. Miller,<sup>7</sup> V. Pavlunin,<sup>7</sup> B. Sanghi,<sup>7</sup> I. P. J. Shipsey,<sup>7</sup> B. Xin,<sup>7</sup> G. S. Adams,<sup>8</sup> M. Anderson,<sup>8</sup> J. P. Cummings,<sup>8</sup> I. Danko,<sup>8</sup> J. Napolitano,<sup>8</sup> Q. He,<sup>9</sup> J. Insler,<sup>9</sup> H. Muramatsu,<sup>9</sup> C. S. Park,<sup>9</sup> E. H. Thorndike,<sup>9</sup> F. Yang,<sup>9</sup> T. E. Coan,<sup>10</sup> Y. S. Gao,<sup>10</sup> F. Liu,<sup>10</sup> M. Artuso,<sup>11</sup> S. Blusk,<sup>11</sup> J. Butt,<sup>11</sup> J. Li,<sup>11</sup> N. Menea,<sup>11</sup> R. Mountain,<sup>11</sup> S. Nisar,<sup>11</sup> K. Randrianarivony,<sup>11</sup> R. Redjimi,<sup>11</sup> R. Sia,<sup>11</sup> T. Skwarnicki,<sup>11</sup> S. Stone,<sup>11</sup> J. C. Wang,<sup>11</sup> K. Zhang,<sup>11</sup> S. E. Csorna,<sup>12</sup> G. Bonvicini,<sup>13</sup> D. Cinabro,<sup>13</sup> M. Dubrovin,<sup>13</sup> A. Lincoln,<sup>13</sup> D. M. Asner,<sup>14</sup> K. W. Edwards,<sup>14</sup> R. A. Briere,<sup>15</sup> I. Brock,<sup>15</sup> J. Chen,<sup>15</sup> T. Ferguson,<sup>15</sup> G. Tatishvili,<sup>15</sup> H. Vogel,<sup>15</sup> M. E. Watkins,<sup>15</sup> J. L. Rosner,<sup>16</sup> N. E. Adam,<sup>17</sup> J. P. Alexander,<sup>17</sup> K. Berkelman,<sup>17</sup> D. G. Cassel,<sup>17</sup> J. E. Duboscq,<sup>17</sup> K. M. Ecklund,<sup>17</sup> R. Ehrlich,<sup>17</sup> L. Fields,<sup>17</sup> L. Gibbons,<sup>17</sup> R. Gray,<sup>17</sup> S. W. Gray,<sup>17</sup> D. L. Hartill,<sup>17</sup> B. K. Heltsley,<sup>17</sup> D. Hertz,<sup>17</sup> C. D. Jones,<sup>17</sup> J. Kandaswamy,<sup>17</sup> D. L. Kreinick,<sup>17</sup> V. E. Kuznetsov,<sup>17</sup> H. Mahlke-Krüger,<sup>17</sup> P. U. E. Onyisi,<sup>17</sup> J. R. Patterson,<sup>17</sup> D. Peterson,<sup>17</sup> J. Pivarski,<sup>17</sup> D. Riley,<sup>17</sup> A. Ryd,<sup>17</sup> A. J. Sadoff,<sup>17</sup> H. Schwarhoff,<sup>17</sup> X. Shi,<sup>17</sup> S. Stroiney,<sup>17</sup> W. M. Sun,<sup>17</sup> T. Wilksen,<sup>17</sup> M. Weinberger,<sup>17</sup> S. B. Athar,<sup>18</sup> R. Patel,<sup>18</sup> V. Potlia,<sup>18</sup> J. Yelton,<sup>18</sup> P. Rubin,<sup>19</sup> C. Cawfield,<sup>20</sup> B. I. Eisenstein,<sup>20</sup> I. Karliner,<sup>20</sup> D. Kim,<sup>20</sup> N. Lowrey,<sup>20</sup> P. Naik,<sup>20</sup> C. Sedlack,<sup>20</sup> M. Selen,<sup>20</sup> E. J. White,<sup>20</sup> J. Wiss,<sup>20</sup> M. R. Shepherd,<sup>21</sup> D. Besson,<sup>22</sup> and T. K. Pedlar<sup>23</sup>

(CLEO Collaboration)

<sup>1</sup>University of Minnesota, Minneapolis, Minnesota 55455, USA

<sup>2</sup>Northwestern University, Evanston, Illinois 60208, USA

<sup>3</sup>State University of New York at Albany, Albany, New York 12222, USA

<sup>4</sup>University of Oklahoma, Norman, Oklahoma 73019, USA

<sup>5</sup>University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA

<sup>6</sup>University of Puerto Rico, Mayaguez, Puerto Rico 00681

<sup>7</sup>Purdue University, West Lafayette, Indiana 47907, USA

<sup>8</sup>Rensselaer Polytechnic Institute, Troy, New York 12180, USA

<sup>9</sup>University of Rochester, Rochester, New York 14627, USA

<sup>10</sup>Southern Methodist University, Dallas, Texas 75275, USA

<sup>11</sup>Syracuse University, Syracuse, New York 13244, USA

<sup>12</sup>Vanderbilt University, Nashville, Tennessee 37235, USA

<sup>13</sup>Wayne State University, Detroit, Michigan 48202, USA

<sup>14</sup>Carleton University, Ottawa, Ontario, Canada K1S 5B6

<sup>15</sup>Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

<sup>16</sup>Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA

<sup>17</sup>Cornell University, Ithaca, New York 14853, USA

<sup>18</sup>University of Florida, Gainesville, Florida 32611, USA

<sup>19</sup>George Mason University, Fairfax, Virginia 22030, USA

<sup>20</sup>University of Illinois, Urbana-Champaign, Illinois 61801, USA

<sup>21</sup>Indiana University, Bloomington, Indiana 47405, USA

<sup>22</sup>University of Kansas, Lawrence, Kansas 66045, USA

<sup>23</sup>Luther College, Decorah, Iowa 52101, USA

(Received 5 December 2007; published 27 June 2008)

By using  $1.8 \times 10^6$   $D\bar{D}$  pairs, we have measured  $\mathcal{B}(D^0 \rightarrow \pi^- e^+ \nu_e) = 0.299(11)(9)\%$ ,  $\mathcal{B}(D^+ \rightarrow \pi^0 e^+ \nu_e) = 0.373(22)(13)\%$ ,  $\mathcal{B}(D^0 \rightarrow K^- e^+ \nu_e) = 3.56(3)(9)\%$ , and  $\mathcal{B}(D^+ \rightarrow \bar{K}^0 e^+ \nu_e) = 8.53(13) \times (23)\%$  and have studied the  $q^2$  dependence of the form factors. By combining our results with recent lattice calculations, we obtain  $|V_{cd}| = 0.217(9)(4)(23)$  and  $|V_{cs}| = 1.015(10)(11)(106)$ .

DOI: [10.1103/PhysRevLett.100.251802](https://doi.org/10.1103/PhysRevLett.100.251802)

PACS numbers: 13.20.Fc, 12.15.Hh, 14.40.Lb

Study of the semileptonic decays of  $D$  mesons plays an important role in determining the Cabibbo-Kobayashi-Maskawa (CKM) matrix [1]. These decays both allow determination of  $|V_{cs}|$  and  $|V_{cd}|$  and provide rigorous tests

[2] of lattice QCD (LQCD) calculations (e.g., [3]). The tests can be approached by comparing measured elements to those constrained by matrix unitarity [4] or by comparing the measured and calculated ratios of semileptonic and

leptonic branching fractions, which are independent of the CKM matrix. Verification of LQCD calculations at the few percent level will provide validation for their application to the  $B$  system.

This Letter presents a study of the  $D^0 \rightarrow K^- e^+ \nu_e$ ,  $D^0 \rightarrow \pi^- e^+ \nu_e$ ,  $D^+ \rightarrow \bar{K}^0 e^+ \nu_e$ , and  $D^+ \rightarrow \pi^0 e^+ \nu_e$  decay modes (charge conjugate modes implied). A companion article [5] provides a more detailed description. The results are based on  $281 \text{ pb}^{-1}$  of  $e^+e^-$  data at the  $\psi(3770)$  resonance ( $1.8 \times 10^6$   $D\bar{D}$  pairs) collected with the CLEO-c detector at the Cornell Electron Storage Ring (CESR) [6]. The data are a superset of those used for the first CLEO-c semileptonic measurements [7]. For each mode, we measure the partial branching fractions in five  $q^2 = m_{\ell\nu}^2$  ranges. Summing the rates yields the total branching fraction; fitting the rates constrains form factor (FF) shapes; comparing to LQCD calculations [3] determines the CKM elements  $|V_{cd}|$  and  $|V_{cs}|$ .

The analysis technique rests upon association of the missing energy and momentum in an event with the neutrino four-momentum [8], enabled by the Hermeticity and excellent resolution of the CLEO-c detector. Charged particles are detected over 93% of the solid angle by two wire tracking chambers within a 1.0 T solenoid magnet. The momentum resolution is 0.6% at 800 MeV/c. Specific ionization and a ring imaging Čerenkov detector (RICH) provide charged particle identification; a CsI(Tl) electromagnetic calorimeter provides photon detection over 93% of  $4\pi$  and a  $\pi^0$  mass resolution of  $\sim 6 \text{ MeV}/c^2$ .

Electron candidates above 200 MeV/c are identified over 90% of the solid angle by combining specific ionization information with calorimetric, RICH, and tracking measurements. To reduce sensitivity to final state radiation (FSR), we add photons within  $3.5^\circ$  of the electron flight direction to the electron momentum. A  $\pi^0$  candidate must have a  $\gamma\gamma$  mass within 2.5 standard deviations ( $\sigma$ ) of the  $\pi^0$  mass.  $K_S^0$  candidates are reconstructed by using a vertex fit to candidate  $\pi^+\pi^-$  daughter tracks. The  $\pi^+\pi^-$  mass must be within  $4.5\sigma$  of the  $K_S^0$  mass.

The missing four-momentum in an event is given by  $p_{\text{miss}} = (E_{\text{miss}}, \vec{p}_{\text{miss}}) = p_{\text{total}} - \sum p_{\text{charged}} - \sum p_{\text{neutral}}$ , where the event four-momentum  $p_{\text{total}}$  is known from the energy and crossing angle of the CESR beams. The charged and neutral particles included in the sums pass selection criteria designed to achieve the best possible  $|\vec{p}_{\text{miss}}|$  resolution by balancing the efficiency for detecting true particles against the rejection of false ones [5].

Association of  $p_{\text{miss}}$  with the neutrino is valid only if the event contains no more than one neutrino and all true particles are detected. We thus exclude events that have either more than one electron [9] or nonzero net charge. The core  $|\vec{p}_{\text{miss}}|$  resolution in our signal Monte Carlo (MC) events satisfying these criteria is  $\sim 15 \text{ MeV}/c$ .

Background sources include hadrons misidentified as electrons (fake electrons), noncharm continuum produc-

tion, and  $D\bar{D}$  processes other than signal. Background suppression criteria were optimized with GEANT-based [10] MC samples independent of those used in fitting.

We require  $M_{\text{miss}}^2 \equiv E_{\text{miss}}^2 - |\vec{p}_{\text{miss}}|^2$  to be consistent with a massless neutrino:  $|M_{\text{miss}}^2/2|\vec{p}_{\text{miss}}|| < 0.2 \text{ GeV}/c^3$ . Since the  $|\vec{p}_{\text{miss}}|$  resolution is roughly half that of  $E_{\text{miss}}$ , in subsequent calculations we take  $p_\nu \equiv (|\vec{p}_{\text{miss}}|, \vec{p}_{\text{miss}})$ .

Semileptonic decays  $D \rightarrow h e \nu_e$ , where  $h = \pi$  or  $K$ , are identified by using four-momentum conservation. Specifically, energy conservation demands  $\Delta E \equiv (E_h + E_e + E_\nu) - E_{\text{beam}}$  to be close to zero within our 20 MeV resolution, and we require  $-0.06 < \Delta E < 0.10 \text{ GeV}$ . Since the  $|\vec{p}_\nu|$  resolution dominates  $\Delta E$  resolution, we improve our  $p_\nu$  measurement by scaling it by the factor  $\zeta$  satisfying  $\Delta E = (E_h + E_e + \zeta E_\nu) - E_{\text{beam}} = 0$ . The  $D$  momentum constraint is recast as the beam-constrained mass  $M_{\text{bc}} \equiv \sqrt{E_{\text{beam}}^2 - |\vec{p}_h + \vec{p}_e + \zeta \vec{p}_\nu|^2}$ , which peaks near the  $D$  mass for signal. Our  $M_{\text{bc}}$  ( $q^2 \equiv [\zeta p_\nu + p_e]^2$ ) resolution is  $4 \text{ MeV}/c^2$  ( $0.01 \text{ GeV}^2/c^4$ ), independent of  $q^2$ .

For the Cabibbo-favored modes, the background remaining after these requirements is only a few percent of the signal. For the Cabibbo-suppressed modes, significant background remains from signal-mode cross feed and from the related modes  $D^+ \rightarrow K_L^0 e^+ \nu_e$  and  $D^+ \rightarrow K_S^0(\pi^0 \pi^0) e^+ \nu_e$ , where the  $(\pi^0 \pi^0)$  indicates the  $K_S^0$  decay mode. Restricting the  $\Delta E$  of the nonsignal side of the event reduces these backgrounds. To further reduce cross-feed backgrounds and simplify statistical interpretation, we limit multiple  $D^0$  ( $D^+$ ) candidates with  $M_{\text{bc}} > 1.794 \text{ GeV}/c^2$  by choosing that with the smallest  $|\Delta E|$ . The  $D^+ \rightarrow \pi^0 e^+ \nu_e$  requirements are stricter: Candidates must have the smallest  $|\Delta E|$  of any final state candidate in the event, and the event must contain no  $D^0 \rightarrow K^- e^+ \nu_e$  candidate. The multiple candidate requirements affect about 13% of  $\pi^0$ , 9% of  $\pi^\pm$ , 8% of  $K_S$ , and 3% of  $K^\pm$  candidates. The average final background fraction ( $q^2$  dependent) in the pion modes is about 20%.

To extract branching fractions, we perform a simultaneous binned maximum likelihood fit [11] to the  $M_{\text{bc}}$  distributions of the four signal modes in five  $q^2$  ranges. The simultaneity guarantees self-consistent rates for misreconstruction of one signal process as another (cross feed) and for background from the related  $K^0$  processes. We use 14 equal  $M_{\text{bc}}$  bins over the range 1.794–1.878 GeV/c<sup>2</sup>.

We fit the data to the signal and five background components. The signal-mode MC components are based on EVTGEN [12] with modified pole-model [Becirevic and Kaidalov (BK) parametrization] FFs [13] and parameters from the most recent unquenched LQCD calculation [3]. Several corrections, relating to inclusive  $D$  decay and reconstruction (see Ref. [5]), are applied to our GEANT-based [10] MC samples. These lead to few percent (or less) changes in the measured yields and are determined precisely enough (by using a large  $D\bar{D}$  sample with one fully

reconstructed  $D$  meson per event) to yield subpercent systematic uncertainties. We are also sensitive to the signal efficiency and kinematic distortions due to FSR. Our signal MC simulation includes FSR distributed according to the leading-order kaon leading-order radiation [14] calculation applied to charm decay.

To reduce our sensitivity to FFs, we extract an independent rate for each of the five  $q^2$  intervals in each reconstructed mode (a total of 20 yields).

We use MC samples to describe the  $D\bar{D}$  background and continuum contributions. We normalize the continuum components by using their cross sections at the  $\psi(3770)$  and the measured data luminosity. The nonsignal  $D\bar{D}$  sample was generated by using EVTGEN, with decay parameters updated to reflect our best knowledge of  $D$  meson decays. This component floats separately for each reconstructed mode but is fixed over the five  $q^2$  regions within that mode, reducing our sensitivity to inaccuracies in the  $D$  decay model. Finally, we input explicit MC components for  $D^+ \rightarrow K_L^0 e^+ \nu_e$  and  $D^+ \rightarrow K_S^0(\pi^0 \pi^0) e^+ \nu_e$ . Their normalization in each  $q^2$  range in the fit is scaled by the appropriate amplitude or branching fraction factor to the corresponding  $D^+ \rightarrow K_S^0(\pi^+ \pi^-) e^+ \nu_e$  parameter.

The contributions of events with fake electrons are evaluated by weighting hadron-momentum spectra in candidate events with misidentification probabilities measured in other CLEO-c data. This component is included with a fixed normalization in the fit.

We allow the fit to adjust the  $M_{bc}$  resolution in the  $D^0 \rightarrow \pi^- e^+ \nu_e$ ,  $D^0 \rightarrow K^- e^+ \nu_e$ , and  $D^+ \rightarrow \pi^0 e^+ \nu_e$  modes by smearing the distributions with a Gaussian. The signal MC  $M_{bc}$  resolution of  $\sim 3.5$  MeV/ $c^2$  thereby increases to match the data resolution of  $\sim 4$  MeV/ $c^2$ .

Figure 1 shows the  $M_{bc}$  distributions summed over the five  $q^2$  ranges, with the fit results overlaid. The two highest bins are not fit. Our  $-2 \ln \mathcal{L}$  behavior should be approximately  $\chi^2$ -like, and our fit yields  $-2 \ln \mathcal{L} = 275.5$  for  $280 - 27 = 253$  degrees of freedom.

We obtain branching fractions (see Table I) for each  $q^2$  region by combining the efficiency-corrected yields from the fit with the number of  $D^0 \bar{D}^0$  ( $N_{D^0 \bar{D}^0}$ ) and  $D^+ D^-$  ( $N_{D^+ D^-}$ ) pairs for our sample. An independent study of hadronic  $D$  decays [15] finds  $N_{D^0 \bar{D}^0} = 1.031(16) \times 10^6$  and  $N_{D^+ D^-} = 0.819(13) \times 10^6$ . We find ratios of branching fractions  $R_0 = \mathcal{B}(D^0 \rightarrow \pi^- e^+ \nu_e) / \mathcal{B}(D^0 \rightarrow K^- e^+ \nu_e) = 8.41(32)(13)\%$  and  $R_+ = \mathcal{B}(D^+ \rightarrow \pi^0 e^+ \nu_e) / \mathcal{B}(D^+ \rightarrow \bar{K}^0 e^+ \nu_e) = 4.37(27)(12)\%$  and partial-width ratios  $I_\pi = \Gamma(D^0 \rightarrow \pi^- e^+ \nu_e) / \Gamma(D^+ \rightarrow \pi^0 e^+ \nu_e) = 2.03(14)(8)$  and  $I_K = \Gamma(D^0 \rightarrow K^- e^+ \nu_e) / \Gamma(D^+ \rightarrow \bar{K}^0 e^+ \nu_e) = 1.06(2)(3)$  by using lifetimes from Ref. [4]. Isospin symmetry predicts  $I_\pi = 2$  and  $I_K = 1$ .

The systematic uncertainty (see Ref. [5]) is dominated by the uncertainty in the number of  $D\bar{D}$  pairs and in neutrino reconstruction simulation. The latter includes inaccuracies in the detector simulation and in the decay




FIG. 1. The  $M_{bc}$  distributions, integrated over  $q^2$ , for data (points), the signal MC (clear), and cross-feed and nonsignal  $D\bar{D}$  MC (gray) and continuum MC (light gray) fit components. The  $e^+$  fake component is negligible on this scale.

model of the nonsignal  $D$ , which were evaluated mainly by using events with a reconstructed hadronic decay. We evaluate  $q^2$ -dependent systematic biases for the efficiency of finding and identifying signal hadrons, identifying signal electrons and fake electron rates, as well as for uncertainties that affect the cross-feed rates, such as those associated with nonsignal  $\pi^0$  and  $\pi^-$  production spectra, and  $K^-$  faking  $\pi^-$ . We correct statistically significant biases and propagate the uncertainty of each study into our measurement uncertainty. The remaining systematic uncertainties include  $M_{bc}$  resolution, the effect of the single-electron requirement, MC FSR modeling, dependence on FFs, and the  $N_{D\bar{D}}$  determinations.

Our primary FF shape analysis utilizes a series expansion that has been widely advocated as a physical description of heavy meson FFs [16–19]:

$$f_+(q^2) = \frac{1}{P(q^2)\phi(q^2, t_0)} \sum_{k=0}^{\infty} a_k(t_0)[z(q^2, t_0)]^k. \quad (1)$$

The expansion results from an analytic continuation of the FF into the complex  $t = q^2$  plane, with a branch cut on the real axis for  $t > (M_D + M_{K,\pi})^2$  that is mapped by  $z(t, t_0) = (\sqrt{t_+ - t} - \sqrt{t_+ - t_0}) / (\sqrt{t_+ - t} + \sqrt{t_+ - t_0})$  onto the unit circle. The constants  $t_{\pm} \equiv (M_D \pm m_{K,\pi})^2$ , and  $t_0$  is the (arbitrary)  $q^2$  value that maps to  $z = 0$ . The physical region is restricted to  $|z| < 1$ , so good convergence is expected.  $P(q^2)$  accommodates subthreshold resonances:  $P(q^2) = 1$  for  $D \rightarrow \pi$  and  $P(q^2) = z(q^2, M_{D_s^*}^2)$  for  $D \rightarrow K$ . The function  $\phi(q^2, t_0)$  can be any analytic function. We report  $a_k$  parameters for  $t_0 = 0$  and the “standard” choice for  $\phi$  (see, e.g., Ref. [19]) that arises in studies of unitarity bounds on  $\sum a_k^2$ .

TABLE I. Branching fractions (top) and FF fits for the series (middle) and simple pole (bottom left) and modified pole (bottom right) parametrizations. The  $\pi^0 e^+ \nu$  FF results are isospin-corrected. Errors are (stat)(syst)[(theor)]. The correlation coefficients ( $\rho$  or  $\rho_{ij}$ ) are for the combined stat + syst uncertainties. The  $a_i$  normalization uses  $|V_{cs}| = 0.976$  and  $|V_{cd}| = 0.224$ .

$q^2$ (GeV <sup>2</sup> /c <sup>4</sup> )	<0.4	0.4–0.8	0.8–1.2	1.2–1.6	$\geq 1.6$	Total	$ V_{cq} $
$\mathcal{B}(\pi^- e^+ \nu_e)(\%)$	0.070(5)(3)	0.059(5)(2)	0.060(5)(2)	0.044(4)(2)	0.066(5)(2)	0.299(11)(9)	0.218(11)(5)(23)
$\mathcal{B}(\pi^0 e^+ \nu_e)(\%)$	0.084(10)(4)	0.097(11)(4)	0.062(9)(3)	0.063(10)(2)	0.067(11)(3)	0.373(22)(13)	0.216(17)(6)(23)
$\mathcal{B}(K^- e^+ \nu_e)(\%)$	1.441(21)(35)	1.048(18)(28)	0.681(15)(18)	0.340(11)(10)	0.048(5)(12)	3.557(33)(90)	1.023(13)(13)(107)
$\mathcal{B}(\bar{K}^0 e^+ \nu_e)(\%)$	3.436(82)(93)	2.544(73)(69)	1.589(58)(44)	0.821(42)(24)	0.139(18)(5)	8.53(13)(23)	1.004(20)(15)(105)
Decay	$a_0$	$a_1$	$a_2$	$\rho_{01}$ $\rho_{02}$ $\rho_{12}$	$ V_{cq} f_+(0)$	$1 + 1/\beta - \delta$	$\rho$ $P(\chi^2)$
$\pi^- e^+ \nu_e$	0.044(2)(1)	-0.18(7)(2)	-0.03(35)(12)	0.81 0.71 0.96	0.140(7)(3)	1.30(37)(12)	-0.85 0.38
$\pi^0 e^+ \nu_e$	0.044(3)(1)	-0.23(11)(2)	-0.60(57)(15)	0.80 0.67 0.95	0.138(11)(4)	1.58(60)(13)	-0.86 0.24
$K^- e^+ \nu_e$	0.0234(3)(3)	-0.009(21)(7)	0.52(28)(6)	0.62 0.56 0.96	0.747(9)(9)	0.62(13)(4)	-0.62 0.89
$\bar{K}^0 e^+ \nu_e$	0.0224(4)(3)	0.009(32)(7)	0.76(42)(8)	0.72 0.64 0.96	0.733(14)(11)	0.51(20)(4)	-0.72 0.44
Decay	$ V_{cq} f_+(0)$	$m_{\text{pole}}$ (GeV/c <sup>2</sup> )	$\rho$ $P(\chi^2)$		$ V_{cq} f_+(0)$	$\alpha_{\text{BK}}$	$\rho$ $P(\chi^2)$
$\pi^- e^+ \nu_e$	0.146(4)(2)	1.87(3)(1)	0.63 0.21		0.142(4)(2)	0.37(8)(3)	-0.75 0.37
$\pi^0 e^+ \nu_e$	0.149(6)(3)	1.97(7)(2)	0.65 0.11		0.147(7)(4)	0.14(16)(4)	-0.75 0.13
$K^- e^+ \nu_e$	0.735(5)(9)	1.97(3)(1)	0.36 0.26		0.732(6)(9)	0.21(5)(3)	-0.42 0.12
$\bar{K}^0 e^+ \nu_e$	0.710(8)(10)	1.96(4)(2)	0.53 0.13		0.708(9)(10)	0.22(8)(3)	-0.59 0.07

For comparison purposes, we provide results based on the simple and modified pole models [13]. The latter introduces the shape parameter  $\alpha_{\text{BK}}$  to give  $f_+(q^2) = f_+(0)/[(1 - q^2)(1 - \alpha_{\text{BK}}q^2)]$ , with  $q^2 \equiv q^2/m_{\text{pole}}^2$ . These parametrizations can typically accommodate the FF shapes observed in previous measurements but only with parameters that deviate from the underlying physical motivation [20]. Note that differing experimental sensitivities across phase space can result in differing parameter values for a nonphysical parametrization.

Each parametrization is fit to our measured rates for the five  $q^2$  regions; each parameter systematic uncertainty is obtained from a fit to the rates for that systematic variation. Table I summarizes the results; Fig. 2 compares the three fits in our most precise mode  $D^0 \rightarrow K^- e^+ \nu_e$ . For the series expansion, we also express our results as physical observables: the intercept  $|V_{cq}|f_+(0)$  and  $1 + 1/\beta - \delta \propto (df_+/dq^2)/f_+|_{q^2=0}$  [19], which represents the effects of gluon hard-scattering ( $\delta$ ) and scaling violations ( $\beta$ ).  $D^0$  and  $D^+$  results agree well.

For the series expansion, our kaon data prefer a nonzero quadratic  $z$  term. The probability of  $\chi^2$  [ $P(\chi^2)$ ] improves from 29% (22%) to 89% (44%) with that additional term for the  $K^-$  ( $\bar{K}^0$ ) fit. The pion measurements lack the sensitivity to probe this term, and two and three parameter fits yield similar results for the first two parameters. Since a quadratic term appears preferred for the kaons, however, we include that term in our series fits to the pion data to improve the probability that our shape uncertainties bracket the true FF shape. While three of the central values for  $a_2$  are an order of magnitude larger than the other terms, regions of parameter space with  $a_2$  of similar magnitude to  $a_0$  and  $a_1$  fall well within the 90% hypercontour for the fit, so no conclusion can be drawn about the size of

$a_2$  or (potential lack of) convergence of the series from these data.

In the simple pole model, we fit for the intercept and the pole mass  $m_{\text{pole}}$ . In the modified pole model, we fix  $m_{\text{pole}}$  at its physical value and fit for the intercept and  $\alpha_{\text{BK}}$ , which determines the effective higher pole contribution. We obtain reasonable  $\chi^2$  values, but the pole masses deviate from  $M_{D_s^*}$  ( $M_{D^*}$ ) in the kaon (pion) modes by over  $3\sigma$  for the most precise fits. The  $1 + 1/\beta - \delta$  value for the series expansion fit to the  $K^- e^+ \nu_e$  data is over  $3\sigma$  from the value of  $\sim 2$  necessary for physical validity of the BK parametrization, while our values for the BK  $\alpha_{\text{BK}}$  parameters from the kaon modes imply  $1 + 1/\beta - \delta$  values tens of  $\sigma$  away. Overall,  $P(\chi^2)$  improves noticeably for our preferred  $z$  expansion fit relative to these pole fits.

We extract (Table I)  $|V_{cd}|$  and  $|V_{cs}|$  from the  $|V_{cq}|f_+(0)$  of our series expansion fits using  $f_+^{(D \rightarrow \pi)}(0) = 0.64(3)(6)$  and  $f_+^{(D \rightarrow K)}(0) = 0.73(3)(7)$  from unquenched LQCD [3].




FIG. 2. The  $D^0 \rightarrow K^- e^+ \nu_e$  form factor fits, normalized to the series expansion result.



The  $D^0$  and  $D^+$  averages (heeding correlations) are  $|V_{cd}| = 0.217(9)(4)(23)$  and  $|V_{cs}| = 1.015(10)(11)(106)$ . Discretization uncertainty in the charm quark action dominates the  $f_+$  uncertainty, listed last.

We also extract the ratio  $|V_{cd}|/|V_{cs}|$  from the ratio of our measured FFs. Averaging over  $D^0$  and  $D^+$  modes, with correlations accounted for, gives  $|V_{cd}|f^{(D\rightarrow\pi)}(0)/|V_{cs}|f^{(D\rightarrow K)}(0) = 0.188(8)(2)$ . A recent light cone sum rules calculation [21] gives  $f^{(D\rightarrow\pi)}(0)/f^{(D\rightarrow K)}(0) = 0.84(4)$ , giving  $|V_{cd}|/|V_{cs}| = 0.223(10)(3)(11)$ .

In summary, we have measured branching fractions and their ratios for four semileptonic  $D$  decay modes in five  $q^2$  bins. The branching fraction results are the most precise ever measured and agree well with world averages. Our modified pole  $\alpha_{\text{BK}}$  parameter results agree within  $1.3\sigma$  with previous determinations by CLEO III [22], FOCUS [23], and  $Ke\nu$  results from Belle [24] but show over  $3\sigma$  disagreement with Belle  $K\mu\nu$  results and LQCD fits. The  $\alpha_{\text{BK}}$  parameters obtained with our individual  $Ke\nu$  results are separated from the recent *BABAR* result [25] by about  $2.5\sigma$ . Our  $z$  expansion results agree with *BABAR*'s at about the  $2\sigma$  level, depending on the total level of correlation between the *BABAR*  $r_1$  and  $r_2$  parameters. We have made the most precise CKM determinations from  $D$  semileptonic decays to date, and the results agree well with neutrino-based determinations of  $|V_{cd}|$  and charm-tagged  $W$  decay measurements of  $|V_{cs}|$  [4]. Overall, these measurements represent a marked improvement in our knowledge of  $D$  semileptonic decay.

We gratefully acknowledge the effort of the CESR staff in providing us with excellent luminosity and running conditions. This work was supported by the A.P. Sloan Foundation, the National Science Foundation, the U.S. Department of Energy, and the Natural Sciences and Engineering Research Council of Canada.

---

[1] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).

- [2] R.A. Briere *et al.* (CESR-c and CLEO-c Taskforces [CLEO-c Collaboration]), LEPP Report No. CLNS 01/1742, 2001 (unpublished).
- [3] C. Aubin *et al.*, Phys. Rev. Lett. **94**, 011601 (2005).
- [4] W.M. Yao *et al.*, J. Phys. G **33**, 1 (2006).
- [5] S. Dobbs *et al.* (CLEO Collaboration), arXiv:0712.1020 [Phys. Rev. D (to be published)].
- [6] G. Viehhauser, Nucl. Instrum. Methods Phys. Res., Sect. A **462**, 146 (2001); D. Peterson *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **478**, 142 (2002).
- [7] G.S. Huang *et al.* (CLEO Collaboration), Phys. Rev. Lett. **95**, 181801 (2005); T.E. Coan *et al.*, Phys. Rev. Lett. **95**, 181802 (2005).
- [8] S.B. Athar *et al.* (CLEO Collaboration), Phys. Rev. D **68**, 072003 (2003).
- [9] The soft muons from semimuonic decay range out before reaching the CLEO muon detectors.
- [10] R. Brun *et al.*, CERN Report No. DD/EE/84-1, 1987.
- [11] R. Barlow and C. Beeston, Comput. Phys. Commun. **77**, 219 (1993).
- [12] D.J. Lange, Nucl. Instrum. Methods Phys. Res., Sect. A **462**, 152 (2001).
- [13] D. Becirevic and A.B. Kaidalov, Phys. Lett. B **478**, 417 (2000).
- [14] T.C. Andre, Ann. Phys. (N.Y.) **322**, 2518 (2007); T. Alexopoulos *et al.*, Phys. Rev. D **71**, 012001 (2005).
- [15] S. Dobbs *et al.* (CLEO Collaboration), Phys. Rev. D **76**, 112001 (2007).
- [16] C.G. Boyd, B. Grinstein, and R.F. Lebed, Phys. Rev. Lett. **74**, 4603 (1995); Nucl. Phys. **B461**, 493 (1996).
- [17] C.G. Boyd and M.J. Savage, Phys. Rev. D **56**, 303 (1997).
- [18] C.M. Arnesen, B. Grinstein, I.Z. Rothstein, and I.W. Stewart, Phys. Rev. Lett. **95**, 071802 (2005).
- [19] T. Becher and R.J. Hill, Phys. Lett. B **633**, 61 (2006).
- [20] R.J. Hill, arXiv:hep-ph/0606023.
- [21] P. Ball, Phys. Lett. B **641**, 50 (2006).
- [22] G.S. Huang *et al.* (CLEO Collaboration), Phys. Rev. Lett. **94**, 011802 (2005).
- [23] J.M. Link *et al.* (FOCUS Collaboration), Phys. Lett. B **607**, 233 (2005).
- [24] L. Widhalm *et al.* (Belle Collaboration), Phys. Rev. Lett. **97**, 061804 (2006).
- [25] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **76**, 052005 (2007).